

Fol. 41 H

Pelagic Fish Committee

Fiskeridirektoratets  
Bibliotek

ICES CM 1997/H:8

Ref. C+G

23 DES. 1997

## REPORT OF THE

# INTERNATIONAL BOTTOM TRAWL SURVEY IN THE NORTH SEA, SKAGERRAK AND KATTEGAT IN 1996: QUARTER 1

by


The International Bottom Trawl Survey Working Group

This report is not to be quoted without prior consultation with the General Secretary. The document is a report of an expert group under the auspices of the International Council for the Exploration of the Sea and does not necessarily represent the views of the Council.

International Council for the Exploration of the Sea

Conseil International pour l'Exploration de la Mer

Palægade 2-4 DK-1261 Copenhagen K Denmark



## TABLE OF CONTENTS

Section	Page
1 INTRODUCTION .....	1
2 SURVEY METHODS AND PARTICIPATION.....	1
3 DATA AVAILABLE.....	1
4 STANDARD OUTPUT FROM THE ICES IBTS DATA BASE.....	1
5 REVISION OF THE AUTUMN SPAWNER HERRING INDEX.....	1
6 RESULTS GOV-TRAWL FOR 1996 .....	2
6.1 North Sea .....	2
6.2 Skagerrak-Kattegat (Division IIIa) .....	3
7 RESULTS OF THE SAMPLING FOR HERRING AND SPRAT LARVAE IN 1996.....	4
8 HYDROGRAPHIC DATA.....	4
8.1 Hydro-chemistry Survey .....	4
9 REFERENCES .....	5
Tables 3.1-8.1 .....	6
Figures 5.1-8.2.....	14



## **1 INTRODUCTION**

This report presents the final results for the International Bottom Trawl Survey (IBTS) in the first quarter of 1996. The survey was formerly called the International Young Fish Survey (IYFS).

In 1990 it was decided to combine the effort of the International Young Fish Survey with a number of national surveys such as the English and Scottish Groundfish Surveys into a quarterly coordinated bottom trawl survey, to be held for a period of 5 years. These quarterly surveys started in 1991. The results for quarters 2-4 will be published elsewhere.

The data in this report comprise the bottom trawl catches of the 8 standard species (herring *Clupea harengus*, sprat *Sprattus sprattus*, mackerel *Scomber scombrus*, cod *Gadus morhua*, haddock *Melanogrammus aeglefinus*, whiting *Merlangius merlangus*, saithe *Pollachius virens* and Norway pout *Trisopterus esmarkii*), as well as the catches of herring larvae. Also summarized results of temperature and salinity sampling are presented. Compared to previous reports of the IBTS Survey results saithe is from now on added to the list of standard species.

## **2 SURVEY METHODS AND PARTICIPATION**

For all matters on survey methodology, the reader is referred to the Manual (Addendum to ICES Doc. C.M.1996/H:1). Details on the participation in the 1996 survey are given below. Except for four rectangles around Shetland the whole survey area has been covered as planned.

Country	Vessel	Period	Number of Hauls	
			GOV	MIK/IKMT
Denmark	"Dana"	2-18/2	46	95
France	"Thalassa"	14-27/2	41	-
Germany, F.R.	"W. Herwig III"	20/1- 6/2	62	-
Netherlands	"Tridens"	29/1-22/2	45	46
Norway	"Michael Sars"	16/1- 5/2	41	71
Sweden	"Argos"	29/1-15/2	45	44
UK Scotland	"Scotia"	8-25/2	49	58

## **3 DATA AVAILABLE**

Table 3.1 shows for which years data are presently available in the ICES IBTS data base for the quarter 1 surveys.

At the time of the analysis of the 1996 data presented in this report all final data were available in the database.

## **4 STANDARD OUTPUT FROM THE ICES IBTS DATA BASE**

For details on the standard analysis of the data the reader is referred to a description by Pedersen (1989). At request, copies of this paper are available at the ICES Secretariat.

## **5 REVISION OF THE AUTUMN SPAWNER HERRING INDEX**

In 1994 the Herring Assessment Working Group for the Area South of 62°N has adopted a new index for 1-ringer abundance of North Sea autumn spawners. The new index is based on daytime catches in all statistical rectangles sampled during the quarter 1 survey, both in the North Sea and in the Skagerrak/Kattegat. In the calculation of this index, catches made in rectangles shallower than 10 m, or deeper than 200 m (250 m in Skagerrak), have been given less weight. The weighting factors are given in Figure 5.1.

It is implicitly assumed that all 1-ringer herring in the North Sea, Skagerrak, and Kattegat are autumn spawners. Unsampled rectangles are allocated the mean catch rate estimated within "roundfish areas" and the index is expressed as the mean catch rate (number per hour) for the entire survey area. The indices for 2+-ringers have been revised in the same way, with the exception that the catches in Skagerrak and Kattegat are assumed to be 0. This implicitly assumed that all 2+-ringers in Skagerrak and Kattegat are local or Baltic spring spawners. The use of "zero" catches instead of "missing" catches of 2+-ringers in this area is convenient because it brings the indices of all age groups on a similar scale so that for instance mortalities can be calculated directly from the indices.

The IBTS Working Group decided at the meeting in November 1995 (ICES 1996/H:1) that saithe should be added to the list of standard species. The indices of saithe for each age group are calculated in a similar way as for 1-ringer herring (see above) with the exception that also night-time hauls are used for saithe. The present report gives for the first time saithe indices and saithe charts.

## 6 RESULTS GOV-TRAWL FOR 1996

### 6.1 North Sea

Preliminary indices based on certain size classes are compared with the final indices in Table 6.1. With the exception of sprat and Norway pout, the preliminary indices are very close to the final ones. Final indices of 1- and 2- group fish of the eighth standard species are given in Table 6.2. Table 6.3 gives the mean age composition of the standard species within the relevant standard areas.

In the analysis only day-light hauls are used for herring, whereas for the other species all valid hauls are used. The number of hauls used for herring and for the other species is shown in Figures 6.1 and 6.2.

Per species a set of figures gives the distributions of the 1-, 2-, and 3+ group and the mean length of 1-group fish per rectangle. In the analysis a specific standard area for each species is used to calculate the index of year class strength. This area is indicated in the figures.

#### HERRING

Indices for 1- and 2-ringed herring are presented in Table 6.2 and mean numbers per rectangle are given in Figures 6.3 to 6.5. It should be noted that the term "age group" in herring refers to number of winter rings and not to years. All juvenile herring in the North Sea are assumed to be autumn spawners, and this means that for instance age group 1 herring in February 1996 represent year class 1994.

The index for the 1994 year class is about average for year classes 1987-1993.

#### SPRAT

Sprat indices of 1- and 2-group are given in Table 6.2. The distribution of the same age groups and the 3+ group, as well as the mean length of 1-group per rectangle, are shown in Figures 6.7-6.10.

The Herring Assessment Working Group has also for sprat adopted a new index series (ICES 1993/Assess:15) in which only hauls between 10 and 150 m depth are included. The standard area has remained the same: Division IVb only. The index for the 1995 year class is the lowest since that for the 1989 year class.

#### MACKEREL

Indices for mackerel are given in Table 6.2, the distributions of 1-, 2- and 3+ groups, and the mean length of 1-group fish are given in Figures 6.11-6.14.

Although the 1995 year class is number 4 in size since 1974 it is still low compared to the earlier periods before the collapse of the North Sea mackerel stock.

## **COD**

Abundance indices are given in Table 6.2, the distributions of 1-, 2-, and 3+ group, and the mean length of 1-group fish are given in Figures 6.15-6.18.

The 1-group index is the 4th lowest on record.

## **HADDOCK**

Abundance indices are given in Table 6.2, the distributions of 1-, 2-, and 3+ group, and the mean length of 1-group fish are given in Figures 6.19-6.22.

After a strong 1994 year class the 1995 year class is weak.

## **WHITING**

Abundance indices are given in Table 6.2, the distributions of 1-, 2-, and 3+ group, and the mean length of 1-group fish are given in Figures 6.23-6.26.

The whiting 1-group index is lower than in the most recent five years.

## **SAITHE**

Abundance indices are given in Table 6.2, the distributions of 1-, 2-, and 3+ group, and the mean length of 1-group fish are given in Figures 6.27-6.30.

Very few 1- and 2-group saithe is found in the North Sea in 1996.

## **NORWAY POUT**

Abundance indices are given in Table 6.2, the distributions of 1-, 2-, and 3+ group, and the mean length of 1-group fish are given in Figures 6.31-6.34.

The 1-group index of the 1994 year class is the second lowest on record.

## **6.2 Skagerrak-Kattegat (Division IIIa)**

The number of hauls per rectangle for herring and gadoids is shown in Figures 6.1 and 6.2.

The final indices for 1- and 2-group herring and 1- and 2+ group sprat are given in Table 6.4. The herring indices are calculated as the mean catches in four depth strata, covering the depth range of 10-150 m, and weighted by the surface area of each stratum.

Age/length keys for cod are available from the 1981 IBTS and onwards. The stocks of cod in Skagerrak and Kattegat are assessed separately and indices are given for each area. Final and preliminary indices for cod and preliminary indices for whiting and haddock are given in Table 6.5.

## **HERRING**

The indices are given in Table 6.4 and the distribution shown in Figures 6.3-6.5. The mean length of 1-ringlers is shown in Figure 6.6.

## **SPRAT**

The standard index for sprat in Division IIIa is shown in Table 6.4. The way of calculating the index has been revised by the Herring Assessment Working Group for the Area South of 62°N (ICES CM 1995/Assess:13).

## **COD**

Preliminary and final indices are given in Table 6.5. The 1-group index for the 1995 year class in the Skagerrak is lower than during the preceding last three years. The 1-group index for the Kattegat is about average. The preliminary index is about 3 times the final index.

## **HADDOCK**

The preliminary index value for haddock <20 cm (1995 year class) indicates a good year class.

## **WHITING**

The preliminary index value for whiting of <20 cm (1995 year class) indicates a moderate year class.

## **7 RESULTS OF THE SAMPLING FOR HERRING AND SPRAT LARVAE IN 1996**

During the IBTS fish larvae are sampled by towing a small meshed ring net (MIK) in oblique hauls. The catches are used in an estimation of fish larval density and abundance, assuming a 100% efficiency of the gear in catching the larvae at night.

Larval density is estimated from:

$$\text{Density (no. m}^{-2}\text{)} = (\text{no. Caught} / [\text{distance towed (m)} * \text{net-opening (m}^2\text{)}]) * \text{water depth (m)}).$$

The number of larvae within a given statistical rectangle is estimated by multiplying the density found by the surface area of a rectangle (approx.  $309 * 10^7 \text{ m}^2$ ). The total number of larvae in the sampled area (the MIK index) is the sum of estimates for all statistical rectangles.

Based on a series of comparative hauls a conversion factor between the IKMT, used in the first period of the series, and the MIK is estimated. This is used to convert earlier catches to total numbers (MIK indices).

### **Herring**

In total 314 hauls were made and this year's coverage of the sampling area was very good.

The abundance of herring larvae was low during this year's survey. The major parts of the larvae were distributed in the western part of the North Sea (Figure 7.1). This dominance of westerly sampling areas has not been seen to the same extent since 1980. The overall index value is calculated to be  $106.2 * 10^9$  (Table 7.1, Figure 7.2).

### **Sprat larvae**

Sprat larvae were sampled in small quantities in the central/southern sections of the sampling area. Because of the limited numbers, no calculation of indices will be carried out.

## **8 HYDROGRAPHIC DATA**

### **8.1 Hydro-chemistry Survey**

Seven ships contributed hydrographic data to the 1996 data set. These consist of 391 stations worked between 14 January and 27 February. Nutrient data were supplied for 125 stations, contributions being received from four ships. Data quality appeared excellent, and as a result, no stations were rejected by the data centre. Some (Denmark, Norway and Sweden) data sets were submitted immediately prior to the preparation of this report.

Charts of the distribution of bottom temperature and salinity are given in Figures 8.1 and 8.2. An updated table, giving the time series of temperature and salinity at 10 locations in the North Sea during IYFS/IBTS (1) surveys from 1970 to 1996 is provided as Table 8.1. The Figures and Table show that temperatures throughout most of the North Sea were at their lowest since at least 1988. This was undoubtedly due to the coldest winter for several years,

especially over continental Europe. The distribution of salinity was markedly similar to those apparent in 1994 and absolute levels were barely indistinguishable from the 1994 and 1995 levels.

Particularly low temperatures were present to the west of Denmark, where temperatures were less than zero for the first time since 1986. According to Table 8.1, the temperature and salinity values at location 7 in the south-eastern North Sea, were amongst the lowest ever recorded during the IYFS, presumably due to the effect of persistent easterly and south-easterly winds.

## **9 REFERENCES**

ICES 1993. Report of the Herring Assessment Working Group for the Area South of 62°N. ICES Doc. CM 1993/Assess:15.

ICES 1995. Report of the Herring Assessment Working Group for the Area South of 62°N. ICES Doc. CM 1995/Assess:13.

ICES 1996. Report of the International Bottom Trawl Survey Working Group. ICES, Doc. C.M.1996/H:1.

**Table 3.1** Data available in the ICES IBTS data base as at 27 June 1997.

**First Quarter.**

✓ = Data available  
 - = No data available  
 x = No survey made

Country	1972	1973	1974	1975	1976	1977	1978	1979	1980
Denmark	✓	✓	✓	✓	✓	✓	✓	✓	x
France	x	x	x	x	✓	x	x	✓	✓
Germany	-	-	-	-	-	-	-	-	-
Netherlands	✓	✓	✓	✓	✓	✓	✓	✓	✓
Norway	-	-	-	-	-	-	-	-	-
Sweden	-	-	-	-	-	-	-	-	-
UK England	-	-	-	-	-	-	-	-	-
UK Scotland	x	x	✓	✓	✓	✓	✓	✓	✓
USSR	x	x	✓	✓	✓	✓	x	✓	x

Country	1981	1982	1983	1984	1985	1986	1987	1988	1989
Denmark	x	✓	✓	✓	✓	✓	✓	✓	✓
France	x	✓	✓	✓	✓	✓	✓	✓	✓
Germany	-	-	✓	✓	✓	✓	✓	✓	✓
Netherlands	✓	✓	✓	✓	✓	✓	✓	✓	✓
Norway	-	-	✓	✓	✓	✓	✓	✓	✓
Sweden	-	-	✓	✓	✓	✓	✓	✓	✓
UK England	✓	✓	✓	✓	✓	✓	✓	✓	✓
UK Scotland	✓	✓	✓	✓	✓	✓	✓	✓	✓
USSR	✓	✓	x	x	x	x	x	x	x

Country	1990	1991	1992	1993	1994	1995	1996	1997
Denmark	✓	✓	✓	✓	✓	✓	✓	✓
France	✓	✓	✓	✓	✓	✓	✓	✓
Germany	✓	✓	✓	✓	✓	✓	✓	✓
Netherlands	✓	✓	✓	✓	✓	✓	✓	✓
Norway	✓	✓	✓	✓	✓	✓	✓	✓
Sweden	✓	✓	✓	✓	✓	✓	✓	✓
UK England	✓	x	x	x	x	x	x	x
UK Scotland	✓	✓	✓	✓	✓	✓	✓	✓
USSR	x	x	x	x	x	x	x	x

**Table 6.1** Preliminary indices for 1-group fish based on a split of the length distribution, and final indices for the North Sea from the first quarter IBTS in 1996.  
For herring the “old” standard area is used.

	Preliminary	Final
Herring	2,145	2,400.2
Sprat	533	232.6
Mackerel	12.8	10.4
Cod	4.4	3.5
Haddock	304	267.4
Whiting	573	502.4
Saithe	-	0
Norway pout	2,148	922.9

**Table 6.2** IBTS indices for 1- and 2-year-old fish of various species. Indices in mean number per hour within the relevant standard area in the North Sea.

Year class	HERRING	SPRAT	MACKEREL	COD	HADDOCK	WHITING	SAITHE	NORWAY POUT						
	1-ring	2-ring	age 1	age 2	age 1	age 2	age 1	age 2	age 1	age 2	age 1	age 2	age 1	age 2
1969	-	-	-	-	-	-	25.9	-	32	-	31	-	-	-
1970	-	-	-	-	-	-	98.3	34.5	855	299	274	190	-	-
1971	-	-	90	-	-	-	4.1	10.6	740	971	332	763	-	-
1972	-	-	123	-	-	-	38.0	9.5	187	110	1,156	496	-	-
1973	-	-	481	-	-	0.1	14.7	6.2	1,092	385	322	153	-	-
1974	-	-	-	-	16.5	0.2	40.3	19.9	1,168	670	893	535	-	4,242
1975	-	-	1,186	-	0.4	+	7.9	3.2	177	84	679	219	-	334
1976	-	-	136	-	1.4	+	36.7	29.3	162	108	418	293	-	4,813
1977	172*	-	1,474	-	2.3	+	12.9	9.3	385	240	513	183	-	1,913
1978	312*	-	248	-	0.2	+	9.9	14.8	480	402	457	391	-	2,690
1979	431*	-	1,402	1,380	+	+	16.9	25.5	896	675	692	485	-	4,081
1980	772*	109	941	502	0.1	0.1	2.9	6.7	268	252	227	232	-	1,375
1981	1,260	161	296	754	0.1	5.2	9.2	16.6	526	400	161	126	-	663
1982	1,443	716	210	387	1.9	0.4	3.9	8.0	307	219	128	179	-	802
1983	2,083	661	382	298	0.1	0.0	15.2	17.6	1,057	828	436	359	-	3,925
1984	2,542	838	660	103	0.7	2.1	0.9	3.6	229	244	341	261	-	384
1985	3,684	4,103	71	74	0.5	+	17.0	28.8	579	326	456	544	-	2,043
1986	4,530	775	803	1,437	8.9	0.1	8.8	6.1	885	688	669	862	-	760
1987	2,313	580	148	442	1.2	1.8	3.6	6.3	92	97	394	542	-	127
1988	1,016	794	4,246	557	1.1	1.2	13.1	15.2	210	114	1,465	887	-	773
1989	1,159	377	177	116	35.0	0.2	3.4	4.1	219	131	509	675	-	677
1990	1,162	762	1,121	340	6.9	0.4	2.4	4.5	679	371	1,014	748	-	902
1991	2,943	1,094	1,561	422	16.0	0.8	13.0	19.9	1,115	543	916	524	-	2,644
1992	1,667	1,285	1,755	1,368	1.0	0.1	12.7	4.4	1,242	504	1,087	637	-	375
1993	1,186	194	4,003	2,716	2.2	+	14.8	22.1	229	205	721	457	-	785
1994	1,735		1,138	558	+	0.7	9.7	8.0	1,375	813	679	486	-	5,941
1995			233		10.4		3.5		267		502	0		923

\* Revised according to new procedures by the Herring Assessment Working Group for the Area South of 62°N (1995/Assess:13), but not based on retrieval from database.

**Table 6.3** Age composition of the standard species in 1996 for the relevant standard areas in the North Sea.

Age group	1	2	3	4	5	6+
Herring	1,734.8	194.3	43.3	12.6	8.5 <sup>1</sup>	-
Sprat	232.6	557.5	176.9	22.4	1.1 <sup>1</sup>	-
Mackerel	10.4	0.7	1.2	0.2	0.2	0.1
Cod	3.5	8.0	6.0	0.7	0.6	0.4
Haddock	267.4	813.3	65.9	46.7	7.7	3.1
Whiting	502.4	486.0	244.7	69.7	23.1	9.8
Saithe	0.5	0.6	1.8	18.1	1.3	0.3
Norway pout	922.9	2,631.0	227.8	4.7	0.1	0.0

<sup>1</sup>Plus-group.

**Table 6.4** IBTS indices for herring and sprat in Skagerrak-Kattegat. The herring indices are weighted by the area of four depth strata and the sprat indices are the mean by hour within the total area (n.a. = not available).

Year	Herring						Sprat		
	Total		Spring-spawners		Autumn-spawners		1-group	2+group	Total
	1-ring	2-ring	1-ring	2-ring	1-ring	2-ring			
1974							1,325		
1975							5,339		
1976							2,069		
1977							5,713	984	6,697
1978							5,119	2,117	7,236
1979							3,338	1,482	4,820
1980	2,311	387	1,607	307	704	80	4,960	3,592	8,558
1981	3,246	1,393	966	1,318	2,250	75	2,809	3,068	5,877
1982	2,560	549	1,408	445	1,152	104	1,577	4,695	6,272
1983	5,419	1,063	1,522	946	3,897	117	1,173	1,685	2,858
1984	6,035	1,947	2,793	1,419	3,242	528	5,818**	861	7,034
1985	7,994	2,473	-*	1,867	-*	606	2,404	2,426	5,388
1986	21,489	2,738	-*	1,562	-*	1,176	670	1,934	4,545
1987	11,733	3,671	-*	2,921	-*	949	2,234	2,219	8,048
1988	67,753	10,095	-*	7,834	-*	2,161	950	5,527	10,634
1989	17,451	4,976	-*	0	-*	4,976	435	1,012	3,310
1990	3,544	3,876	0	3,192	3,544	684	510	243	944
1991	3,588	3,749	-*	480	-*	3,269	659	468	1,945
1992	5,057	1,934	0	771	5,057	1,163	5,897	634	7,122
1993	26,738	3,165	0	203	26,738	2,962	1,593	4,237	7,186
1994	8,777	2,333	0	0	8,777	2,333	1,494	586	3,361
1995	7,114	535	0	0	7,114	535	2,006	1,726	3,802
1996	n.a.	n.a.	n.a.	n.a.	n.a.	n.a.	255	5,099	5,312

\* Separation not valid.

\*\* From 1984 onwards only hauls taken in depths between 10 and 150 m are included in the estimates

**Table 6.5** Indices of 1- and 2-group cod and 1-group whiting and haddock in Skagerrak-Kattegat.  
The preliminary indices are the mean number per hour for cod < 25 cm,  
whiting and haddock < 20cm.


Year Class	Cod 1-group Skagerrak		Cod 1-group Kattegat		Cod 2-group		Whiting 1-group Prel.	Haddock 1-group Prel.
	Prel.	Final	Prel.	Final	Skagerrak	Kattegat		
1974							499	
1975							236	
1976							99	
1977							392	
1978							561	
1979	79		386		93	171	722	40
1980	18	15	42	26	31	63	968	4
1981	36	36	126	104	30	258	690	48
1982	32	28	113	96	19	143	262	34
1983	24	23	49	39	52	106	500	72
1984	18	14	18	9	10	72	940	161
1985	82	78	229	213	113	372	1379	57
1986	15	5	48	11	18	28	2178	251
1987	81	77	76	68	24	48	2978	125
1988	62	56	6	3	8	16	478	20
1989	25	31	131	153	25	112	2255	8
1990	42	9	39	19	5	20	1636	74
1991	119	96	64	64	16	38	1796	288
1992	171	110	40	31	7	13	1359	405
1993	77	82	22	19	47	114	525	180
1994	111	97	390	300	35	272	1105	233
1995	37	36	172	52			883	490

**Table 7.1** Density and abundance of herring larvae caught in February during the IBTS. Values for year classes by areas are density estimates in number per square metre. Total abundance is found by multiplying density by surface area and summing up.

Area	North west	North east	Central west	Central east	South west	South east	Division IIIa	Southern Bight	O-ringrs abundance
Area m <sup>2</sup> x 10 <sup>9</sup>	83	34	86	102	37	93	31	31	N 10 <sup>9</sup>
<b>Year class</b>									
1976	0.054	0.014	0.122	0.005	0.008	0.002	0.002	0.016	17.1
1977	0.024	0.024	0.050	0.015	0.056	0.013	0.006	0.034	13.1
1978	0.176	0.031	0.061	0.020	0.010	0.005	0.074	0.000	52.1
1979	0.061	0.195	0.262	0.408	0.226	0.143	0.099	0.053	101.1
1980	0.052	0.001	0.145	0.115	0.089	0.339	0.248	0.187	76.7
1981	0.197	0.000	0.289	0.199	0.215	0.645	0.109	0.036	133.9
1982	0.025	0.011	0.068	0.248	0.290	0.309	0.470	0.140	91.8
1983	0.019	0.007	0.114	0.268	0.271	0.473	0.339	0.377	115.0
1984	0.083	0.019	0.303	0.259	0.996	0.718	0.277	0.298	181.3
1985	0.116	0.057	0.421	0.344	0.464	0.777	0.085	0.084	177.4
1986	0.317	0.029	0.730	0.557	0.830	0.933	0.048	0.244	270.9
1987	0.078	0.031	0.417	0.314	0.159	0.618	0.483	0.495	168.9
1988	0.036	0.020	0.095	0.096	0.151	0.411	0.181	0.016	71.4
1989	0.083	0.030	0.040	0.094	0.013	0.035	0.041	0.000	25.9
1990	0.075	0.053	0.202	0.158	0.121	0.198	0.086	0.196	69.9
1991	0.255	0.390	0.431	0.539	1.500	0.369	0.298	0.395	200.7
1992	0.168	0.039	0.672	0.444	0.734	0.268	0.345	0.285	190.1
1993	0.358	0.212	0.260	0.187	0.120	0.119	0.223	0.028	101.7
1994	0.148	0.024	0.417	0.381	0.332	0.148	0.252	0.169	126.9
1995	0.260	0.086	0.699	0.092	0.266	0.018	0.001	0.020	106.2


**Table 8.1****Time series data of bottom temperature and salinity during IYFS/IBTS(1) 1970-1996**

Location	1		2		3		4		5		6		7		8		9		10	
Position	60° 0'N 2°E		57° 30'N 0°E		57° 30'N 2°E		57° 30'N 4°E		55° 0'N 0°E		55° 0'N 2°E		55° 0'N 4°E		55° 0'N 8°E		54° 0'N 3°E		52° 30'N 3°E	
Year	t°C	S-30	t°C	S-30	t°C	S-30	t°C	S-30	t°C	S-30	t°C	S-30	t°C	S-30	t°C	S-30	t°C	S-30	t°C	S-30
1970	5.5	5.08	5.8	4.95	5.3	5.00	4.7	4.92	5.9	4.75	4.5	4.82	4.0	4.72	0.5	3.00	4.0	4.72	4.0	4.62
1971	7.1	5.15	7.0	5.05	6.9	5.15	6.0	5.10	7.0	4.82	6.2	4.88	5.5	4.80	3.5	3.00	5.9	4.55	7.0	4.95
1972	5.8	5.22	6.9	5.08	5.9	5.20	4.5	4.78	6.5	4.91	4.8	4.86	5.2	4.80	2.5	3.80	5.2	4.70	6.9	5.10
1973			7.4	5.02	7.2	5.20	6.7	5.10	7.0	5.05	6.1	5.00	6.0	4.86	5.0	3.00	6.4	4.80	6.5	5.05
1974	6.9	5.28	6.5	5.11	6.5	5.08	6.3	5.04	6.5	4.90	6.0	4.90	5.6	4.90	4.7	3.00	6.1	4.78	8.0	5.20
1975	7.3	5.20	6.6	5.05	6.6	5.15	6.4	5.13	6.6	4.95	6.4	4.90	6.1	4.85	5.2	3.50	5.9	4.62	6.9	4.62
1976	6.7	5.20	6.5	5.00	6.5	5.15	5.6	5.12	6.1	4.81	4.9	4.95	4.9	4.85	2.2	1.00	5.1	4.78	5.1	4.80
1977	6.0	5.18	6.2	5.02	5.1	5.00	4.8	4.92	6.0	4.98	4.9	4.85	5.0	4.80	3.1	3.60	5.6	4.78	7.1	5.22
1978	6.4	4.88	6.6	5.00	6.0	4.90	4.7	4.88	5.6	4.78	4.9	4.88	4.2	4.80	2.2	2.50	4.6	4.68	5.5	4.90
1979	6.4	5.15	6.0	4.80	4.1	4.88	4.0	4.98	4.5	4.64	2.8	4.62	2.8	4.62	-1.5	2.00	3.0	4.62	4.2	4.95
1980	5.9	5.12	6.6	5.00	5.5	5.00	4.5	4.70	6.1	4.60	3.8	4.65	4.5	4.50	3.1	3.50	5.1	4.70	6.1	5.11
1981	6.9	5.22	6.6	4.90	6.2	5.05	5.8	5.15	6.5	4.80	5.8	4.82	5.1	4.82	3.4	2.50				
1982	6.6	5.28	6.1	5.02	5.9	5.05	5.5	5.10	5.5	4.72	4.8	4.82	4.5	4.62	2.8	2.50	4.7	4.30	6.0	4.65
1983	6.9	5.22	6.5	5.00	6.4	5.10	6.2	5.15	5.6	4.62	6.1	4.95	5.2	4.90	3.0	3.00	5.2	4.80	6.4	4.70
1984	6.3	5.18	6.4	5.10	6.4	5.10	5.2	5.12	5.9	4.80	5.0	4.84	4.9	4.90	3.5	3.00	4.9	4.65	7.4	4.95
1985	6.9	5.17	6.8	5.10	6.5	5.18	5.9	5.05	6.5	4.70	4.7	4.91	5.0	4.90	1.0	2.50	4.0	4.70	6.0	4.80
1986	6.6	5.25	5.8	5.05	5.4	5.08	5.2	5.05	5.2	4.65	3.9	4.72	3.6	4.60	0.0	2.50	4.0	4.60	4.0	4.65
1987	6.5	5.28	6.1	4.90	5.9	5.08	4.9	5.00	5.0	4.75	4.2	4.80	4.3	4.60	0.8	0.00	4.9	4.60	4.8	4.90
1988	7.6	5.18	7.6	4.95	7.4	5.03	7.0	4.96	7.1	4.70	6.6	4.80	6.5	4.50	5.9	3.50	6.9	4.60	7.7	4.90
1989	8.5	5.29	8.0	4.85	7.8	4.89	7.6	5.05	7.5	4.76	7.1	4.81	6.8	4.80	6.0	4.10	6.5	4.68	7.5	4.62
1990	8.5	5.29	7.6	5.00	7.6	5.12	7.6	5.15	7.5	4.70	7.5	4.85	7.5	4.80	6.5	4.10	7.4	4.70	7.4	4.60
1991	7.9	5.30	6.7	5.10	7.1	5.22	6.1	4.97	6.6	4.65	5.8	4.85	5.5	4.80	3.0	4.00	5.8	4.60	6.1	5.30
1992	8.1	5.29	7.6	5.10	7.1	5.16	7.1	5.19	7.4	4.80	6.6	4.80	6.5	4.80	6.6	2.00	4.5	4.80	6.0	5.20
1993	7.4	5.31	6.5	4.92	6.4	5.18	6.5	5.30	6.5	5.05	6.2	5.00	5.4	4.95	4.3	3.50	5.6	4.80	6.0	5.00
1994	6.2	5.20	6.5	5.05	5.5	4.93	4.3	4.80	6.3	4.90	5.4	4.90	5.2	4.80	4.0	2.00	5.5	4.70	7.0	5.00
1995	7.5	5.23	7.0	4.92	7.1	5.00	6.7	5.09	6.7	4.71	6.0	4.87	5.6	4.81	4.0	0.03	6.0	4.65	7.9	4.51
1996	7.1	5.24	6.5	4.91	5.0	4.94	4.7	4.87	6.0	4.59	4.6	4.71	3.0	4.44	-0.2	2.12	3.4	4.71	3.8	4.83



**Figure 5.1** Weighting factors used in the revised IBTS herring index and for saithe. Numbers denote the relative weight of a given rectangle. Only weighting factors less than 1 are shown.


International Bottom Trawl Survey 1996Q1



Herring, CLUP HAR  
Number of Hauls.

Figure 6.1 Number of hauls used in the analysis of herring 1996 data.


# International Bottom Trawl Survey 1996Q1



Saithe, POLL VIR  
Number of Hauls.


**Figure 6.2** Number of hauls used in the analysis of herring 1996 data on sprat, mackerel, cod, haddock, saithe, whiting and Norway pout.

International Bottom Trawl Survey 1996Q1



Herring, CLUP HAR  
Number per Hour , Age Group 1.


International Bottom Trawl Survey 1996Q1



Herring, CLUP HAR  
Number per Hour , Age Group 2.

Figure 6.4 Herring: number per hour, 2-ringers.


International Bottom Trawl Survey 1996Q1



Herring, CLUP HAR  
Number per Hour , Age Group 3+.

**Figure 6.5** Herring: number per hour, 3+ringers.

# International Bottom Trawl Survey 1996Q1




Herring, CLUP HAR  
Mean Length, Age Group 1.

Figure 6.6

Herring: mean length, age 1.

# International Bottom Trawl Survey 1996Q1




Sprat, SPRA SPR  
Number per Hour , Age Group 1.

**Figure 6.7**

Sprat: number per hour, age-group 1.


# International Bottom Trawl Survey 1996Q1



Sprat, SPRA SPR  
Number per hour , Age Group 2.

Figure 6.8 Sprat: number per hour, age-group 2.


# International Bottom Trawl Survey 1996Q1



Sprat, SPRA SPR  
Number per Hour , Age Group 3+.

**Figure 6.9** Sprat: number per hour, age-group 3+.


# International Bottom Trawl Survey 1996Q1



Sprat, SPRA SPR  
Mean Length, Age Group 1.


Figure 6.10 Sprat: mean length, age-group 1.

International Bottom Trawl Survey 1996Q1




Mackerel, SCOM SCO  
Number per Hour , Age Group 1.

# International Bottom Trawl Survey 1996Q1



Mackerel, SCOM SCO  
Number per Hour , Age Group 2.


International Bottom Trawl Survey 1996Q1



Mackerel, SCOM SCO  
Number per Hour , Age Group 3+.

Figure 6.13 Mackerel: number per hour, age 3+.


# International Bottom Trawl Survey 1996Q1



Mackerel, SCOM SCO  
Mean Length, Age Group 1.


Figure 6.14 Mackerel: mean length, age 1.

# International Bottom Trawl Survey 1996Q1



Cod, GADU MOR  
Number per Hour , Age Group 1.


International Bottom Trawl Survey 1996Q1



Cod, GADU MOR  
Number per Hour , Age Group 2.

Figure 6.16 Cod: number per hour, age group 2.

# International Bottom Trawl Survey 1996Q1




Cod, GADU MOR

Number per Hour , Age Group 3+.

Figure 6.17 Cod: number per hour, age group 3+.


International Bottom Trawl Survey 1996Q1



Cod, GADU MOR  
Mean Length, Age Group 1.

**Figure 6.18** Cod: mean length, age group 1.


# International Bottom Trawl Survey 1996Q1



Haddock, MELA AEG  
Number per Hour , Age Group 1.

Figure 6.19 Haddock: number per hour, age group 1.


# International Bottom Trawl Survey 1996Q1



Haddock, MELA AEG  
Number per Hour , Age Group 2.


Figure 6.20 Haddock: number per hour, age group 2.

International Bottom Trawl Survey 1996Q1




Haddock, MELA AEG  
Number per hour, Age Group 3+.

International Bottom Trawl Survey 1996Q1



Haddock, MELA AEG  
Mean Length, Age Group 1.


# International Bottom Trawl Survey 1996Q1



Whiting, MERL MNG  
Number per Hour , Age Group 1.

Figure 6.23 Whiting: number per hour, age group 1.


# International Bottom Trawl Survey 1996Q1



Whiting, MERL MNG  
Number per Hour , Age Group 2.

Figure 6.24 Whiting: number per hour, age group 2.


# International Bottom Trawl Survey 1996Q1



Whiting, MERL MNG  
Number per Hour , Age Group 3+.

Figure 6.25 Whiting: number per hour, age group 3+.


# International Bottom Trawl Survey 1996Q1



Whiting, MERL MNG  
Mean Length, Age Group 1.


**Figure 6.26** Whiting:mean length, age group 1.

International Bottom Trawl Survey 1996Q1




Saithe, POLL VIR  
Number per Hour , Age Group 1.

# International Bottom Trawl Survey 1996Q1



Saithe, POLL VIR  
Number per Hour , Age Group 2.

International Bottom Trawl Survey 1996Q1




Saithe, POLL VIR  
Number per Hour , Age Group 3+.

Figure 6.29

Saithe: number per hour, age 3+.


International Bottom Trawl Survey 1996Q1



Saithe, POLL VIR  
Mean Length, Age Group 1.

**Figure 6.30** Saithe: mean length, age 1.


# International Bottom Trawl Survey 1996Q1



Nor. Pout, TRIS ESM  
Number per Hour , Age Group 1.

Figure 6.31 Norway pout: number per hour, age group 1. 45


# International Bottom Trawl Survey 1996Q1



Nor. Pout, TRIS ESM  
Number per Hour , Age Group 2.

Figure 6.32 Norway pout: number per hour, age group 2.


# International Bottom Trawl Survey 1996Q1



Nor. Pout, TRIS ESM  
Number per Hour , Age Group 3+.

**Figure 6.33** Norway pout: number per hour, age group 3+.


# International Bottom Trawl Survey 1996Q1




Nor. Pout, TRIS ESM  
Mean Length, Age Group 1.

Figure 6.34 Norway pout:mean length, age group 1.


0-ringers year class 1993



0-ringers year class 1994




0-ringers year class 1995


**Figure 7.1**

Distribution of 0-ringer herring, year classes 1993-1995. Abundance estimates of 0-ringers within each statistical rectangle based on MIK catches during IBTS in February. Areas of filled circles illustrate densities in  $\text{no m}^{-2}$ , the area of a circle extending to the border of a rectangle represents  $1.5 \text{ m}^{-2}$ .


### Time series of recruitment indices



**Figure 7.2** Trend in MIK 0-ringer and IBTS 1-ringer indices for the year classes 1977-1995



**Figure 8.1**



**Figure 8.2**



