

LONG TERM CHANGES OF GROWTH AND MATURATION  
IN THE NORWEGIAN SPRING SPAWNING HERRING

by

Reidar Toresen

Institute of Marine Research  
P.O. Box 1870 Nordnes  
N-5024 Bergen, Norway

ABSTRACT

Two periods with great differences in year class abundance and recruitment were chosen for growth and maturation studies, 1950 - 1960 and 1973 - 1983. In the first period, the mean year class strength at 3 years was  $6.8 \times 10^9$  individuals while in the second period it was  $0.4 \times 10^9$  ind.. The growth in each period as observed by the length at age, is compared to the year class strength and the abundance of the stock at the time. Variations in growth are observed and before maturation strong year classes has remarkably less growth than weak year classes. The latter period is therefore characterized by far better growth, especially in the early years of life, than the former. According to growth rates of the various year classes, the age at maturity is substantially higher in the first period than in the latter. However, the length at maturity is found to be fairly constant showing the maturation to be size dependent rather than dependent on the age.

## INTRODUCTION

After the great reduction in the abundance of the Norwegian spring spawning herring in the late sixties, the growth as observed by the length at age and weight at age increased significantly, (Jørgensen 1979). At the same time, the adult stock totally changed its migration pattern as it ceased migrating into the Norwegian Sea after spawning, (Dragesund et al. 1980). Since 1970 the distribution has been limited to the fjords of the western and northern Norway (except for the 1983 year-class). The observed change in growth and maturation is explained by the severe reduction of the abundance of the stock as observed in other herring stocks, (Jakobsson and Halldorsson 1984). However, this change in growth is not necessarily density dependent.




Fig. 1. The areas in which growth is studied.

It should be noted that growth in the Norwegian spring spawners is extremely variable, depending strongly on the area in which they spend their adolescent phase. Herring located off Finnmark and in the Barents Sea during the adolescent phase have a markedly slower growth

than those found off the coast between Møre and Lofoten. As the maturing process is dependent on the length rather than the age in the herring (Toresen 1986), the parts of the stock spending their first years of life south of Lofoten mature earlier than those found further to the north or in the Barents Sea. Accordingly, immature herring older than three years are scarce south of Lofoten, whereas off Finnmark herring in the adolescent stage are common up to four years of age. This pattern is probably independent of the size of the year-classes. What is most certainly dependent on the year class strength is the relative abundance of the stock in the various areas and the growth within these areas.

Since the recruitment to this stock is so variable, it is of great importance in the assessment that the prediction of growth is correct, and it is therefore valuable to know the mechanisms that govern the growth.

In this paper various aspects of growth in the Norwegian spring spawners are enlightened especially with respect to the question of growth and year class abundance. Two periods with quite significant differences in abundance, recruitment and growth are chosen for this purpose.

#### MATERIAL AND METHODS

The immature part of the stock is treated separately from the adults because the biological samples from the two parts of the stock are not taken in the same period of the year. The samples from the spawning stock are taken during the spawning in spring while the samples from the immatures are taken during autumn.

##### The immatures

Growth, as expressed by the mean length at age in the autumn (Sept-Nov) is studied within 4 regions along the Norwegian coast (Fig. 1). Average values of length at age for the full periods are compared between the regions. The 1983 year class is treated separately because it is a very strong one showing a tendency to a level of growth which is similar to that of the fifties.

The material from the fifties is very poor, as no systematic sampling of the juvenile part of the stock was carried out at that time. No weight at age data are available.

##### The adults

The geographical regions are here cut down to one, the main spawning grounds in area 07 (Fig 1). Sampling is done during spawning in spring (Feb-Apr). Mean values of length at age each year in the two periods are held up against spawning stock abundance. In addition, the mean length at age for each year class is compared to the year class strength at three years. Further on, growth as expressed by length increments in the age intervall from 5-10 years is compared between the periods and the mean length at age for all year classes in each period are compared in a plot.

At last, the growth as expressed by the weight at age relationship is discussed briefly.

## Maturation

Age at maturity. The herring stock in the seventies is divided in two components, one northern type and one southern type. These components were characterized by differences in the growth pattern, as the southern one was more fast growing. Maturity ogives are calculated for both components separately.

When a year class is fully recruited to the spawning stock the ratio between the number of individuals in this year class and the number in the older year classes is constant if total mortality (Z) by age group is constant.

In the spawning stock, the year classes mix very well and in the samples taken from the spawning stock the ratio between the various year classes is quite constant although it may vary somewhat from one year to another. However, no trend in this variance was detected and the proportions of each year class in the spawning stock was calculated as follows:

$$p_i = \frac{N_i}{\sum_{i=1}^k N_i}$$

where  $p_i$  is the proportion of year-class  $i$  in the spawning stock,  $N_i$  is the number of fish in the samples from this year-class and  $k$  is the number of year-classes in the spawning stock that is fully recruited. The relative abundance of the fully recruited year-classes were then calculated as the mean of the age groups 6, 7 and 8 years ( $p_i$ ). The maturity ogive by year-class is then calculated as the proportions of the number of fish in the younger age groups to  $p_i$  in the samples of the spawning stock.

## RESULTS AND DISCUSSIONS

## The immatures

The observed data of length at age for the two periods in four areas along the Norwegian coast are shown in Table 1 a-d. The data given are mean values of all samples of young herring from September till November each year. The mean values of length at age might have been affected by the time the samples are taken in the autumn. However, the growth during these months is marginal and it is assumed that the growth for the season is finished. As can be seen from the tables, the data from the fifties and early sixties is very poor. However, treated together, the few samples present give an idea of the growth level in the full period.

Comparing the length at age within the two periods but between the different regions show a clear trend of increasing growth when moving from north to south along the coast (Table 1 a-d). This phenomenon is earlier described by Dragesund (1970), and is probably related to environmental conditions. The length distribution of the 0-group herring may be similar all along the coast, but a difference in length at age is initiated very soon and this difference increases through the years. This feature results in earlier maturation of the components of the stock spending the adolescence further to the south




Fig. 2a-d. Plots of length at age in the four regions in the two periods.

than those found in the north or in the Barents Sea, as the maturation process is related to growth rather than to age. The recruitment to the spawning stock is therefore largely dependent on the geographical distribution of the year classes in the early stages of life.

The difference in growth between the two periods are studied by comparing the average length at age values for each period and region. Figure 2. a-d shows the plots of length at age in the four regions in each period. The 1983 year-class is plotted separately. In all the regions, growth is far better in the seventies than in the fifties. As the plots clearly show, the growth rate is also higher in this latter period leading to greater differences in length at age through the years. This higher growth also leads to maturation at a much younger age for the year classes in the seventies, which was observed down to 3 years.

Is this large difference in growth between these two periods density dependent? The annual recruitment was at a much higher level in the fifties than in the seventies. The mean year class strength at 3 years was  $6\ 790 \times 10^6$  individuals in the fifties, while in the seventies this number was reduced to  $436 \times 10^6$ . However, it is a fact that big year classes of Norwegian spring-spawning herring have quite another geographical distribution than small ones. Weak year classes spend more or less entirely the adolescence in the fjords and along the coast. The relative distribution of each year class in the different regions is determined at the 0-group stage. Abundant year classes also distribute in the fjords, but at a certain level of abundance, the 0-group herring also get distributed in the open watermasses of the Barents Sea. The mechanism governing this distribution pattern is not yet known. However, the conditions for high growth rates are less favourable in the open watermasses of the oceans than in the fjords and along the coast, where the temperature is higher and the density of zooplankton is greater. The density of herring in the fjords for these big year classes is still much higher than for small year classes, probably leading to reduced growth due to higher density. In this way, we have two main reasons for reduced growth in big year classes of Norwegian spring-spawners; density dependent growth in the fjords and reduced growth due to distribution in the open watermasses of the Barents Sea. The latter factor is a secondary effect of high density but the reduced growth is not density dependent itself.

#### Maturation

The results are shown in Table 2. It is quite clear that the recruitment to the spawning stock set in earlier in the seventies than in the fifties and sixties. Especially the year classes, 1973 and 1974 mature to spawn at a very early age and they are almost fully recruited at four. For the year classes 1959-61, they partly mature at three, but only to a very small extent. Full recruitment doesn't occur until they are six or seven.

These maturity ogives are strongly related to the growth. Fast growing herring mature to spawn early, but more or less at the same length as the slow growing ones.





Fig. 3a & b. Mean length at age per year and spawning stock abundance.

## The adults

Length at age. The observed data of mean length at age for the two periods at the spawning grounds in Area 07 is shown in Table 3. All samples taken in the actual time, February - April, are present in the material.

Mean length at age per year in the two periods and spawning stock abundance is shown in Fig. 3, a and b. The spawning stock decreases from a level of about 10 mill tonnes in 1957 till about 3 mill tonnes in 1965 and further to nearly zero in 1969. Through these years, the growth, as expressed by the mean length at age per year, show an increasing trend throughout the fifties. In the beginning of the sixties however, the growth suddenly start to fall shown by the low values of mean length at age for the year classes 1958-59 and -60. The correlation between this pattern of growth and the stock abundance is not very clear because the change of growth pattern differs from one age group to another. In fact, it is obvious that the growth pattern changes in correspondance with the year class strength rather than the stock abundance.




Fig. 4. Mean length at age per year class strength at three years in the two periods.

Figure 4 shows the mean length at age per year class and the corresponding year class strength in the two periods. It is clear, from the plot that there is a positive correlation between the length at age for the different year classes and the year class abundance. The overall growth pattern shows an increasing trend until 1957. The 1958 year class has far less growth than the two or three older year classes and this decreasing tendency to growth continues for the 1959 year class flattening out for the 1960 year class. This pattern is set before the herring mature. As shown in Fig. 4, the pattern is clear at



4 years, smoothening a little until the age of ten. The observed drop in growth for the weak 1958 year class is most certainly caused by interactions between this year class and the very strong 1959 year class.

In the second period, only weak year classes occur, except for the 1983 year class which is comparable to strong year classes in the fifties. It is difficult to draw any conclusions from the appearing growth pattern. However there is a tendency to decreasing values of length at age throughout the years (Fig. 3 b). This decrease might have been affected by the increase in the abundance of the total stock. However, a clear drop in growth is observed for the strong 1983 year class.

Comparing the curves of mean length at age by year class between the two periods in Figure 4 gives an impression of better growth in the seventies than in the fifties. However, the better growth is restricted to the immature part of the stock only, as the mean length increments for the age intervall 5-10 years is higher in the former period than in the latter, 5.87 and 5.08 cm respectively. The length increment in the age intervall 5-10 years for the year classes with lowest values of length at 4 years is much higher than for year-classes with high values of mean length at 4. This phenomenon is earlier described by Watkin (1927), and Østvedt (1958) and Jakobsson & Halldorsson (1984) has described this for Atlanto-scandian herring. It leads to less variation in length within the same year class as the fish grow older.

Figure 5 shows the plots of mean length at age for all year year classes in the two periods. The plot illustrates the higher values of mean length at age in the seventies than in the fifties and the difference is constant in the whole age interval 5-10 years. The Figures 2 and 5 compared show very well that the differences in growth in length is initiated early in life, before maturation.




Fig. 5. Mean length at age for all year classes in the two periods.

Weight at age. Table 2 shows the mean weight at age each year for the two periods. These data show exactly the same pattern of growth as the growth of length data did (Fig. 4 and 5). There is a more or less constant difference of 35-60 g between the mean weight at age values for the age groups in the two periods. These differences follows the length at age differences and varies randomly during the age intervall 5-10 years.

#### MANAGEMENT CONSIDERATIONS

Without any doubt, there is a relationship between year class abundance and growth. The reduced growth due to high abundance might be density dependent in certain areas, but in most cases, it is most certainly due to dispersion of young herring in areas with lower concentrations of food particles and lower temperatures.

The reduced growth leads to later maturation because the maturation is size dependent rather than age dependent. The recruitment of large year classes to the spawning stock therefore tend to be delayed compared to the recruitment of small year classes. This delay may be 2-3 years or even more. The recruitment to the fishery will also be later for larger year classes than for less abundant year classes since the fishery is regulated by a minimum landing size regulation. If the prediction of growth is not thoroughly evaluated in the assessment work, the delay in recruitment may lead to an over-exploitation of the older year classes in the stock. Since the TAC is given in weight rather than number of individuals, a reduced growth also may lead to a general over-exploitation of the stock.

#### REFERENCES

- Dragesund, O. 1970. Distribution, abundance and mortality of young and adolescent Norwegian spring spawning herring (Clupea harengus L.) in relation to subsequent year-class strength. Fisk. Dir. Skr. Ser. Havunders., 15(4): 451 - 556.
- Dragesund, O., Hamre, J. and Ulltang, Ø. 1979. Biology and population dynamics of the Norwegian spring spawning herring. ICES Symposium on the biological basis of pelagic fish stock management, 3: 80 pp. (mimeo).
- Jakobsson, J. and Halldorsson, O. 1984. Changes in biological parameters in the Icelandic summer spawning herring. ICES CM 1984/H:43, 12 pp. (mimeo).
- Jørgensen, K. O. 1979. Norsk vårgytende sild (Clupea harengus L.) Endringer i biologiske parametre i perioden 1950 - 1978. Thesis, University of Bergen, 75 pp. (In Norwegian) (mimeo).

Toresen, R. 1986. Length and age at maturity of Norwegian spring-spawning herring for the year-classes 1959-61 and 1973-78. ICES CM 1986/H:42, 4 pp. (mimeo).

Watkin, E.E. 1927. Investigations on Cardigan Bay Herring. Part V. Rep. Mar. and Fresh Water Inv. Dept. Zool. Univ. Coll. Wales, 2.

Østvedt, O.J. 1958. Some considerations concerning the homogeneity of the Atlanto-scandian herring. Rapp. P.-v. Reun. Cons. int. Explor. Mer, 143: 53-57.

Table 1 a. Mean length at age.

AREA 04

| Year | Age   |       |       |       |       |
|------|-------|-------|-------|-------|-------|
|      | 0     | 1     | 2     | 3     | 4     |
| 1950 |       |       |       | 19.4  | 21.1  |
| 51   |       |       |       | 20.8  |       |
| 52   |       |       | 19.8  | 23.3  |       |
| 53   |       |       | 17.6  |       | 24.0  |
| 54   |       |       |       | 23.4  |       |
| 55   |       |       | 22.3  |       |       |
| 56   |       |       | 20.7  |       |       |
| 57   |       | 16.5  | 16.8  |       |       |
| 58   | 10.8  | 15.6  |       |       |       |
| 59   | 12.1  |       |       |       |       |
| 60   |       |       |       |       |       |
| Mean | 11.45 | 16.40 | 19.44 | 21.73 | 22.55 |
| 1973 |       | 17.1  | 23.5  | 28.7  | 31.5  |
| 74   | 9.9   |       |       | 29.8  |       |
| 75   |       |       | 26.8  |       |       |
| 76   | 9.9   |       |       |       |       |
| 77   |       |       |       |       |       |
| 78   |       |       | 23.3  |       |       |
| 79   |       | 16.9  |       |       |       |
| 80   | 10.9  | 17.4  | 22.8  | 27.5  | 32.2  |
| 81   |       |       |       | 29.0  |       |
| 82   | 9.0   | 18.0  | 24.2  | 28.4  | 30.8  |
| 83   | 9.6   | 16.6  | 20.5  | 23.6  |       |
| Mean | 9.93  | 17.35 | 24.12 | 28.68 | 31.50 |

Table 1 b. Mean length at age.

AREA 05

| Year | Age   |       |       |       |       |
|------|-------|-------|-------|-------|-------|
|      | 0     | 1     | 2     | 3     | 4     |
| 1950 |       |       | 22.2  |       |       |
| 51   |       |       | 19.0  |       |       |
| 52   |       | 17.4  |       | 21.7  |       |
| 53   |       |       | 19.3  | 24.4  |       |
| 54   |       | 18.2  | 21.6  |       |       |
| 55   |       | 18.8  |       |       |       |
| 56   |       |       | 19.6  | 23.5  |       |
| 57   |       | 14.5  |       |       |       |
| 58   | 10.9  | 18.4  |       |       |       |
| 59   |       |       |       |       |       |
| 60   |       |       |       |       |       |
| Mean | 10.9  | 17.55 | 20.34 | 23.20 |       |
| 1973 |       | 19.2  | 25.2  | 29.7  | 33.2  |
| 74   | 13.5  | 22.6  | 29.2  | 32.4  | 34.1  |
| 75   |       | 21.2  | 29.0  |       |       |
| 76   | 12.6  | 16.8  | 28.2  |       |       |
| 77   |       | 20.2  |       |       |       |
| 78   | 9.8   | 19.4  | 23.3  |       | 30.4  |
| 79   | 10.3  | 18.2  |       | 28.5  | 32.4  |
| 80   | 12.2  |       | 24.7  | 29.5  | 34.2  |
| 81   |       |       | 28.2  | 31.3  | 32.7  |
| 82   | 9.7   | 18.5  | 24.4  | 29.3  | 31.5  |
| 83   | 9.6   | 19.4  | 21.9  | 24.3  | 29.3  |
| Mean | 11.35 | 19.51 | 26.53 | 30.12 | 32.64 |

Table 1 c. Mean length at age.

AREA 06

| Year | Age   |       |       |       |       |
|------|-------|-------|-------|-------|-------|
|      | 0     | 1     | 2     | 3     | 4     |
| 1950 |       |       | 22.3  |       |       |
| 51   |       | 17.2  |       |       |       |
| 52   |       |       |       |       |       |
| 53   |       |       |       | 23.0  |       |
| 54   |       |       | 21.7  |       |       |
| 55   |       | 19.2  |       |       |       |
| 56   |       |       |       |       |       |
| 57   |       | 17.1  |       |       |       |
| 58   | 12.1  |       |       |       |       |
| 59   | 12.7  | 17.6  |       |       |       |
| 60   | 9.7   |       |       |       |       |
| Mean | 11.50 | 17.78 | 22.00 | 23.00 |       |
| 1973 |       | 21.1  | 28.5  | 31.0  | 33.2  |
| 74   | 13.6  | 25.4  | 28.9  | 31.7  |       |
| 75   |       | 22.7  | 27.5  |       | 32.0  |
| 76   | 11.2  | 19.4  | 26.4  | 31.0  | 33.5  |
| 77   | 10.8  | 21.3  | 27.0  | 31.5  | 33.2  |
| 78   | 10.4  | 18.9  | 25.7  | 28.2  | 31.6  |
| 79   | 10.7  | 14.1  | 22.2  | 28.7  | 33.1  |
| 80   | 9.5   | 15.3  | 21.4  | 28.2  | 31.8  |
| 81   |       | 14.4  | 27.0  |       |       |
| 82   | 11.4  | 22.8  | 23.9  | 28.7  |       |
| 83   | 9.3   | 18.8  | 24.3  | 26.3  |       |
| Mean | 11.09 | 19.54 | 25.85 | 29.88 | 32.63 |

Table 1 d. Mean length at age.

AREA 07

| Year | Age   |       |       |       |       |
|------|-------|-------|-------|-------|-------|
|      | 0     | 1     | 2     | 3     | 4     |
| 1950 |       |       |       |       |       |
| 51   |       | 15.8  |       |       |       |
| 52   |       |       | 20.3  |       |       |
| 53   |       |       | 20.4  |       |       |
| 54   |       | 17.7  | 25.6  |       |       |
| 55   |       | 19.7  |       |       |       |
| 56   |       |       |       |       |       |
| 57   |       | 15.2  |       |       |       |
| 58   | 11.9  |       |       |       |       |
| 59   |       |       |       |       |       |
| 60   | 11.6  |       |       |       |       |
| Mean | 11.75 | 17.10 | 22.10 |       |       |
| 1973 |       | 23.6  | 29.6  | 32.0  | 33.3  |
| 74   |       | 24.9  | 30.2  | 32.3  | 33.9  |
| 75   | 13.1  | 18.1  | 28.9  | 32.0  | 34.0  |
| 76   | 12.3  | 20.9  | 29.2  | 32.7  | 33.9  |
| 77   | 10.8  | 19.8  | 29.3  | 31.5  | 33.2  |
| 78   | 11.3  | 24.1  | 27.1  | 29.6  | 32.2  |
| 79   | 9.2   | 20.3  | 26.9  | 30.6  | 32.7  |
| 80   | 15.6  | 21.2  | 27.0  | 28.5  | 32.8  |
| 81   | 12.6  |       | 27.8  | 31.8  | 32.3  |
| 82   |       | 22.3  | 28.0  | 30.7  | 32.3  |
| 83   |       | 22.0  | 28.1  | 27.8  | 30.3  |
| Mean | 12.13 | 21.69 | 28.40 | 31.17 | 33.06 |

Table 4. Weight at age in the spawning stock.

| Year  | Age   |       |       |       |       |       |       |
|-------|-------|-------|-------|-------|-------|-------|-------|
|       | 4     | 5     | 6     | 7     | 8     | 9     | 10    |
| 1950  | 181   | 213   | 249   | 262   | 293   | 303   | 314   |
| 51    | 195   | 245   | 272   | 295   | 308   | 321   | 322   |
| 52    | 208   | 268   | 292   | 307   | 306   | 323   | 334   |
| 53    | 136   | 242   | 260   | 288   | 304   | 326   | 334   |
| 54    | 234   | 252   | 284   | 302   | 320   | 327   | 350   |
| 55    | 221   | 265   | 291   | 313   | 328   | 354   | 370   |
| 56    |       | 251   | 300   | 301   | 359   |       | 369   |
| 57    | 229   | 294   | 293   | 317   |       | 354   |       |
| 58    | 219   | 253   | 264   |       | 306   | 323   | 383   |
| 59    | 187   | 211   | 235   | 249   | 294   | 288   | 319   |
| 60    | 192   | 197   | 222   | 270   | 277   | 309   | 325   |
| Mean  | 200.2 | 244.6 | 269.3 | 290.4 | 309.5 | 322.8 | 342.0 |
| 1973  | 231   | 298   | 369   | 385   | 391   | 385   | 430   |
| 74    | 267   | 338   | 366   | 379   | 361   | 418   | 380   |
| 75    | 282   | 340   | 347   | 322   | 438   | 366   | 390   |
| 76    | 262   | 299   | 321   | 386   | 371   | 398   | 384   |
| 77    | 278   | 262   | 338   | 347   | 382   | 375   | 330   |
| 78    | 181   | 274   | 307   | 347   | 350   | 321   |       |
| 79    | 235   | 278   | 313   | 327   | 286   |       |       |
| 80    | 234   | 238   | 275   | 275   |       |       |       |
| 81    | 212   | 260   | 250   |       |       |       |       |
| 82    | 213   | 224   |       |       |       |       |       |
| 83    | 151   |       |       |       |       |       |       |
| Mean; | 239.5 | 281.1 | 320.6 | 346.0 | 368.4 | 377.2 | 382.8 |
| Diff; | 39.3  | 36.5  | 51.3  | 55.6  | 58.9  | 54.4  | 40.8  |