

1 Long-term variation in numbers and biomass of silver eels being produced in 2 two European river systems

3 Running page head: Annual silver eel production and migration

4 W. Russell Poole^{1*}, Ola H. Diserud², Eva B. Thorstad², Caroline Durif³, Conor Dolan⁴, Odd
5 Terje Sandlund², Knut Bergesen⁵, Gerard Rogan¹, Sean Kelly¹ and L. Asbjørn Vøllestad⁶

6 ¹ Marine Institute, Furnace, Newport, Co. Mayo, Ireland

7 ² Norwegian Institute for Nature Research (NINA), NO-7485 Trondheim, Norway

8 ³ Institute of Marine Research, Austevoll Research Station, Storebø, Norway

9 ⁴ School of Biological Sciences, Queen's University Belfast, 97 Lisburn Road, Belfast BT9 7BL,
10 UK

11 ⁵ NINA Research Station, NO-4308 Sandnes, Norway

12 ⁶ Centre for Ecological and Evolutionary Synthesis, Department of Biosciences, University of
13 Oslo, Oslo, Norway

14 *Corresponding Author: Dr. Russell Poole, russell.poole@marine.ie, Phone: +353-98-42300;
15 FAX: +353-98-42340

16

17

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

1

2

3 **Abstract**

4 The European eel (*Anguilla anguilla*) population has been in decline at least since the 1960s
 5 and reliable regional information, particularly on the spawner production and escapement
 6 (i.e. the silver eel life stage), is a requirement of the EU stock recovery regulation. Two
 7 comparable time series exist in Burrishoole (Ireland) and Imsa (Norway), with monitoring of
 8 total silver eel production since the early 1970s. Numbers of emigrating silver eels fell
 9 significantly ($p < 0.0001$) in the 1980s (breakpoints: Burrishoole 1982; Imsa 1988), in both
 10 catchments, from more than 4,000 eels per annum to approximately 2,000 eels per annum.
 11 The proportion of male eels dropped and the average size of female eels increased. Biomass
 12 of silver eels escaping has remained similar in Burrishoole (1.1/1.2 kg/ha), but not in Imsa
 13 (2.1/0.9 kg/ha), between the early period and the 2000s. Factors that govern the onset of
 14 eel maturation (silvering) and the annual production of silver eels are little understood. In
 15 this paper, the influence of time lagged environmental variables on silver eel production is
 16 examined. Annual variation in the time series was partly (r^2 Burrishoole = 0.43, Imsa = 0.46)
 17 explained by variation in water temperature and water level. Annual number of migrating
 18 eels in both catchments was positively related to summer temperature and summer water
 19 flow, negatively related to summer temperatures in the previous year, and in the
 20 Burrishoole, also negatively related to high water levels in September/October. The models

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

did not transfer well between catchments, indicating likely catchment specific environmental factors impacting on eel production. The reduction in eel numbers observed in both catchments, accompanied by the change in sex ratio and mean weight of females that contribute to maintain biomass production, calls into question the advisability of basing a spawner escapement recovery target on biomass alone, while numbers and proportions of males decline.

Keywords: Silver European eel, *Anguilla anguilla*, annual production, biomass, temperature, water level

Introduction

European eel (*Anguilla anguilla*) spawn in the Sargasso Sea and recruit to the European and north African continental habitats following a trans-Atlantic larval migration. On arrival, the larvae metamorphose into glass eel and a proportion migrates upstream into inland waters where they are subject to many natural and anthropogenic pressures. The growth phase, when they are known as yellow eels, takes many years (sometimes decades) and varies between sexes, with males typically smaller and younger than females. The onset of sexual maturation or “silvering” occurs towards the end of their life cycle and is associated with preparation for the downstream and oceanic migration back to the Sargasso Sea (Tesch, 2003). Triggers for the silvering process, apart from size of the eel, are little understood, but are likely to be linked to growth (Durif *et al.*, 2005). Little is known about the influence of environmental factors on this process. The long-term datasets in this paper provide an

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

opportunity to examine the role of time lagged environmental drivers, i.e., water temperature and level, on the amount of annual silver eel production.

The eel population, which is genetically panmictic (Palm *et al.*, 2009), has been in decline at least since the 1960s, with a severe reduction in glass eel recruitment occurring in the early 1980s (Moriarty, 1990a; Dekker, 2003; Aalto *et al.*, 2016; ICES, 2016). This prompted eel conservation and stock recovery measures within the European Union (European Council, 2007). EU Member States are now required by legislation (EC No. 1100/2007) to contribute to the restoration of the European eel stock by implementing eel management plans. The Regulation seeks to reduce anthropogenic mortalities to permit the escapement of 40% of the silver eel biomass relative to the best estimate of the escapement that would have existed if no anthropogenic influences had affected the stock (European Council, 2007). Setting the historical baselines and monitoring in biomass, rather than numbers, adds an extra challenge for Member States when assessing compliance with the Regulation - given the overall paucity of relevant historical data and limited opportunities for ground truthing outputs from modelling exercises (e.g. Walker *et al.*, 2011).

Long-term datasets of silver eel production and escapement are few, and many are based on fishery yield or some form of escapement estimate, using one or a combination of methods such as mark and recapture, acoustics and fisheries landings (Dekker, 2000a; Rosell *et al.*, 2005; ICES, 2010). It is often difficult to determine the representativeness (proportion of run captured, bias in size or sex ratio) of such monitoring (Feunteun *et al.*, 2000; Allen *et al.*, 2006; MacNamara and McCarthy, 2013; McCarthy *et al.*, 2013). Biases can occur when the

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

fishing season does not cover the full migration period, when there is significant eel production downstream of the fishery area assessed, or when the implementation of management measures on fisheries introduces discontinuities in the data.

Direct assessments of silver eel leaving a given catchment are difficult to obtain due to the migration behaviour of the eel. Silver eels tend to migrate at night, during dark moon periods and in high water discharges making high capture efficiency difficult, especially in larger rivers (Vøllestad *et al.*, 1986; Bruijs and Durif, 2009; Sandlund *et al.*, 2017). Wolf traps (Wolf, 1951) or similar capture systems, can be used to provide precise estimates of migrating eels and in some instances allow all silver eels to be counted and measured. Full quantification of the number, size and sex of silver eel leaving catchments has only been carried out on a few systems with trapping facilities (Vøllestad and Jonsson, 1986, 1988; Poole *et al.*, 1990; Feunteun *et al.*, 2000; Laffaille *et al.*, 2006; Acou *et al.*, 2009). Only the time series in the Imsa (Norway) and Burrishoole (Ireland) presented in this paper include data collected before the collapse in recruitment during the 1980s. Such historical data are a requirement for setting historical baselines for the EU (Piggins, 1985; Vøllestad and Jonsson, 1986, 1988; Poole *et al.*, 1990).

This study brings together the silver eel data sets from two contrasting watersheds in Ireland and Norway describing the stock composition and long-term trends of the silver eel production over a more than 40 year time span in rivers that have experienced little or no exploitation. Neither river has been stocked, nor had hydropower obstruction to migration.

The eel data represent the complete annual number of escaping silver eels which are

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

combined with environmental variables (temperature, water level) to explain how time lagged environmental drivers influence the year-to-year variation in annual numbers of migrating silver eel being produced in each catchment. Similarities and differences between the catchments are discussed, and the use of biomass as a management indicator in the light of continually falling numbers and changing sex ratios is questioned.

Materials and Methods

Site Descriptions

The Burrishoole system in western Ireland (53° 56' N, 9° 35' W: Figure 1; and Supplementary Material) has a catchment area of 100 km² and drains through two channels, the Mill Race and the Salmon Leap, into the North-east Atlantic. The catchment has a total productive wetted area of 474 ha (449 ha lacustrine, 25 ha fluvial). Feeagh and Bunaveela, the two largest freshwater lakes, have mean depths >12 m, are oligotrophic (TP <10 µg l⁻¹), coloured (c. 80 mg l⁻¹ PtCo), and have low alkalinity (<20 mg l⁻¹ CaCO₃) and pH (c. 6.7). Average combined water discharge at the river outlets is 4.2 m³ s⁻¹, ranging between 0.4 and 13 m³ s⁻¹, although higher discharges have been observed but not quantified. Strongly influenced by the Atlantic Ocean (Jennings *et al.*, 2000), the climate is temperate and oceanic, with mild winters and cool summers. Maximum air temperature rarely exceeds 20 °C, while minimum winter temperatures are usually between 2 °C and 4 °C. The geology is predominantly of low buffering capacity, leading to acidic runoff, and overlaid with poorly drained predominantly

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

peaty soils. There has been no commercial or recreational yellow or silver eel fishery and no stocking of eel.

The Imsa River in southwestern Norway (58° 54' N, 5° 57' E: Figure 1; and Supplementary Material) has a catchment area of 128 km². The catchment has a productive wetted area of 1160 ha (>800 ha of lake and approx. 360 ha fluvial habitat). Maximum depths of the five major lakes vary between 27 and 48 m. Water discharge at the river outlet varies between 0.5 and 30 m³ s⁻¹ (mean 4.5 m³ s⁻¹). Similar to Burrishoole, climatic conditions are influenced by the Atlantic Ocean, and it is relatively warm, considering the northern location. Winter air temperatures are rarely below -4 °C and summer temperatures rarely above 24 °C. The Imsa has an average pH >6.8 and ANC ranging between 150 and 230 µeq l⁻¹. There has been no stocking of eel, but there was a restricted seasonal yellow and silver eel fishery upstream of the trapping station, which in the 1990s became relatively small due to fewer landowners participating in eel fishing and only hobby fishing, and the fishery closed from 2006. The catches, when known, from this limited fishery were taken into account in the production data in this study.

[Figure 1.]

Silver Eel Trapping and Recording of Environmental Variables

Trapping of downstream migrating silver eel has been in operation in Burrishoole since 1958 and in Imsa since 1975. The traps at the freshwater outflows of both catchments include Wolf type downstream traps of similar design (Wolf, 1951; McGrath, 1969) employing

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

horizontal grids with 10 mm gaps on a 1:10 inclination. Trapping at Burrishoole involves a fish fence and wolf trap on the Mill Race outflow installed in 1958 and a full flow controlled Wolf trap on the Salmon Leap outflow (McGrath, 1969 and map in the Supplementary material) installed in 1970 (Poole *et al.*, 1990; Poole, 1994). Total annual captures probably represent about 90% \pm 10% of the total run (Piggins, 1985). Years when extreme flood events led to known major losses of eel were 1978, 1984 and 1989 and these have been removed from the data.

In Imsa, trapping commenced in 1975 (Vøllestad and Jonsson, 1986, 1988). The total weight of the catch biomass from commercial silver eel fishing in the river upstream of Imsa was reported annually to the Research Station at Ims (Vøllestad and Jonsson, 1988). This was converted to numbers using the mean weight data of the eels caught in the trap that year, and added to the trap catch to give total silver eel production (in numbers and weight (kg) per ha). No other fishing takes place on a regular basis (Vøllestad and Jonsson, 1988; Bergesen pers obs.).

Daily water level and temperature ($^{\circ}\text{C}$) were recorded on both the Burrishoole and Imsa outflow rivers. In Imsa, water discharge is estimated based on an empirical relationship between water level and discharge. The water level is recorded 15 m above the fish fence on the Mill Race and this is influenced by the amount of debris in the water. For this project, we used the lake water level, taken at the outflow of the lake 75 m upstream and away from any influence of the traps. Since the water level measurements from Burrishoole were

obtained from two locations and with varying recorder setups and influences, a standardized *This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record* Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

Lake Feeagh water level time series was reconstructed, based on relationships estimated from periods with parallel recordings. Over the time-period, daily water temperature ranged from 1.5 °C to 23.5 °C in Burrishoole and 0.2 °C to 24.2 °C in the Imsa, with corresponding annual means of 9.2-12.4 °C and 7.7-10.7 °C respectively. Imsa has warmer summer and colder winters than Burrishoole (Figure 2 top graph). An examination of the temperature anomalies compared to the period 1971-2000 (1975-2000 for Imsa) indicated a period of warming in both catchments since 1997/'98 (Figure 2 bottom graph).

[Figure 2.]

Data Collection and Analysis

In both catchments, traps were attended at least once a day and all silver eels were counted and recorded. Annual numbers analysed in this paper are the summation of the daily trap counts from 1st May to the 30th April in the following year, called 'Silver Season'. Samples of descending silver eels were taken throughout the run, measuring total length (TL: to the nearest 0.1 cm Burrishoole, 0.5 cm Imsa) and weight to the nearest 1 g (Imsa) or 5 g (Burrishoole). Length and weight data were not collected in the Imsa between 1993 and 2011 and only batch weights were taken in Burrishoole between 1971 and 1975, 1978/'79 and 1980 to 1983. For consistency, the sex ratio was determined from length frequency analysis of the sampled eels. The overlap between the male and female modes was separated using the technique described by Bhattacharya (1967). These analyses were

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

confirmed by random dissections (Burrishoole; $n = 1303$; 1986-2016; Imsa; see Vøllestad and Jonsson 1986) using gross macroscopic examination (Frost, 1945; Bertin, 1956).

Production of silver eel was calculated in numbers and biomass (kg) per hectare (wetted area) of the watershed upstream of the trap. Biomass was calculated using the mean weight and total count, including the reported catches upstream of the trap in the Imsa.

Structural changes, or breakpoints, in the time series were dated using the function breakpoint in the R package *strucchange* (R Core Team, 2016) where we assume that

coefficients shift from one stable relation to another (e.g. Bai and Perron 2003). A Two

Sample t-test was used to verify that the difference in the means of the two time series

before and after the breakpoint was significant. After the structural breaks in the data were

accounted for, time series analyses were performed (e.g. Wei, 2006) to evaluate possible

serial dependencies. For each catchment, autoregressive integrated moving average

(ARIMA) models were fitted to each time series segment and tested whether the same AR

structure existed within a catchment before and after the break. Finally, with the break

points and potential time series structures established, Generalised Linear Models (GLM)

with a Poisson error distribution were fitted to look for covariates that together with, or

instead of, the AR-terms explained the variance in the number of emigrating eels.

Statistical modelling of variation in eel run and the relationship to environmental parameters

was carried out using the statistical software R (R Core Team, 2016). In the initial models of

the annual eel count in both catchments, monthly averages for temperature ($T_{Y,M}$: monthly

mean water temperature from May the preceding year until November this year; the

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

subscript Y denotes year, either the year of the eel counts (t) or the preceding year ($t-1$), while the subscript M denotes month (*Jan - Dec*) and standardized water level ($WL_{Y,M}$: monthly mean standardized water level from May the preceding year until November this year; the subscripts have the same interpretation as for water temperature) were used as explanatory variables. The water level time series was standardized, since we assumed that it is the variation around the mean that is informative and not the absolute value at some arbitrary location. The two-year period was chosen, assuming that the various underlying processes preparing for migration may last for an extended period. Different river systems likely have different bottlenecks or optimal periods for maturation and migration, so we should initially include a long period in the analysis. The effects of temperature and water level were assumed to be accumulated over a few months, but rather than determining these periods a priori, subsequent months that had the same effect on the model response were pooled to quantify the mean value over a longer period. For example, $T_{t-1, May-Nov}$ will denote the mean water temperature from May to November the preceding year.

A maximum linear regression model including all explanatory variables was simplified by stepwise reduction based on the AIC (Sakamoto *et al.*, 1986). If consecutive monthly averages for either water level or temperature had the same effect on the response, a model where these months were pooled together was also fitted and evaluated. A decrease in AIC of more than two was considered as sufficient support for retaining a variable in the model. Residuals were checked for normality and autocorrelation, since the migrating part

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

of a population in subsequent years may be correlated, especially for species with a wide age distribution such as eel.

Results

Annual variation in number of silver eels

There was considerable annual variation in silver eel counts in both Burrishoole and Imsa rivers (Figure 3). In both rivers, there was a downward shift in the 1980s in the annual numbers of migrating eels. In Burrishoole, breakpoint analysis indicated a break in the time series after 1982 (95% CI = 1980-1985), when the average count changed significantly (two sample t-test; $p < 0.0001$) from 4719 eels to 2821 eels. In Imsa, breakpoint analysis indicated a break after 1988 (95% CI = 1987-1990) when the average count changed significantly (two sample t-test; $p < 0.0001$) from 5815 eels to 2201 eels (data supplied in the Supplementary Material Tables 1 and 2). It should be noted that before the abrupt shifts, the average number of eels in Imsa was higher than in Burrishoole, while after the shift, the counts were higher in Burrishoole (Figure 3).

For Burrishoole, after the break in 1982 had been accounted for, no significant AR terms were found in either of the time series segments. For the Imsa time series with a break in 1988, the AR model with lag one year was close to being significant before the break ($AR(1) = -0.404$; $p = 0.085$), and not significantly different from the model fitted to the time series after the break. Assuming the same autoregressive structure before and after the break, the AR model still showed a weak significance ($AR(1) = -0.281$; $p = 0.061$). A negative

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

1 AR(1) coefficient indicates a temporal pattern where high emigration in one year may lead to
 2 a reduced emigration the next year.

3 [Figure 3.]

4 Influence of water temperature and level on annual eel run

5 Finally, with break points accounted for and weak serial dependencies established for the
 6 time series, we fitted GLMs with a Poisson error distribution to the annual eel runs. The one
 7 year lagged autocorrelations of the time series were better explained by lagged covariates of
 8 water temperature, so no AR terms were retained in the models.

9 By fitting a model to the whole detrended time series, i.e. breaks accounted for, we assume
 10 that the modelled relationships were the same in both periods and that the covariates
 11 cannot explain the shift in mean level. Models fitted to the two periods separately gave
 12 similar results, and there was no obvious relationship between the shift in eel numbers and
 13 the temperature or water level variables, which supported these assumptions. The fit of the
 14 models (Table 1) shows that the eel run in most years is well described by the model (Figure
 15 4), although a few years have large residuals, indicating that there are some environmental
 16 variables affecting the eel run in some years that are not accounted for by the model. The
 17 years with large residuals occur in different years in the two systems (Figure 4).

18 The maximum model for Burrishoole, including all explanatory variables, had an AIC of
 19 683.5. Stepwise removal of variables that did not improve the AIC, and biologically sensible

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

pooling of months that had the same effect on the response, resulted in a simplified model with four variables and an AIC of 670.6 and $R^2 = 0.43$ (Table 1).

The detrended eel numbers in Burrishoole can be explained by a negative relationship with a warm previous season ($T_{t-1, May-Nov}$ – mean water temperature over all months from May to November), and a positive relationship to the current year's conditions in August (both temperature and water level), while high mean water levels in September and October caused a decrease in the number of migrating eels (Table 1).

The maximum model for Imsa, including all explanatory variables, had an AIC = 652.6.

Applying the same variable selection approach as for Burrishoole resulted in a model with an AIC = 644.8 and $R^2 = 0.48$ (Table 1). The detrended eel numbers in Imsa have a negative relationship with the mean temperature in June and July the previous year, and a positive relationship with mean temperatures in September and October the previous year. There was a positive relationship with mean temperatures in the summer (June and July) of the same calendar year, and with the mean standardized water level in January and February in the same year (Table 1).

When Imsa data were used to validate the Burrishoole model, the validation R^2 was close to zero. A similar result was obtained using Burrishoole data to validate the Imsa model, i.e. R^2 almost equal to zero. Thus, predicting the quantity of annual eel runs in one of the catchments from a model fitted in the other catchment gave poor results.

[Figure 4.]

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

1 [Table 1].

2 Size and Sex

3 We used length as the indicator of eel size, while noting that there is a highly significant
4 exponential relationship between length and weight allowing conversion to biomass.

5 Although the Burrishoole eels were on average 10% heavier than the Imsa eels of the same
6 length, the slopes in the log-log models were similar (Burrishoole $\log(\text{Weight}) =$
7 $3.11 \cdot \log(\text{Length}) - 6.72$, $R^2 = 0.94$; Imsa $\log(\text{Weight}) = 3.12 \cdot \log(\text{Length}) - 6.83$, $R^2 = 0.97$).

8 There was a strong sex related bimodal distribution of eel length in both Burrishoole and
9 Imsa although the proportions differed between locations and over time (see Figure 5 for
10 sample years). In Burrishoole, male eels ranged from 27.6 cm to 45.4 cm body length
11 (dissected eels, $n = 445$) and females ranged from 37.5 cm to 105.3 cm (dissected eels, $n =$
12 859). Only three females were shorter than 40 cm. In Imsa, male eels ranged from 32.5 cm
13 to 46.0 cm body length ($n = 119$) and females from 39.0 cm to 107.0 cm ($n = 4164$).

14 A multivariate linear regression model showed differences in the size of eels for each sex,
15 both over time and between locations (see Supplementary Material Table 2). For male eels,
16 and noting the large disparity in numbers of males between the two catchments, the males
17 in Burrishoole were significantly smaller than those in the Imsa (-3.7cm ; $p < 0.0001$) in all
18 periods. The males in Burrishoole decreased significantly in size by decade ($p < 0.001$) over
19 the sample period, with males averaging 37.0 cm in the 1980s, 36.7 cm in the 1990s and 35.7

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

cm after 2000 (Figure 6). There were too few males in the Imsa to examine any change over time.

The female length model was highly significant ($p < 0.001$) with both decadal differences in each catchment and a significant interaction between catchments, noting the missing data in the Imsa in the 1990s and 2000s, and with competing tendencies in the 2010s when the Burrishoole females decreased in length while Imsa females increased. The female eels in Burrishoole were significantly shorter (-11.6cm; $p < 0.001$) than those in the Imsa in all years sampled (Figure 5; and Supplementary Material). The female eels in Burrishoole increased in length from the 1970s until 1998, stabilised until 2005 and then started to decline in size again (Figure 7), averaging 50.3 cm in the 1970s/80s, 52.0 cm in the 90s, 52.8 cm in the 2000s and 49.6 cm in the 2010s (Figure 7; and Supplementary Material Table 3). In the Imsa, females were similar in length from 1982 to 1991 (61.9 cm in the 80s and 62.8 cm in 91/92) and then increased to 68.7 cm in the last four years of the 2010s (Figure 7).

There has been a drop in the proportion of male eels in both catchments, in Burrishoole from >55% male before 1985 to about 35% in the last decade, and in the Imsa from 5.6% in the early 1980s to <0.2% in the last five years (Table 2).

[Figure 5.]

[Figure 6.]

[Figure 7.]

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

1 Production

2 The average silver eel production in numbers per ha was higher in Burrishoole than in Imsa
3 for both periods (ANOVA, $p < 0.0001$) (pre- and post-1982 in Burrishoole; pre- and post-1988
4 in Imsa), but the Imsa had a significantly (ANOVA, $p < 0.0001$) higher production in biomass
5 per ha due to the larger size of the female eels (Table 2, Figure 8). In Burrishoole, the
6 reduction in numbers over time was accompanied by a slight, but not significant ($p = 0.134$)
7 increase in biomass while, in contrast, both the numbers and biomass fell significantly
8 (ANOVA, $p < 0.0001$) in the Imsa (Figure 8; ANOVA outputs supplied in the Supplementary
9 Material Table 4). This was due to the larger reduction in the number of males in the
10 Burrishoole and the increase in the average weight of the females while the number of
11 females remained similar in both periods (Table 2). In the Imsa, males were relatively few in
12 both periods. The reduction in numbers of females outweighed their increased average size,
13 leading to the reduction in both numbers and biomass in the Imsa.

14 [Figure 8.]

15 [Table 2]

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

1 Discussion

2 The Burrishoole (Ireland) and Imsa (Norway) rivers have been independently monitored
 3 using full river trapping systems of similar design (see McGrath, 1969) for more than 45
 4 years. While the eel stock in each catchment has its own particular characteristics, some of
 5 the similarities are striking. In the 1980s, both catchments experienced a sudden drop in
 6 numbers of eels emigrating, a reduction in the proportion of males in the silver eel run and
 7 an increase in the size of females. It seems likely that maturation (silvering), and hence the
 8 level of annual production of silver eel, was positively related to summer temperature and
 9 summer water flow, negatively related to summer temperatures in the previous year, and in
 10 the Burrishoole, also negatively related to high water levels in September/October.

11 Both catchments have experienced higher temperatures in the last 20 years compared to
 12 the 30 years before that, the Imsa more so than Burrishoole (Fealy *et al.*, 2014; this paper).
 13 Burrishoole has also shifted towards mesotrophic conditions since the 1950s, associated
 14 with the commencement of commercial coniferous plantations (Dalton *et al.*, 2014).

15 Untangling the relative influence of different factors, such as temperature, eel density and
 16 trophic changes, on eel stocks is difficult. Changing eel growth across Europe has been
 17 associated with rising temperature over the last century (Daverat *et al.*, 2012) and the silver
 18 eel run has been commencing earlier (on average 0.8 days/year) in Burrishoole (Sandlund *et*
 19 *al.*, 2017). However, to our knowledge there are no previous studies showing how the level

20 of annual production of silver eel might be influenced by time-lagged environmental factors.
This is a pre-copyedited, author-produced version of an article accepted for publication in ICES
Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud,
 Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut
 Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river
 systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is
 available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

1 Little is known about the mechanism that governs the decision to mature and migrate, once
 2 an eel is in the optimum size range for the river, or about the factors that influence the
 3 physiological readiness to migrate, although these are likely to involve significant changes in
 4 body constitution and energy (e.g. fat, protein and dry matter) and also raised cortisol levels
 5 in the autumn linked to energy mobilisation (Van Ginneken *et al.*, 2007). The environmental
 6 or physiological conditions that trigger the onset of maturation and silvering are still poorly
 7 understood (Durif *et al.*, 2009).

8 Environmental variables (e.g. water level, water temperature) likely to influence the onset of
 9 silvering, using variables from the year of the migration and from the preceding year,
 10 explained approximately half the variation in annual eel count in each catchment. While
 11 there was little evidence of a regular pattern of low years following high years in either
 12 catchment, high summer temperatures were associated with greater numbers migrating in
 13 that year, and consequently lower numbers migrating the following year. This was shown by
 14 a significant relationship with temperature in June/July (Imsa) and May/November
 15 (Burrishoole) in the previous year, and high temperatures in the current year in June/July
 16 (Imsa) and in August (Burrishoole). Water level in August in Burrishoole was also positively
 17 related to eel count. The onset of migration in the Loire River (France) was correlated with
 18 August temperature, discharge and sunshine hours (Durif and Elie, 2008). Temperature-
 19 related increases in growth and physiological activity could encourage a higher proportion of
 20 eels to reach the required physiological threshold switch for the onset of silvering in that
 21 year. Consequently, in the following year, there will be a lower proportion of eels near that

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

threshold as they had migrated in the previous season and therefore less eels are available to migrate in those years following high temperatures. The significant positive relationship in Imsa, with high temperatures in September/October leading to a higher migration in the following year, also supports that argument, where eels that were not quite at the threshold in the current season may have a longer growing season thereby facilitating silvering in the following year.

The relationship with water level in the Imsa in January and February may be related to an interaction between warm winters and higher precipitation or less ice, and therefore earlier influences on eel physiology than in colder drier winters. Such cold winters usually do not occur in Burrishoole. In Burrishoole, eel counts were lower than expected with increasing water level in September/October. This is counter-intuitive, unless we consider that this was an artefact related to high floods reducing trapping efficiency, which is considered unlikely.

The size of female silver eel and proportion of males observed in the two catchments in this study are consistent with the cline in these parameters demonstrated for American eel *A. rostrata* (Helfman, 1988; Oliveira, 1999; Jessop, 2010) and European eel (Vøllestad, 1992), with less males and larger females with increasing latitude and distance from the Sargasso Sea. In the 1960s, partial trapping in Burrishoole indicated a silver eel male sex ratio of over 94% (Piggins, 1985), a fact supported by a yellow eel fyke net survey (Moriarty, 1974). The male ratio declined to circa 30% in the 1980s (Poole *et al.*, 1990), and is currently around 44% in the most recent decade (this study). In contrast, the silver eel run in Imsa was

dominated by females in the 1970s and 1980s, with the male sex ratio between 1% and 7%.

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

(Vøllestad and Jonsson, 1986, 1988), and this has now decreased to less than 0.5% males. These changes in sex ratio in both catchments have been accompanied by an increase in the average size of females. The number of females per annum in Burrishoole has remained similar while it has fallen in the Imsa since the mid-1980s. Andersson *et al.* (2012) noted a considerable decline in escapement of eels in the southern areas of Sweden over the past 50 years, but particularly in the 1960s and '70s, inferred from fishing records and fisheries independent surveys. It seems likely that falling density of eel leads to lower proportions of males and some compensation by an increase in mean weight of silver eel (De Leo and Gato, 1996; Andersson *et al.*, 2012; this study). The ratios of number to biomass observed in this study are probably catchment specific and are likely to vary with latitude (Vøllestad, 1992) depending on the proportion and size of the female eels present in the catchments.

Production and escapement estimates are rare, especially before the 1980s (see review table 4 in the Supplementary material) with the majority based on fishing yield or on estimates of escapement using mark-recapture or partial counts. In recent years, the Burrishoole has been producing about 1.1-1.4 kg/ha or 5.3-6.9 eels/ha, which is a similar weight per hectare as estimated for the Fremur catchment (1.3 kg/ha; Feunteun *et al.*, 2000), the Shannon (1.5-1.6 kg/ha; MacNamara and McCarthy, 2013) and the Erne (1.5-1.7 kg/ha, McCarthy *et al.*, 2013). However, it is a lower production in weight per hectare than estimated for the Imsa (3.51 kg/ha; Vøllestad and Jonsson, 1988), the Oir (4.8-6.9 kg/ha; Acou *et al.*, 2009) and the Lake Ijsslemeer (4.4 kg/ha; Dekker, 2000b). Production in the Burrishoole and these other freshwater catchments is much lower than estimated for

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

coastal lagoons in south-western France (Bages-Sigean: 30 kg/ha; Amilhat *et al.* (2008) and the Mediterranean (Comacchio: 20 kg/ha; Rossi 1979, Sardinia: 19 kg/ha; Rossi and Cannas, 1984). High yields have also been reported from some freshwater catchments where the yield was maintained by active transport and stocking of glass eels and elvers, such as Lough Neagh in Northern Ireland (>20 kg/ha; Moriarty, 1988).

While the biomass production in Burrishoole has remained similar since the 1970s, the number of eels has decreased from an average of 11.5 eels/ha in the late 1970s to an average of 5.8 eels/ha in the 2000s. Production in 2015 was the lowest on record in the Burrishoole and second lowest in Imsa. In contrast, both biomass and numbers have fallen in the Imsa over a similar period. A decline in silver eel production has also been reported in the Irish River Shannon, at least since 1992, but due to changing fishery management practices it is difficult to understand the dynamics of that stock (Moriarty 1990b; McCarthy and Cullen, 2000; MacNamara and McCarthy, 2013).

Mature eels are sexually dimorphic in size, with males being smaller than females. Both genetic and environmental factors have been implicated in sex determination and, for European eel, it has been related to various factors including population density, levels of recruitment and catchment characteristics (Davey and Jellyman, 2005). In general, male silver eels are more abundant at high densities whereas females predominate at lower densities (Parsons *et al.*, 1977; Vøllestad and Jonsson, 1988). It is generally understood that in the European eel, optimum size for each sex is promoted, rather than age, and that this optimum size, especially in females, varies with distance from the Sargasso and with latitude

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

1 (Vøllestad, 1992). The majority of female silvers in Imsa are larger than in Burrishoole (this
 2 paper; Haraldstad *et al.*, 1985; Vøllestad and Jonsson, 1986; Poole *et al.*, 1990; Poole and
 3 Reynolds, 1996), and in recent years this difference has increased.

4 Sex ratios can vary widely between catchments and over time (Parsons *et al.*, 1977; Jessop,
 5 1987; Oliveira, 1999; Oliveira and McCleave, 2000, 2001; Poole *et al.*, 1990; Tesch, 2003).
 6 Changes in stocking densities (Parsons *et al.*, 1977; Wickström *et al.*, 1996) and natural
 7 recruitment (Rossi *et al.*, 1988; Bark *et al.*, 2007) have been implicated in altered sex ratios in
 8 some systems. It seems likely, therefore, that the changes observed in the Burrishoole and
 9 Imsa Rivers are a response to the observed low recruitment, which has been consistent with
 10 the decline observed across Europe since the early 1980s (Poole, 1994; Bornarel *et al.*, 2017;
 11 Durif *et al.*, in prep), with falling local stock densities and maybe also changes induced by
 12 rising temperatures and catchment enrichment. Bark *et al.*, (2007) illustrated temporal
 13 changes in population density and population structure in the River Piddle and other rivers
 14 in England and Wales, where a decline in population density since the mid-1970s was
 15 accompanied by a profound change in population structure, from one dominated by small
 16 eels to one currently dominated by large eels, clearly suggestive of a major recruitment
 17 failure. In rivers where eel populations were stable, the populations were male biased, while
 18 rivers with declining populations were female biased (Bark *et al.*, 2007). Similarly, Laffaille *et*
 19 *al.* (2006) associated a shift from male to female dominance with a falling number of silver
 20 eels in the Fremur in France in an eight-year period after 2000, likely due to a number of

This is a pre-copiedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

factors including the installation of eel passes on the main hydraulic engineering structures in 1992 and 1996.

While sex ratio and eel size have been changing over a longer time scale in Burrishoole (Poole *et al.*, 1990), likely influenced by environmental changes in the catchment (Dalton *et al.*, 2014), it is not easy to find an explanation for the sudden drop in numbers of eels in Burrishoole and in Imsa, particularly as the annual silver eel run is composed of up to 20 or more age classes of eel (Poole and Reynolds, 1996; Durif *et al.*, in prep). The six-year difference between catchments in the main drop in numbers during the 1980s is also difficult to explain.

The collapse in recruitment observed in the 1980s across Europe was preceded by a longer-term decline in landings, indicating a probable decline in the continental potential spawning stock (Dekker, 2003) and leading to a conclusion that insufficient spawning stock biomass might have caused the recruitment collapse. More current observations indicate possible Allee-effects (Allee, 1931), or compensatory mechanisms (Hillborn and Walters, 1992), taking place in the stock-recruitment relationship and likely caused by a disruption in the social mating system below a minimum spawning density (Dekker, 2004, 2008). The spawning target set in the EU Regulation is based on biomass of the combined sexes (European Council, 2007). If the demographics (e.g. numbers, size and sex ratio) of the eel stock are changing, as indicated in these catchments (this paper), and other areas (e.g. De Leo and Gato, 1996; Bark *et al.*, 2007; Andersson *et al.*, 2012), it is possible that by concentrating on a biomass only approach we may overlook social aspects of eel reproduction and miss the

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

1 opportunity to positively influence the spawning stock recovery and further contribute to
2 the decline in the stock.

3 In conclusion, annual counts and biomass of silver eels in the Burrishoole and Imsa rivers
4 were remarkably similar: the numbers of eels emigrating from the 1980s to present
5 decreased in both catchments, the Imsa lagging six years behind the Burrishoole. In both
6 catchments, the proportion of silver males decreased and the size of the females increased,
7 likely due to a combination of stock density and changing productivity of the catchments.
8 Silver eel runs were more important after warm summers and this resulted in a reduction of
9 the number of silver eels in the following year. Warm autumn temperatures in the Imsa may
10 extend the growing season, facilitating more eels to silver the following year. The reduction
11 in numbers observed in both catchments, accompanied by the change in sex ratio and mean
12 weight of females that contribute to maintain biomass production, especially in the
13 Burrishoole, calls into question the advisability of basing the silver eel recovery target under
14 the EU Regulation solely on biomass, while numbers and proportions of males continue to
15 decline.

16 **Supplementary Material**

17 The following supplementary material is available at ICESJMS online, detailed maps of the
18 Burrishoole and Imsa Catchments, annual silver eel count and size data for each catchment
19 and a review table of published eel yield and production/escapement data.

20 **Acknowledgements**

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

This study was funded by the Irish Marine Institute, the Norwegian Institute for Nature Research, the Norwegian Environment Agency and the European Inland Fisheries and Aquaculture Advisory Commission (EIFAAC) and the authors would like to acknowledge the long-term commitment to maintaining the monitoring stations. We would like to gratefully thank the technical and field staff at both Imsa and Burrishoole for their considerable support and Kari Sivertsen (NINA) for support with the graphics.

References

- Aalto, E., Capoccioni, F., Terradez Mas, J., Schiavina, M., Leone, C., De Leo, G., and Ciccotti, E. 2016. Quantifying 60 years of declining European eel (*Anguilla anguilla* L., 1758) fishery yields in Mediterranean coastal lagoons. *ICES Journal of Marine Science*, 73: 101-110.
- Acou, A., Gabriel, G., Laffaille, P. and Feunteun, E. 2009. Differential production and condition indices of premigrant eels in two small Atlantic coastal catchments of France. *In*: Casselman, J. M. & Cairns, D. K., eds. *Eels at the edge: science, status, and conservation concerns*, 2009 Bethesda, Maryland. American Fisheries Society Symposium, 157-174.
- Allee, W. C. 1931. *Animal Aggregations. A study in General Sociology*. University of Chicago Press, 431pp.

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

- 1 Allen, M., Rosell, R., and Evans, D. 2006. Predicting catches for the Lough Neagh (Northern
2 Ireland) eel fishery based on stock inputs, effort and environmental variables.
3 Fisheries Management and Ecology, 13: 251-260.
- 4 Amilhat, E., Farrugio, H., Lecomte-Finiger, R., Simon, G., and Sasal, P. 2008. Silver eel
5 population size and escapement in a Mediterranean lagoon: Bages-Sigean, France.
6 Knowledge and Management of Aquatic Ecosystems, 05: 390-391.
- 7 Andersson, J., Florin, A.-B., and Petersson, E. 2012. Escapement of eel (*Anguilla anguilla*) in
8 coastal areas in Sweden over a 50-year period. ICES Journal of Marine Science, 69:
9 991-999.
- 10 Bai, J., and Perron, P. 2003. Computation and Analysis of Multiple Structural Change
11 Models. Journal of Applied Econometrics, 18: 1-22.
- 12 Bark, A., Williams, B., and Knights, B. 2007. Current status and temporal trends in stocks of
13 European eel in England and Wales. ICES Journal of Marine Science, 64: 1368-1378.
- 14 Bertin, L. 1956. Eels: A Biological Study, Cleaver-Hume Press Ltd, London.
- 15 Bhattacharya, C. G. 1967. A simple method of resolution of a distribution into Gaussian
16 components. Biometrics, 23: 115-135.
- 17 Bornarel, V., Lambert, P., Briand, C., Antunes, C., Belpaire, C., Eleonora, C., Diaz, E., *et al.*
18 2017. Modelling the recruitment of European eel (*Anguilla anguilla*) throughout its
19 European range. ICES Journal of Marine Science, doi:10.1093/icesjms/fsx180: 12pp

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. ICES Journal of Marine Science 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

1

2 Bruijs, M. C. M., and Durif, C. M. F. 2009. Silver eel migration and behaviour. In Spawning

3 migration of the European eel: Reproduction index, a useful tool for conservation

4 management, G. Van den Thillart, S. Dufour and J. C. Rankin, eds. (Fish & Fisheries

5 Series 30, Springer), pp. 65-95.

6 Dalton, C., O'Dwyer, B., Taylor, D., de Eyto, E., Jennings, E., Chen, G., Poole, R., *et al.* 2014.

7 Anthropocene environmental change in an internationally important oligotrophic

8 catchment on the Atlantic seaboard of western Europe. *Anthropocene*, 5: 9-21.

9 Daverat, F., Beaulaton, L., Poole, R., Lambert, P., Wickstrom, H., Andersson, J., Aprahamian,

10 M., *et al.* 2012. One century of eel growth: changes and implications. *Ecology of*11 *Freshwater Fish*, 21: 325-336.

12 Davey, A. J. H., and Jellyman, D. J. 2005. Sex determination in freshwater eels and

13 management options for manipulation of sex. *Reviews in Fish Biology and Fisheries*

14 15, 37-52.

15 De Leo, G. A., and Gatto, M. 1996. Trends in vital rates of the European eel: evidence for

16 density dependence? *Ecological Applications*, 6: 1281-1294.17 Dekker, W. 2000a. The fractal geometry of the European eel stock. *ICES Journal of Marine*18 *Science*, 57: 109-121.

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

- 1 Dekker, W. 2000b. Impact of yellow eel exploitation on spawner production in Lake
2 IJsselmeer, the Netherlands. *Dana*, 12: 17-32.
- 3 Dekker, W. 2003. Did lack of spawners cause the collapse of the European eel, *Anguilla*
4 *anguilla*? *Fisheries Management and Ecology*, 10: 365-376.
- 5 Dekker, W. 2004. Slipping through our hands: population dynamics of the European Eel. PhD
6 Thesis, Faculteit der Natuurwetenschappen, University of Amsterdam, Amsterdam:
7 186pp.
- 8 Dekker, W. 2008. Coming to grips with the eel stock slip-sliding away. Pages 335-355 in M.G.
9 Schlecter, N.J. Leonard and W.W. Taylor, editors. *International Governance of*
10 *Fisheries Eco-systems: Learning from the Past, Finding Solutions for the Future.*
11 *American Fisheries Society, Symposium 58, Bethesda, Maryland.*
- 12 Durif, C., Dufour, S., and Elie, P. 2005. The silvering process of *Anguilla anguilla*: a new
13 classification from the yellow resident to the silver migrating stage. *Journal of Fish*
14 *Biology*, 66: 1025-1043.
- 15 Durif, C. M. F., and Elie, P. 2008. Predicting downstream migration of silver eels in a large
16 river catchment based on commercial fishery data. *Fisheries Management and*
17 *Ecology*, 15: 127-137.
- 18 Durif, C., Guibert, A., and Elie, P. 2009. Morphological discrimination of the silvering stages
19 of the European eel. *In Eels at the edge: science, status, and conservation concerns,*

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

- 1 pp. 103-111. Ed. by C. J. M, and D. K. Cairns. American Fisheries Society Symposium,
2 Bethesda, Maryland.
- 3 European Council. 2007. Council Regulation (EC) No 1100/2007 of 18 September 2007
4 establishing measures for the recovery of the stock of European eel. Official Journal
5 of the European Union, L 248/17: 17-23.
- 6 Fealy, R., Allott, N., Broderick, C., de Eyto, E., Dillane, M., Erdil, R. M., Hancox, L., *et al.* 2014.
7 RESCALE: Review and Simulate Climate and Catchment Responses at Burrishoole.
8 409 pp. Marine Institute, Galway.
- 9 Feunteun, E., Acou, A., Laffaille, P., and Legault, A. 2000. European eel (*Anguilla anguilla*):
10 prediction of spawner escapement from continental population parameters.
11 Canadian Journal of Fisheries and Aquatic Sciences, 57: 1627-1635.
- 12 Frost, W. E. 1945. The age and growth of eels (*Anguilla anguilla*) from the Windermere
13 catchment area. Journal of Animal Ecology, 14: 106-124.
- 14 Haraldstad, Ø., Vøllestad, L. A., and Jonsson, B. 1985. Descent of European silver eels,
15 *Anguilla anguilla* L., in a Norwegian watercourse. Journal of Fish Biology, 26: 37-41.
- 16 Helfman, G. S., 1988. XI. Fish and Fisheries. Patterns in the life history of anguillid eels.
17 Verhandlungen des Internationalen Verein Limnologie, 23; 1663-1669.
- 18 Hilborn, R., and Walters, C. J. 1992. Quantitative fisheries stock assessment: choice, dynamics
19 and uncertainty. Reviews in Fish Biology and Fisheries, 2: 177-186.

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

- 1 ICES. 2010. Report of the Joint EIFAAC/ICES Working Group on Eels (WGEEL), 9-14
2 September 2010: 201 pp.
- 3 ICES. 2016. Report of the Joint EIFAAC/ICES Working Group on Eels (WGEEL), 15-22
4 September 2016: 106 pp.
- 5 Jennings, E., Allott, N., McGinnity, P., Poole, R., Quirke, W., Twomey, H., and George, G.
6 2000. The North Atlantic Oscillation: Effects on Freshwater Systems in Ireland.
7 Biology and Environment: Proceedings of the Royal Irish Academy, 100B: 149-157.
- 8 Jessop, B. 1987. Migrating American Eels in Nova Scotia. Transactions of the American
9 Fisheries Society, 116: 161-170.
- 10 Jessop, B. M. 2010. Geographic effects on American eel (*Anguilla rostrata*) life history
11 characteristics and strategies. Canadian Journal of Fisheries and Aquatic Sciences,
12 67: 326-346.
- 13 Laffaille, P., Acou, A., Guioulet, J., Mounaix, B., and Legault, A. 2006. Patterns of silver eel
14 (*Anguilla anguilla* L.) sex ratio in a catchment. Ecology of Freshwater Fish, 15: 583-
15 588.
- 16 MacNamara, R., and McCarthy, T. K. 2013. Silver eel (*Anguilla anguilla*) population dynamics
17 and production in the River Shannon, Ireland. Ecology of Freshwater Fish, 23: 181-
18 192.

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

- 1 McCarthy, T. K., and Cullen, P. 2000. The River Shannon silver eel fisheries: variations in
2 commercial and experimental catch levels. *Dana*, 12: 59-68.
- 3 McCarthy, T. K., Nowak, D., Grennan, J., Bateman, A., Conneely, B., and Macnamara, R. 2013.
4 Spawner escapement of European eel (*Anguilla anguilla*) from the River Erne,
5 Ireland. *Ecology of Freshwater Fish*, 23: 21-32.
- 6 McGrath, C. J. 1969. The fish trapping installation at the Salmon Leap, Furnace, Co. Mayo.
7 Annual Report of the Salmon Research Trust, Appendix V: 107 pp.
- 8 Moriarty, C. 1974. Eel research in 1973. An Roinn Talmhaíochta agus Iascaigh, Fishery
9 Leaflet, 60: 9pp.
- 10 Moriarty, C. 1988. The eel in Ireland. *In* Went Memorial Lecture 1987, Royal Dublin Society,
11 Occasional Papers in Irish Science and Technology: 9pp.
- 12 Moriarty, C. 1990a. European Catches of Elver of 1928-1988. *Internationale Revue der*
13 *gesamten Hydrobiologie*, 75: 701-706.
- 14 Moriarty, C. 1990b. Short note on the silver eel catch on the lower River Shannon.
15 *Internationale Revue der gesamten Hydrobiologie*, 75: 817-818.
- 16 Oliveira, K. 1999. Life history characteristics and strategies of the American eel, *Anguilla*
17 *rostrata*. *Canadian Journal of Fisheries and Aquatic Sciences*, 56: 795-802.

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

- 1 Oliveira, K., and McCleave, J. D. 2000. Variation in population and life history traits of the
2 American eel, *Anguilla rostrata*, in four rivers in Maine. *Environmental Biology of*
3 *Fishes*, 59: 141-151.
- 4 Oliveira, K., and McCleave, J. 2001. Regional variation and the effect of lake: river area on sex
5 distribution of American eels. *Journal of Fish Biology*, 58: 943-952.
- 6 Palm, S., Dannewitz, J., Prestegard, T., and Wickstrom, H. 2009. Panmixia in European eel
7 revisited: no genetic difference between maturing adults from southern and
8 northern Europe. *Heredity*, 103, 82–89.
- 9 Parsons, J., Vickers, K. U., and Warden, Y. 1977. Relationship between elver recruitment and
10 changes in the sex ratio of silver eels *Anguilla anguilla* L. migrating from Lough
11 Neagh, Northern Ireland. *Journal of Fish Biology*, 10: 211-229.
- 12 Piggins, D. J. 1985. The silver eel runs of the Burrishoole river system: 1959-1984; numbers,
13 weights, timing and sex ratios. *ICES CM 1985/M:5*. Mayo, Ireland: International
14 Council for the Exploration of the Sea, Copenhagen: 12pp.
- 15 Poole, W. R. 1994. A population study of the European eel (*Anguilla anguilla* (L.)) in the
16 Burrishoole system, Ireland, with special reference to growth and movement.
17 Dublin University, Dublin, Ireland: 416pp.
- 18 Poole, W. R., and Reynolds, J. D. 1996. Growth rate and age at migration of *Anguilla anguilla*.
19 *Journal of Fish Biology*, 48; 633-642.

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

- 1 Poole, W. R., Reynolds, J. D., and Moriarty, C. 1990. Observations on the silver eel migrations
2 of the Burrishoole river system, Ireland, 1959 to 1988. *Internationale Revue der*
3 *gesamten Hydrobiologie*, 75: 807-815.
- 4 R Core Team (2016). R: A language and environment for statistical computing. R Foundation
5 for Statistical Computing, Vienna, Austria. URL <http://www.R-project.org/>
- 6 Rosell, R., Evans, D., and Allen, M. 2005. The eel fishery in Lough Neagh, Northern Ireland -
7 an example of sustainable management? *Fisheries Management and Ecology*, 12:
8 377-385.
- 9 Rossi, R. 1979. An estimate of the production of the eel population in the Valli of Comacchio
10 (Po Delta) during 1974–1976. *Bolletino di zoologia*, 46: 217-223.
- 11 Rossi, R., and Cannas, A. 1984. Eel fishing management in a hypersaline lagoon of southern
12 Sardinia. *Fisheries Research*, 2: 285-298.
- 13 Rossi, R., Carrieri, A., Franzoi, P., Cavallini, G., and Gnes, A. 1988. A study of eel (*Anguilla*
14 *anguilla* L.) population dynamics in the Comacchio lagoons (Italy) by mark-recapture
15 method. *Oebalia*, 14: 87-106.
- 16 Sakamoto, Y., Ishiguro, M., and Kitagawa G. 1986. *Akaike Information Criterion Statistics*.
17 Dordrecht: D. Reidel Publishing Company.
- 18 Sandlund, O. T., Diserud, O. H., Poole, R., Bergersen, K., Dillane, M., Rogan, G., Durif, C.,
19 Thorstad, E. B., and Vøllestad, L. A. 2017. Timing and pattern of annual silver eel

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

- 1 migration in two European watersheds are determined by similar cues. Ecology and
2 Evolution, 7: 5956-5966.
- 3 Tesch, F. W. 2003. The Eel, Blackwell Science Ltd, London.
- 4 Van Ginneken, V., Durif, C., Balm, S. P., Boot, R., Verstegen, M. W. A., Antonissen, E., and Van
5 den Thillart, G. 2007. Silvering of European eel (*Anguilla anguilla* L.): seasonal
6 changes of morphological and metabolic parameters. Animal Biology, 57: 63-77.
- 7 Vøllestad, L. A. 1992. Geographic variation in age and length at metamorphosis of maturing
8 European eel: Environmental effects and phenotypic plasticity. Journal of Animal
9 Ecology, 61: 41-48.
- 10 Vøllestad, L. A., and Jonsson, B. 1986. Life-history characteristics of the European eel
11 *Anguilla anguilla* in the Imsa River, Norway. Transactions of the American Fisheries
12 Society, 115: 864-871.
- 13 Vøllestad, L. A., Jonsson, B., Hvidsten, N. A., Næsje, T. F., Haraldstad, Ø., and Ruud-Hansen, J.
14 1986. Environmental factors regulating the seaward migration of European silver
15 eels (*Anguilla anguilla*). Canadian Journal of Fisheries and Aquatic Sciences, 43:
16 1909-1916.
- 17 Vøllestad, L. A., and Jonsson, B. 1988. A 13-year study of the population dynamics and
18 growth of the European eel *Anguilla anguilla* in a Norwegian river: Evidence for

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. ICES Journal of Marine Science 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

density-dependent mortality, and development of a model for predicting yield.

Journal of Animal Ecology, 57: 983-997.

Walker, A. M., Andonegi, E., Apostolaki, P., Aprahamian, M., Beaulaton, L., Bevacqua, D.,

Briand, C., *et al.* 2011. Lot 2: Pilot project to estimate potential and actual

escapement of silver eel. Final project report, Service Contract S12.539598, Studies

and Pilot Projects for Carrying out the Common Fisheries Policy, European

Commission Directorate General for Maritime Affairs and UK EMP Progress Report

2015: 125 pp.

Wei, W. W. S. 2006. Time Series Analysis: Univariate and Multivariate Methods. 2nd Edition,

Pearson Education Inc.: 614pp

Wickström, H., Westin, L., and Clevestam, P. 1996. The biological and economic yield from a

long-term eel-stocking experiment. Ecology of Freshwater Fish, 5: 140-147.

Wolf, P. 1951. A trap for the capture of fish and other organisms moving downstream.

Transactions of the American Fisheries Society, 80: 41-45.

Figure Captions

Figure 1. Map of northwestern Europe, showing the location of the two study sites (stars),

Burrishoole in Ireland and Imsa in Norway.

This is a pre-copiedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

Figure 2. Mean winter, annual and summer water temperatures (upper graph), and water temperature anomalies with reference to the 1971-2000 period for the Burrishoole (Burr) and Imsa (1975-2000) (lower graph). In the lower graph, bars indicate annual water temperature anomalies and lines 5-year moving averages.

Figure 3. Annual counts of eels emigrating from the Burrishoole and Imsa Rivers between the early 1970s and 2015, with means for each period indicated by dashed lines (Burrishoole in blue, Imsa in red). The breakpoints are indicated by vertical dotted grey lines and the 95% confidence intervals around the breakpoints are shown above the x-axis.

Figure 4. Model run predicted outputs plotted against detrended silver eel counts for the Burrishoole (left) and Imsa (right) catchments. The straight line gives the 1-1 relationship.

Figure 5. Length frequency distributions for Burrishoole 1988 (n = 3003) and Imsa 1982-1988 (n = 1830) (upper graph) and for 2012-2015 Burrishoole (n = 5665) and Imsa (n = 1720) (lower graph).

Figure 6. Boxplots for length (cm) of male silver eels in Burrishoole. The central box depicting the middle half of the data (25th-75th percentiles), the horizontal line across the box marks

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

the median, the whiskers indicate the main extent of the connected data ($\sim 1.5 \times \text{IQR}$) and extreme values are indicated by starbursts.

Figure 7. Boxplots for length (cm) of female silver eels in Burrishoole and Imsa. The central box depicting the middle half of the data (25th-75th percentiles), the horizontal line across the box marks the median, the whiskers indicate the main extent of the connected data ($\sim 1.5 \times \text{IQR}$) and extreme values are indicated by starbursts.

Figure 8. Production in numbers (left panel) and biomass (right panel) per hectare for the Burrishoole and Imsa catchments, averaged for the periods before and after their respective change in the 1980s (1982 Burrishoole; 1988 Imsa). The black bars indicate the 95% confidence intervals.

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).

1 Table Captions

2 Table 1. Significant variables and their coefficients and significance levels for the Burrishoole
3 and Imsa run models. T = Water Temperature ($^{\circ}\text{C}$), WL = Water Level (m in Imsa,
4 standardised lake water level in Burrishoole), t = year and $t-1$ = preceding year.

5

6 Table 2. Five year mean total biomass (kg), % males and production in weight and number
7 per hectare of the total run of silver eels, and separately for males and females in the
8 Burrishoole and Imsa Rivers. Standard deviations are in brackets. Blanks signify absent data
9 and * is for 1976 only.

10

This is a pre-copyedited, author-produced version of an article accepted for publication in ICES Journal of Marine Science following peer review. The version of record Poole, W. Russell; Diserud, Ola Håvard; Thorstad, Eva Bonsak; Durif, Caroline; Dolan, Conor; Sandlund, Odd Terje; Bergesen, Knut Aanestad; Rogan, Gerard; Kelly, Sean D.; Vøllestad, Leif Asbjørn.

Long-term variation in numbers and biomass of silver eels being produced in two European river systems. *ICES Journal of Marine Science* 2018 [insert complete citation information here] is available online at: 20180509 <http://www.ices.dk/products/icesjournal.asp> [10.1093/icesjms/fsy053](https://doi.org/10.1093/icesjms/fsy053).