



The Acute and Delayed Mortality of the Northern Krill (*Meganyctiphanes norvegica*) When Exposed to Hydrogen Peroxide

Rosa H. Escobar-Lux¹ · Ole B. Samuelsen²

Received: 2 July 2020 / Accepted: 14 September 2020 / Published online: 26 September 2020
© The Author(s) 2020

Abstract

Bath treatment pharmaceuticals used to control sea lice infestations in the salmonid industry, such as hydrogen peroxide (H_2O_2), are released directly into the environment where non-target organisms are at risk of exposure. The aim of this study was to determine the threshold concentrations for mortality of the Northern krill, *Meganyctiphanes norvegica*, a major component of the north Atlantic marine ecosystem. To assess the lethal effects of H_2O_2 , we carried out a series of 1 h acute toxicity tests and assessed mortality through a 48 h post-exposure period. One-hour exposure to 170 mg/L, corresponding to 10% of the recommended H_2O_2 treatment, caused 100% mortality and a subsequent acute median-lethal concentration LC_{50} value of 32.5 mg/L. Increased mortality was observed with time in all exposed groups, resulting in successively lower LC_{50} values during the post-exposure period. The suggested H_2O_2 concentrations have the potential of causing negative effects to the Northern krill.

Keywords Crustacean · Toxicity · LC_{50} · Aquaculture

Sea lice (*Lepeophtheirus salmonis* and *Caligus rogercresseyi*), naturally occurring parasitic copepods affecting both farmed and wild salmonid populations, are a major challenge for the salmonid industry worldwide (Costello 2006; Torrissen et al. 2013; Vollset et al. 2016). The parasites feed on the mucous, skin, and blood of its host, and if present in significant numbers they can cause damage associated with osmotic stress and secondary infections (Finstad et al. 2000; Johnson et al. 2004; González et al. 2015). Norwegian wild salmonid populations, migrating post smolts from Atlantic salmon and local populations of sea trout (*Salmon trutta*), can suffer high mortality if there is high density of salmon lice larvae in the surrounding water (Costello 2009; Vollset et al. 2016). In farmed fish, salmon lice infestations reduce the general welfare of the fish and lead to an increase of the overall cost of the industry due to reduced growth and marketability due to skin lesions, and high costs associated with delousing treatments (Costello 2009). Therefore, both

the economic and ecological impact of salmon lice infestations are significant challenges for the salmonid industry.

In order to control salmon lice infestations, the industry has relied on the use of different chemotherapeutants, through the application of bath treatments and the use of in-feed drugs. Bath treatments can be applied either by enclosing the fish cages with an impervious tarpaulin or transferring the fish into well-boats, and after treatment the waste water is directly released into the surrounding water (Ernst et al. 2001; Burrige et al. 2010). At a global level, hydrogen peroxide (H_2O_2) was introduced as an antiparasitic agent after the loss of sensitivity in both *L. salmonis* and *C. rogercresseyi* to other delousing agents (Bravo et al. 2015; Urbina et al. 2019). In Norway alone, H_2O_2 is still the most used bath treatment therapeutant with a consumption of 4523 tons in 2019 (www.fhi.no/hn/legemiddelbruk).

Hydrogen peroxide acts on salmon lice by hydroxyl radicals attacking lipid and cellular organelles resulting in inactivation of enzymes and DNA replication (Cotran et al. 1989; Urbina et al. 2019). Previous studies have also shown that decomposition of hydrogen peroxide to water and O_2 bubbles in the gut and the haemolymph may cause mechanical paralysis leading to detachment of the pre-adult and adult salmon lice from the fish and causing them to float towards the surface (Bruno and Raynard 1994;

✉ Rosa H. Escobar-Lux
rosa.escobar@hi.no

¹ Institute of Marine Research, Austevoll Research Station, Sauganeset 16, 5392 Storebø, Norway

² Institute of Marine Research, Nordnes, P.O. Box 1870, 5817 Bergen, Norway

Aaen et al. 2014). A bath treatment involves the release of a large volume of H₂O₂ containing waste water and the chemical can potentially be dispersed over a wide area (Burrige et al. 2010, 2014, Parsons et al. 2020, Refseth et al. 2017). Therefore, there is a growing concern about the possible toxic effects of H₂O₂ on non-target aquatic invertebrate species living in the vicinity of fish farms, and specifically crustaceans which has been proven as particularly vulnerable (Smit et al. 2008; Burrige et al. 2014; Van Geest et al. 2014; Gebauer et al. 2017; Hansen et al. 2017; Bechmann et al. 2019; Escobar-Lux et al. 2019).

The pelagic zooplankton, *Meganyctiphanes norvegica*, Northern krill, is a species at risk as its distribution overlaps with the location to many salmon farms in Norway, as it inhabits both coastal and offshore waters (Kaartvedt et al. 2002; Melle et al. 2004; Tarling et al. 2010). Furthermore, the distribution of this boreal krill species has been described to be seasonal, with a predominant coastal distribution between the months of January and May (Grover 1952). In Norway, during this period of the year, pharmaceuticals are being used to keep the level of salmon lice below 0.2 female lice per fish as specified in the Norwegian Ministry of Trade, Industry and Fisheries (FOR-2012-12-05-1140, 2012) (Grefsrud et al. 2019). The total biomass of euphasiid stocks in the Norwegian Sea has been previously estimated to 42 million tons (Mt), with around 40–75% of this stock being Northern krill (Lindley 1982; Melle et al. 2004). Thus, the northern krill is a major component of the north Atlantic marine ecosystem, acting as a keystone organism between lower trophic levels and larger predators and plays an important role in the sequestration of carbon (Kaartvedt et al. 2005; Tarling et al. 2010). It is preyed upon by several commercially important fish species (Sameoto et al. 1994; Onsrud et al. 2004), seabirds (Montevecchi et al. 1992; Stevick et al. 2008), and marine mammals (Brodie et al. 1978). Moreover, the commercial exploitation of Northern krill is gaining interest in the salmonid industry as a potential protein alternative to the fishmeal (Tarling et al. 2010). Mass death of krill washed up on a beach can occur and is considered a natural phenomenon. Previously the mass stranding of *M. norvegica* has been explained as predation events in which predators' chase krill ashore (MacDonald 1927), transported to land by oceanic currents or by special events like upwellings (Aitken 1960; Cox 1975), or because special lightning conditions that might interfere with the krill's behavior (Wiborg 1966). However, in recent years there has been a higher frequency of reports in Norway describing this phenomenon near areas with salmon farms. This started a debate in public media of what might have caused the mass mortality and one of the most frequently cited suggestions has been the use of pesticides for delousing of the salmon farms, and especially H₂O₂. However, the

effects of H₂O₂ exposure on the Northern krill have until now been unknown.

For treating salmon, the recommended concentration for a H₂O₂ bath treatment is 1500–2100 mg/L for 20 min depending on temperature (<https://www.felleskatalogen.no/medisin-vet>). Typically, toxicity studies use exposure times that vary from 24 to 96 h. However, these may not be representative of the real-life scenarios following a release of waste water after a bath treatment on a salmon farm (Ernst et al. 2001; Urbina et al. 2019). The use of 1 h exposures, is considered a more realistic exposure scenario, but to date only a limited number of species have been tested under those conditions (Medina et al. 2004; Fairchild et al. 2010; Burrige et al. 2014; Van Geest et al. 2014; Escobar-Lux et al. 2019; Parsons et al. 2020). What these previous studies also have shown is that the mortality observed immediately after exposure tends to be lower than the mortalities registered if a post-exposure period is included in the experimental set-up. A longer post-exposure observation period is therefore recommended.

The main objective of this study was to examine the toxicity of H₂O₂ to *M. norvegica*, a non-target crustacean and keystone species of the Norwegian marine environment. Our objective was to expose the Northern krill to a short 1 h pulse of H₂O₂ and assess the acute and delayed mortality during a post-exposure period of 48 h in clean seawater.

Materials and Methods

In the present study, krill (*M. norvegica*) were collected from the dock at Austevoll Research Station, Institute of Marine Research Norway (60° 05' 20" N 5° 15' 57" E) using light traps. The light traps (mesh size 500 µm; 0.45 m in diameter; BellaMare USA) were equipped with a white LED light and deployed at a depth of 20 m overnight. The research station is at least 3 km away from the nearest commercial salmon farm. Krill from the traps were transported to the laboratory at Austevoll Research Station and kept overnight in 10 L buckets supplied with sand filtered seawater from a depth of 160 m (Bjørnafjorden) holding a temperature of 8 °C (salinity of 34.2 ppt; pH 7.94). The experiment was performed within 48 h of capture and prior to exposure the krill were sorted and only krill in excellent physical condition were used in the experiments.

Commercial H₂O₂ (Nemona, Akzo Nobel Pulp and Performance Chemicals AB Sweden) at a concentration of 49.50% (600 g/L) was purchased from Akzo Nobel, Pulp and Performance Chemicals, AB Sweden. Since no previous studies had assessed the toxicity of H₂O₂ on *M. norvegica*, the chosen concentrations were based on the recommended dose for treating salmon (1700 mg/L). The krill were exposed to concentrations of 1.7, 8.5, 17, 170, 850 and

1700 mg/L H_2O_2 , corresponding to 0.1, 0.5, 1, 10, 50 and 100% of the recommended treatment dose. All exposures were conducted in glassware units with a volume of 500 mL. A total of 140 krill were randomly divided into seven treatment groups, including a control group, with four replicates for each treatment and each replicate counting five individuals. After the 1 h exposure, acute mortality was recorded and the krill were transferred to 10 L recovery tanks where mortality was checked successively at 6, 24 and 48 h post-exposure using a dissecting microscope. Krill were considered dead if there was no movement of the pereopods, pleopods or antenna after a gentle stimulus. Mortality that occurred during the 1 h exposure was defined as acute mortality. Total mortality was defined as the cumulative mortality after the 48 h post-exposure period.

The statistical analyses for mortality were done in the software R (Version 3.5.3 (2019-03-11) Copyright © 2019 The R Foundation for Statistical Computing). The LC_{50} values, and their 95% confidence intervals (CI), were calculated using generalized linear models (GLM) with binomial error structures and probit links, according to Finney (1971). Hydrogen peroxide concentrations were log₁₀ transformed to linearize the data.

Results and Discussion

This study clearly show that H_2O_2 was acutely toxic to wild-captured Northern krill *M. norvegica*. While no mortality was recorded in the group exposed to the lowest dose of 1.7 mg/L or in the control group, a 1 h exposure to 170 mg/L, i.e. 10% of recommended dose, caused 100% mortality and a subsequent acute LC_{50} value of 32.5 mg/L (16.8–48.2) was calculated (Fig. 1a). During the post-exposure period, increased mortality with time was observed in all exposed groups resulting in successively lower LC_{50} values with 14.11 mg/L after 6 h (7.3–20.9), 4.92 mg/L (1.2–7.9) after 24 h and finally 0.86 mg/L after 48 h (Fig. 1b–d). No mortality was registered in the control groups during the post-exposure period. The calculated LC_{50} value at 24 h represents a threefold dilution of the acute 1 h LC_{50} value. These findings clearly support the recommendations suggested in previous studies to include a post-exposure period following the exposure to H_2O_2 to assess any delayed effects (Van Geest et al. 2014; Brokke 2015; Escobar-Lux et al. 2019).

While several studies have examined the toxicity of H_2O_2 on marine crustacean species, the number of




Fig. 1 The toxicity of hydrogen peroxide to *M. norvegica* following 1 h exposure. Dose–response curves showing mortality amongst the northern krill at 0 h, 6 h, 24 h, and 48 h post-exposure to H_2O_2 . Each point on the graphs represent an individual replicate tank containing

4 to 6 krill and the line represent the best fit model for the data calculated using a binomial log-probit GLM in R. The shadowed area represents the 95% confidence intervals

studies using an exposure time of 1 h is more limited. A review of those studies reveals that some crustaceans

have a relatively high tolerance to H_2O_2 exposure and is reflected in low mortality when exposed to concentrations similar to or higher than the recommended treatment dose. This applies to both newly hatched larvae and adult of American lobster (*Homarus americanus*), sand shrimp (*Crangon septemspinosa*), the mysid *Mysid* sp. (Burridge et al. 2014), rock pool shrimp (*Palaemon elegans*) and chameleon shrimp (*Praunus flexuosus*) (Brokke 2015). For some species, low mortality was observed even when a post-exposure period was included in the study. Following an exposure of 1 h and a 95 h post-exposure period, the calculated LC_{50} values were 1673 mg/L for *H. americanus* larvae, > 3750 mg/L for adult American lobster, 3182 mg/L for sand shrimps and 973 mg/L for *Mysid* sp. (Burridge et al. 2014; Van Geest et al. 2014). For rock pool shrimps and chameleon shrimps the acute mortality after 1 h exposure was low indicating LC_{50} values higher than the highest exposure concentration of 1700 mg/L for both species (Brokke 2015). However, a significant mortality occurred during the 24 h post-exposure period, resulting in LC_{50} values of 174.1 mg/L and 77.5 mg/L for rock pool shrimp and chameleon shrimps respectively, classifying these species as highly sensitive. In the study by Bechmann et al. (2019), the Northern shrimp (*Pandalus borealis*) was exposed to 15 mg/L H_2O_2 for 1 h. The very low acute mortality observed immediately after exposure did however increase during the post-exposure period (7 days) but as the total mortality never exceeded 30%, no LC_{50} could be calculated. Damage on the gills was observed in the shrimps exposed to H_2O_2 and suggested as the major cause of the delayed mortality (Bechmann et al. 2019).

In comparison, species like the copepods *Acartia Hudsonica* and *Calanus* spp. have shown higher sensitivity to H_2O_2 exposure, resulting in EC_{50} and LC_{50} values of 2.6–10 mg/L and 30.6 mg/L respectively, following a 24 h post-exposure period (Van Geest et al. 2014; Escobar-Lux et al. 2019). In the case of the European lobster (*Homarus gammarus*) larvae (stage I–IV), a 1 h exposure to 1530 mg/L followed by a 24 h post-exposure period, resulted in mortalities between 75 and 100% (Escobar-Lux et al. 2020) and calculated LC_{50} values of 177 mg/L, 404 mg/L, 676 mg/L and 738 mg/L, for stages I, II, III and IV respectively. For species other than crustaceans, the polychaete *Ophryotrocha* sp. and the sugar kelp *Saccharina latissima* are amongst the more sensitive marine species with LD_{50} values of 64.3 mg/L and 80.7 mg/L following 72 h and 7 days' post-exposure periods, respectively (Fang et al. 2018; Haugland et al. 2019). The LC_{50} values calculated for northern krill are therefore, to our knowledge the most sensitive species examined so far.

This study has shown that a bath treatment with H_2O_2 has a detrimental effect on *M. norvegica*. However, it is important to assess whether these laboratory-based concentrations are likely to pose a significant risk to krill at

the proximity of salmonid aquaculture sites. Due to differences in experimental set-ups the variation in half-lives reported for H_2O_2 in seawater in large, with results between 1 and 58 days (Bruno and Raynard 1994; Lyons et al. 2014; Fagereng 2016; Parsons and Samuelson unpubl. data). Several factors affect both the toxicity and the degradation of H_2O_2 , for example the water temperature or the irradiance (Stratford et al. 1984; Treasure et al. 2000)". However, even the shortest degradation time reported (1 day) is significantly longer than the 1 h exposure needed in the present study to cause considerable mortality of the Northern krill. Even though H_2O_2 is extensively used around the world as an anti-sea lice bath treatment, few studies have initiated the use of mathematical models to predict its' dispersal and its' impact on non-target species. One such study from Norway has indicated that the spread of H_2O_2 may be larger than previously thought (Refseth et al. 2017). According to the model, concentrations up to 300 mg/L may occur within a 1 km radius from the farm and 100 mg/L within a radius of 2 km. Furthermore, the model also suggested that a concentration of 100 mg/L can be present in surface waters for several hours after discharge. The presented model simulations therefore suggest that the Northern krill within 2 km of a salmonid farm may be exposed to lethal concentration of H_2O_2 .

Parsons et al. (2020) used dispersion models to predict the spreading of pharmaceuticals from salmonid farms in Norway, following bath treatment. Based on the models and LC_{50} values (1 h exposure followed by 24 h post-exposure period) for European lobster larvae (stage I and II) they calculated impact zones around 23 Norwegian fish farms for the pesticides azamethiphos and deltamethrin. This model however, did not take into account the degradation of the compounds due to the presence of organic matter in the water. While the azamethiphos impact zones around farms were relatively small (mean area of 0.04–0.2 km²), deltamethrin impact zones covered much larger areas (mean area of 21.1–39.0 km²). The difference in impact zone is due to the difference in toxicity between the two drugs. For azamethiphos the 1 h- LC_{50} values (95% CIs) for stage I and II larvae were 43.1 µg/L (13.0–131.0 µg/L) and 20.5 µg/L (13.2–30.9 mg/L), respectively, representing approximately 2- and fivefold dilutions of the treatment concentration (100 µg/L) used on Norwegian fish farms. For deltamethrin the 1 h- LC_{50} values (with 95% CIs) for stage I and II larvae were estimated to be 2.6 ng/L (0.6–11.0 ng/L) and 2.9 ng/L (1.5–5.7 ng/L), representing approximately 800-fold dilution of the treatment concentration of 2000 ng/L. Considering the sensitivity of krill towards H_2O_2 found in the present study, where the LC_{50} ranged from 52- to 2000-fold dilution with increasing post-exposure period, impact zones like those

calculated for deltamethrin in Parsons et al. (2020) will be most relevant for impact zones for H₂O₂ and krill.

Meganyctiphanes norvegica can be found around the North Atlantic, with the Norwegian sea being a major hotspot for its distribution (Melle et al. 2004). Due to their distribution, krill can often be found in waters close to aquaculture sites and therefore be negatively impacted by the dispersal of effluent plumes after treatments. Based on our findings and the information from previous mathematical models, H₂O₂ may cause a larger impact than it was previously believed. Therefore, that some cases of mass mortality of krill observed in past years may have been caused by H₂O₂ exposure, cannot be overlooked.

Acknowledgements This work was funded by the Norwegian Institute of Marine Research /Havforskningsinstituttet internal funding (Project # 14907) led by O.B.S. We thank all staff at IMR Austevoll for their assistance with these experiments, in particular Florian Freydet.

Funding Open Access funding provided by Institute Of Marine Research.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

- Aaen SM, Aunsmo A, Horsberg TE (2014) Impact of hydrogen peroxide on hatching ability of egg strings from salmon lice (*Lepeophtheirus salmonis*) in a field treatment and in a laboratory study with ascending concentrations. *Aquaculture* 422:167–171
- Aitken JJ (1960) Swarming in *Meganyctiphanes norvegica* (M. Sars) in Strangford Lough, Co. Down. *Ir Nat J* 13:140–142
- Bechmann RK, Arnberg M, Gomiero A, Westerlund S, Lyng E, Berry M, Agustsson T, Jager T, Burrige LE (2019) Gill damage and delayed mortality of Northern shrimp (*Pandalus borealis*) after short time exposure to anti-parasitic veterinary medicine containing hydrogen peroxide. *Ecotoxicol Environ Saf* 180:473–482
- Bravo S, Silva MT, Agusti C, Sambra K, Horsberg TE (2015) The effect of chemotherapeutic drugs used to control salmon louse on the hatching viability of egg strings from *Caligus rogercresseyi*. *Aquaculture* 443:77–83
- Brodie PF, Sameoto DD, Sheldon RW (1978) Population densities of euphausiids off Nova Scotia as indicated by net samples, whale stomach contents, and sonar. *Limnol Oceanogr* 23(6):1264–1267
- Brokke KE (2015) Mortality caused by de-licing agents on the non-target organisms chameleon shrimp (*Praunus flexuosus*) and grass prawns (*Palaemon elegans*). M.Sc. Thesis, University of Bergen, Bergen, Norway
- Bruno DW, Raynard RS (1994) Studies on the use of hydrogen peroxide as a method for the control of sea lice on Atlantic salmon. *Aquacult Int* 2(1):10–18
- Burrige L, Weis JS, Cabello F, Pizarro J, Bostick K (2010) Chemical use in salmon aquaculture: a review of current practices and possible environmental effects. *Aquaculture* 306(1–4):7–23
- Burrige LE, Lyons MC, Wong DKH, MacKeigan K, VanGeest JL (2014) The acute lethality of three anti-sea lice formulations: AlphaMax®, Salmosan®, and Interlox® Paramove™ 50 to lobster and shrimp. *Aquaculture* 420:180–186
- Cotran RS, Kumar V, Robbins SL (1989) *Pathological basis of disease*, 4th edn. Saunders, Toronto
- Costello MJ (2006) Ecology of sea lice parasitic on farmed and wild fish. *Trends Parasitol* 22(10):475–483. <https://doi.org/10.1016/j.pt.2006.08.006>
- Costello MJ (2009) The global economic cost of sea lice to the salmonid farming industry. *J Fish Dis* 32(1):115–118
- Cox SJ (1975) Shore stranding of *Meganyctiphanes norvegica* (M. Sars). *Estuar Coast Mar Sci* 3:483–484
- Ernst W, Jackman P, Doe K, Page F, Julien G, MacKay K, Sutherland T (2001) Dispersion and toxicity to non-target aquatic organisms of pesticides used to treat salmon louse on salmon in net pen enclosures. *Mar Pollut Bull* 42(6):432–443
- Escobar-Lux RH, Fields DM, Browman HI, Shema SD, Bjelland RM, Agnalt AL, Skiftesvik AB, Samuelsen OB, Durif CM (2019) The effects of hydrogen peroxide on mortality, escape response, and oxygen consumption of *Calanus* spp. *FACETS* 4(1):626–637
- Escobar-Lux RH, Parsons AE, Samuelsen OB, Agnalt AL (2020) Short-term exposure to hydrogen peroxide induces mortality and alters exploratory behaviour of European lobster (*Homarus gammarus*). *Ecotoxicol Environ Saf* 204:111111
- Fagereng MB (2016) Bruk Av Hydrogenperoksid i Oppdrettsanlegg; fortytningstudier Og Effekter På blomsterreke (*Pandalus montagui*). M.Sc. Thesis. University of Bergen, Norway, p. 104. <https://bora.uib.no/handle/1956/13008>. (In Norwegian)
- Fairchild WL, Doe KG, Jackman PM, Arsenault JT, Aubé JG, Losier M, Cook AM (2010) Acute and chronic toxicity of two formulations of the pyrethroid pesticide deltamethrin to an amphipod, sand shrimp and lobster larvae. *Can Tech Rep Fish Aquat Sci* 2876:34p
- Fang J, Samuelsen OB, Strand Ø, Jansen H (2018) Acute toxic effects of hydrogen peroxide, used for salmon lice treatment, on the survival of polychaetes *Capitella* sp. and *Ophryotrocha* spp. *Aquac Environ Interact* 10:363–368
- Finney DJ (1971) *Probit analysis*, 3rd edn. Cambridge University Press, Cambridge
- Finstad B, Bjørn PA, Grimnes A, Hvidsten NA (2000) Laboratory and field investigations of salmon lice [*Lepeophtheirus salmonis* (Krøyer)] infestation on Atlantic salmon (*Salmo salar* L.) post-smolts. *Aquac Res* 31(11):795–803
- Gebauer P, Paschke K, Vera C, Toro JE, Pardo M, Urbina M (2017) Lethal and sub-lethal effects of commonly used anti-salmon louse formulations on non-target crab *Metacarcinus edwardsii* larvae. *Chemosphere* 185:1019–1029
- Grefsrud ES, Svåsand T, Glover K, Husa V et al. (2019) Risikorapport norsk fiskeoppdrett 2019. Fisken og havet, særnr. 1–2019. Havforskningsinstituttet, Bergen. <https://www.hi.no/hi/nettrappporter/fisken-og-havet-2019-5>. (In Norwegian)
- Grover RS (1952) Continuous plankton records: the Euphausiacea of the Northeastern Atlantic and North Sea 1946–48: *Hull Bull. Mar Ecol* 3(23):185–214
- González MP, Marín SL, Vargas-Chacoff L (2015) Effects of *Caligus rogercresseyi* (Boxshall and Bravo, 2000) infestation on physiological response of host *Salmo salar* (Linnaeus 1758): Establishing physiological thresholds. *Aquaculture* 438:47–54

- Hansen BH, Hallmann A, Altin D, Jensen BM, Ciesielski M (2017) Acute hydrogen peroxide (H₂O₂) exposure does not cause oxidative stress in late-copepodite stages of *Calanus finmarchicus*. *J Toxicol Environ Health A* 80(16–18):820–829
- Haugland BT, Rastrick SP, Agnalt AL, Husa V, Kutti T, Samuelsen OB (2019) Mortality and reduced photosynthetic performance in sugar kelp *Saccharina latissima* caused by the salmon-lice therapeutant hydrogen peroxide. *Aquac Environ Interact* 11:1–17
- Johnson SC, Bravo S, Nagasawa K, Kabata Z, Hwang J, Ho J, Shih CT (2004) A review of the impact of parasitic copepods on marine aquaculture. *Zool Stud* 43(2):229–243
- Kaartvedt S, Larsen T, Hjelmseth K, Onsrud MS (2002) Is the omnivorous krill *Meganyctiphanes norvegica* primarily a selectively feeding carnivore? *Mar Ecol Prog Ser* 228:193–204
- Kaartvedt S, Røstad A, Fiksen Ø, Melle W, Torgersen T, Breien MT, Klevjer TA (2005) Piscivorous fish patrol krill swarms. *Mar Ecol Prog Ser* 299:1–5
- Lindley JA (1982) Population dynamics and production of euphausiids. *Mar Biol* 66(1):37–46
- Lyons MC, Wong DKH, Page FH (2014) Degradation of hydrogen peroxide in seawater using the anti-sea louse formulation interox paramove 50. Fisheries and Oceans Canada, Maritimes Region, St. Andrews Biological Station
- Macdonald R (1927) Food and habits of *Meganyctiphanes norvegica*. *J Mar Biol Assoc U K* 14(3):753–784
- Medina M, Barata C, Telfer T, Baird DJ (2004) Assessing the risks to zooplankton grazers of continuous versus pulsed cypermethrin exposures from marine cage aquaculture. *Arch Environ Contam Toxicol* 47(1):67–73
- Melle W, Ellertsen B, Skjoldal HR (2004) Zooplankton: the link to higher trophic levels. The Norwegian Sea ecosystem. Tapir Academic Press, Trondheim, pp 137–202
- Montevicchi WA, Birt-Friesen VL, Cairns DK (1992) Reproductive energetics and prey harvest of Leach's storm-petrels in the north-west Atlantic. *Ecology* 73(3):823–832
- Onsrud MSR, Kaartvedt S, Røstad A, Klevjer TA (2004) Vertical distribution and feeding patterns in fish foraging on the krill *Meganyctiphanes norvegica*. *ICES J Mar Sci* 61(8):1278–1290
- Parsons AE, Escobar-Lux RH, Sævik PN, Samuelsen OB, Agnalt AL (2020) The impact of anti-sea lice pesticides, azamethiphos and deltamethrin, on European lobster (*Homarus gammarus*) larvae in the Norwegian marine environment. *Environ Pollut*. <https://doi.org/10.1016/j.envpol.2020.114725>
- Refseth GH, Sæther K, Drivdal M, Nøst OA, Augustine S, Camus L, Tassara L, Agnalt A-L, Samuelsen OB (2017) Miljørisiko ved bruk Av Hydrogenperoksid. Økotoksikologisk vurdering Og Grenseverdi for Effekt. Assessment. CRC Press, Boca Raton. <https://www.fhf.no/prosjekter/prosjektbasen/901249/>. (In Norwegian)
- Sameoto D, Neilson J, Waldron D (1994) Zooplankton prey selection by juvenile fish in Nova Scotian Shelf basins. *J Plankton Res* 16(8):1003–1019
- Smit MG, Ebbens E, Jak RG, Huijbregts MA (2008) Time and concentration dependency in the potentially affected fraction of species: the case of hydrogen peroxide treatment of ballast water. *Environ Toxicol Chem* 27(3):746–753
- Stevick PT, Incze LS, Kraus SD, Rosen S, Wolff N, Baukus A (2008) Trophic relationships and oceanography on and around a small offshore bank. *Mar Ecol Prog Ser* 363:15–28
- Stratford HK, Quimby PC, Ouzts JD (1984) Photo enhancement of hydrogen peroxide toxicity to submersed vascular plants and algae. *Aquat Plant Manage* 22:25–34
- Tarling GA, Ensor NS, Fregin T, Goodall-Copestake WP, Fretwell P (2010) An introduction to the biology of northern krill (*Meganyctiphanes norvegica* Sars). *Advances in marine biology*, vol 57. Academic Press, Cambridge, pp 1–40
- Torrissen O, Jones S, Asche F, Guttormsen A, Skilbrei OT, Nilssen F, Horsberg TE, Jackson D (2013) Salmon lice—impact on wild salmonids and salmon aquaculture. *J Fish Dis* 36(3):171–194
- Treasure JW, Gran A, Davi PJ (2000) Physical constraints of bath treatments of Atlantic salmon (*Salmo salar*) with a sea lice burden (Copepoda: Caligidae). *Contrib Zool* 69(1–2):129–136
- Urbina MA, Cumillaf JP, Paschke K, Gebauer P (2019) Effects of pharmaceuticals used to treat salmon lice on non-target species: evidence from a systematic review. *Sci Total Environ* 649:1124–1136
- Van Geest JL, Burrige LE, Fife FJ, Kidd KA (2014) Feeding response in marine copepods as a measure of acute toxicity of four anti-salmon louse pesticides. *Mar Environ Res* 101:145–152
- Vollset KW, Barlaup BT, Mahlum S, Bjørn PA, Skilbrei OT (2016) Estimating the temporal overlap between post-smolt migration of Atlantic salmon and salmon lice infestation pressure from fish farms. *Aquac Environ Interact* 8:511–525
- Wiborg KF (1966) Undersøkelser av krill (lyskreps) i Hardangerfjorden og tilstøtende områder, samt på stasjon M i Norskehavet. *Fiskets Gang* 41(754):761 (In Norwegian)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.