



Deformities in larvae and juvenile European lobster (*Homarus gammarus*) exposed to lower pH at two different temperatures

A.-L. Agnalt, E. S. Grefsrud, E. Farestveit, M. Larsen, and F. Keulder

Ann-Lisbeth Agnalt, Institute of Marine Research, P.O. Box 1870 Nordnes, 5817 Bergen, Norway

Correspondence to: A.-L. Agnalt (ann-lisbeth.agnalt@imr.no)

Received: 31 March 2013 – Published in Biogeosciences Discuss.: 2 May 2013

Revised: 28 October 2013 – Accepted: 30 October 2013 – Published: 4 December 2013

Abstract. The ongoing warming and acidification of the world's oceans are expected to influence the marine ecosystems, including benthic marine resources. Ocean acidification may especially have an impact on calcifying organisms, and the European lobster (*Homarus gammarus*) is among those species at risk. A project was initiated in 2011 aiming to investigate long-term effects of ocean acidification on the early life-cycle of lobster under two temperatures. Larvae were exposed to $p\text{CO}_2$ levels of ambient water (water intake at 90 m depth), medium 750 ($\text{pH} = 7.79$) and high 1200 μatm $p\text{CO}_2$ ($\text{pH} = 7.62$) at temperatures 10 and 18 °C. The water parameters in ambient water did not stay stable and were very low towards the end of the experiment in the larval phase at 10 °C, with pH between 7.83 and 7.90. At 18 °C, pH in ambient treatment was even lower, between 7.76 and 7.83, i.e. close to medium $p\text{CO}_2$ treatment. Long-term exposure lasted 5 months. At 18 °C the development from stage 1 to 4 lasted 14 to 16 days, as predicted under optimal water conditions. Growth was very slow at 10 °C and resulted in three larvae reaching stage 4 in high $p\text{CO}_2$ treatment only.

There were no clear effects of $p\text{CO}_2$ treatment, on either carapace length or dry weight. However, deformities were observed in both larvae and juveniles. The proportion of larvae with deformities increased with increasing $p\text{CO}_2$ exposure, independent of temperature. In the medium treatment about 23 % were deformed, and in the high treatment about 43 % were deformed. None of the larvae exposed to water of $\text{pH} > 7.9$ developed deformities. Curled carapace was the most common deformity found in larvae raised in medium $p\text{CO}_2$ treatment, irrespective of temperature, but damages in the tail fan occurred in addition to a bent rostrum. Curled carapace was the only deformity found in high $p\text{CO}_2$ treatment at both temperatures. Occurrence of deformities after

five months of exposure was 33 and 44 % in juveniles raised in ambient and low $p\text{CO}_2$ levels, respectively, and 21 % in juveniles exposed to high $p\text{CO}_2$. Deformed claws were most often found in ambient and medium treatment (56 %), followed by stiff/twisted walking legs (39 %) and puffy carapace (39 %). In comparison, at high $p\text{CO}_2$ levels 71 % of the deformed juveniles had developed a puffy carapace. Overall, about half of the deformed juveniles from the ambient and medium $p\text{CO}_2$ treatment displayed two or three different abnormalities; 70 % had multiple deformities in the high $p\text{CO}_2$ treatment. Some of the deformities in the juveniles may affect respiration (carapace), the ability to find food, or sexual partners (walking legs, claw and antenna), and ability to swim (tail-fan damages).

1 Introduction

The world's atmosphere is increasingly becoming more saturated with the concentration of CO_2 as carbon emissions from burning fossil fuels keep increasing (IPCC, 2007; Caldeira et al., 2005). Atmospheric CO_2 is currently around 380 ppmv, but is predicted to increase to 780 ppmv and 1200 ppmv by 2100 and 2200, respectively (IPCC, 2007). Increased absorption of atmospheric CO_2 into the marine environment leads to an increase in total dissolved inorganic carbon (DIC), which changes the chemistry and acid-base balance and results in a decreased seawater pH (Dickson et al., 2007). Currently the global average seawater pH is about 8.05 units, and is associated with a DIC of 2026 (Fabry et al., 2008). It is predicted to drop by 0.3 to 0.4 pH units by 2100 (Orr et al., 2005; Feely et al., 2009). At the same time, depending on the emission scenarios, ocean temperature in

the upper 100 m is predicted to increase by 0.6° (RCP2.6) to 2 °C (RCP8.5) by 2100, (Blunden and Arndt, 2013). The absorption of CO₂ by the ocean and ocean acidification (OA) occurs mostly in the upper 100 m, and varies with latitude and temperature (Orr et al., 2005; Fabry et al., 2008).

Regional monitoring of DIC at Station M in the eastern Norwegian Sea shows a value of 2140 (Skjelvan et al., 2008), which is above the global average of 2026. This agrees with recent modeling that shows CO₂ uptake is higher and pH lower in the eastern Norwegian Sea (Olsen et al., 2006). pH levels will decrease with water depth at which aragonite becomes undersaturated, from ~ 100 to 50 m by 2100 at high latitudes (Fabry et al., 2008). Marine organisms in colder regions are therefore at a greater risk of being affected by the effects of both warming and OA, especially calcifying marine organisms. One such species is an important predator, the European lobster *Homarus gammarus*.

H. gammarus is found along the continental shelf in the northeast Atlantic, which extends from the warm waters off Morocco to the colder areas near the Arctic Circle, i.e. Tysfjord and Nordfolda (68° N) (Agnalt et al., 2009). This distribution covers a large latitudinal and temperature range. With CO₂ absorption varying greatly with latitude and temperature, the effects of acidification may vary greatly for subpopulations across the distribution range. The life cycle of *H. gammarus* consists of four larval stages, a juvenile stage, a sub-adult stage (~ 50 mm carapace length: CL) and adult of > 60 mm CL (Factor, 1995); larvae are pelagic in the first three stages (stage 1–3), after which they settle during larval stage 4. Little is known about the benthic stages of *H. gammarus* juveniles less than 40 mm CL in the wild (Linnane et al., 2001). The European lobster is esteemed as a valuable marine resource and has supported the coastal fishery in northern Europe for several centuries (Agnalt, 2008). In the 1960s, lobster populations in Norway were depleted below sustainable levels and due to low recruitment, the recovery has been slow. Any additional factors that reduce recruitment and population size further may push this species to the brink of extinction in these areas.

Only one study has investigated the effects of OA (*p*CO₂) in *H. gammarus*, focusing on larvae at the predicted future scenario of *p*CO₂ 1200 µatm (Arnold et al., 2009). Calcification was significantly reduced in stages 3 and 4, with no direct effect on growth observed. The decrease in calcification observed in stage 4 at a *p*CO₂ of 1200 µatm may have been due to energy being channelled towards growth and possibly acid-base regulation (Arnold et al., 2009). Growth in crustaceans requires replacement of the old exoskeleton by a new and larger exoskeleton, in a process called moulting, occurring more frequently in juveniles compared with adults. Moulting is highly temperature dependent, making juveniles vulnerable to temperature changes (Waddy et al., 1995). Moulting also involves depositing CaCO₃ to harden the shell, which is energetically costly and therefore puts great physiological stress on the animal. Low pH resulting

from OA increases physiological stress and may be devastating to moulting juveniles already under metabolic stress. Warmer temperatures, predicted to co-occur with increased OA, may have its added metabolic stress on juvenile lobsters (as seen in the crab *Hyas araneus*) if thermal tolerance limits are exceeded (Walther et al., 2010). Ries et al. (2009) investigated the impact of OA on a range of benthic marine calcifiers and found that *Homarus americanus* juveniles had the highest net calcification at high *p*CO₂ levels (2856 ppm). The increase in calcification is thought to be initiated by actively increasing pH at the calcifying centres, therefore reducing H⁺ and converting bicarbonate (HCO₃⁻) to carbonate (CO₃⁻²). CO₂-induced acidosis in crustaceans is usually compensated for by increasing bicarbonate production (Truchot, 1978; Pörtner et al., 2004; Spicer et al., 2007). Bicarbonate production is energetically costly and may be reduced over the long term if energy reserves are low.

The one study on OA in *H. gammarus* and studies concerning other crustacean species suggest different impacts occurring in different stages of the life cycle. The synergistic effects of warming and OA on the life cycle of *H. gammarus* (and other lobster species) are unknown and urgently need to be studied. A project was therefore initiated in 2011 aiming to investigate long-term synergistic effects of temperature and projected increases in ocean acidification on the early life cycle of lobster, i.e. the larval and juvenile phase of *H. gammarus*. Here we use 18 °C, similar temperatures to Arnold et al. (2009), which are optimal for hatching and growth (Wickins and Lee, 2002; Kristiansen et al., 2004) and 10 °C which are less optimal conditions (Schmalenbach and Buchholz, 2013). The temperatures chosen are not a likely climate change scenario but a physiological test.

2 Material and methods

Experiments combining OA with temperature were conducted at IMR-Matre (60°52' N, 05°35' E) over a period of five months, lasting from 28 September 2011 to 22 March 2012. Raw water was pumped from 90 m depth, thus representing ambient water. Each experimental unit consisted of a 400 L tank of 0.87 × 0.87 × 0.53 m. Six of these units were used for the ambient, medium (750 µatm) and high (1200 µatm) *p*CO₂ treatment run at two temperatures (10 °C and 18 °C). For the larval rearing, two 40 L incubators (Hughes kreisel) were placed in each unit (12 in total, Fig. 1a). For on-growing, the juveniles were kept individually in trays, placed in the experimental units (Fig. 1b). From 5 months of age, the surviving juveniles were continuously monitored but under ambient conditions at 14 °C, until 25 October 2012, by which time they were about 1 year old. The larvae and juveniles were all kept in 8 to 10 h light and 16 to 18 h dark.

2.1 Water parameters

The salinity during the course of the experiment was on average 33.7 ± 0.2 ppt. Temperature was run at 10 and 18 °C, to simulate a lower limit and an optimum threshold for homarid lobsters, respectively (Wickins and Lee, 2002; Kristiansen et al., 2004; Schmalenbach and Buchhoz, 2013). Initially a middle temperature of 14 °C was also included, but due to the limited number of larvae available this had to be excluded. The experiments were run in ambient water, believed to be at a current $p\text{CO}_2$ of 380 μatm occurring in the natural oceanic environment; treatments with a medium $p\text{CO}_2$ of approximately 750 μatm and with a high $p\text{CO}_2$ aiming at 1200 μatm , to simulate low and medium OA scenarios predicted for 2100 and 2200, respectively. Multiple 2×2 AGA 25 kg gas bottles were used to induce CO_2 into two cones, creating seawater with $\text{pH} = 6$. From each of these cones, CO_2 -rich water was mixed with oxygen-rich seawater and the flow rates from the cones were regulated by controllers to obtain the desired seawater quality; $\text{pH} = 7.79$ in medium $p\text{CO}_2$ treatment and $\text{pH} = 7.62$ in high $p\text{CO}_2$ treatment. Water from the enclosures were supplied to and circulated through the experimental tanks. Temperature, pH and oxygen were monitored continuously in each experimental tank by probes connected to a computer. The pH was measured several times daily using Orbisint CPS11D from Endress + Hauser electrodes, using the National Bureau of Standards (NBS) scale. pH was calibrated once a week with buffers of $\text{pH} = 4.0$ and 7.0 . In addition, calibration was also made by using a spectrophotometry Hitachi U-2900 connected to a Refrigerated Heating Circulator Julabo F12 combined with temperature sensors TD301A from SAIV A/S. These measurements deviated from 0.001 to 0.053 from pH_{NBS} and were considered acceptable for this study. Samples for analysis of CO_2 content in seawater in μatm , total alkalinity (A_{T}) and total dissolved inorganic carbon (C_{T}) were collected in 350 ml amber glass bottles with minimal headspace. 300 μL of saturated HgCl_2 solution was added to preserve the sample. Aragonite and calcite saturation state (Ω) and pH on the total hydrogen ion scale (pH_{T}) were then calculated.

In the medium $p\text{CO}_2$ treatment the content of $p\text{CO}_2$ was 727 ± 12 μatm , and correspondingly 1198 ± 157 μatm in the high $p\text{CO}_2$ treatment (Table 1), combined for both temperatures. The water quality in ambient water, i.e. intake of raw water at 90 m depth, did not stay stable. At 10 °C, pH in the larval experiments was stable at an average 8.01 from 28 September until 11 November, then dropped to 7.92 and even dropped further and was between 7.83 and 7.90 towards the end of the experiment (Fig. 2). In ambient water at 18 °C the pH was relatively stable but at 7.80 ± 0.02 throughout the larval phase. In other words, pH was above 7.9 only in two-thirds of the larval phase at 10 °C. pH was relatively stable in the experiments run at medium and high $p\text{CO}_2$ at both temperatures, however with a few outliers (Fig. 2). In the juvenile experiments $p\text{CO}_2$ in ambient water was higher than

Table 1. Seawater parameters during the OA experiment. A_{T} : total alkalinity; C_{T} : total dissolved inorganic carbon. Analysis of nutrition gave estimates of nitrate ($12.3 \mu\text{mol kg}^{-1}$), phosphate ($1.3 \mu\text{mol kg}^{-1}$) and silicate ($7.1 \mu\text{mol kg}^{-1}$) used in the calculations.

	Ambient	Medium $p\text{CO}_2$ treatment	High $p\text{CO}_2$ treatment
$p\text{CO}_2$ (μatm)	692 ± 26	727 ± 12	1198 ± 157
Salinity (ppt)	33.37 ± 0.12	33.37 ± 0.12	33.34 ± 0.09
A_{T} ($\mu\text{mol kg}^{-1}$)	2314.4 ± 2.5	2310.3 ± 1.1	2318 ± 14.3
C_{T} ($\mu\text{mol kg}^{-1}$)	2168.3 ± 6.6	2171.3 ± 1.5	2266.4 ± 31.8
HCO_3^-	2028.5 ± 9.2	2034.6 ± 2.6	2154.7 ± 35.4
CO_3^{2-}	116.2 ± 3.7	112.0 ± 1.5	66.1 ± 11.0
pH	7.84 ± 0.01	7.82 ± 0.01	7.62 ± 0.05
Ω_{calcite}	2.81 ± 0.09	2.71 ± 0.04	1.60 ± 0.27
$\Omega_{\text{aragonite}}$	1.81 ± 0.06	1.75 ± 0.02	1.02 ± 0.18

expected (Fig. 3) and only slightly lower than the medium $p\text{CO}_2$ treatment. In the experiments monitoring the surviving juveniles from the exposed treatments grown in ambient water until 8 months of age, pH varied around 7.95 to 7.96 (data not shown).

2.2 Brood stock

Ovigerous females were collected from the *H. gammarus* population in Øy garden ($60^\circ 35' \text{N}$; $4^\circ 50' \text{E}$) during the commercial fishing season October to November 2011, and transported to IMR-Matre. They were acclimatized at 6 °C in individual 75 L tanks ($52 \times 52 \times 28$ cm) in a CT room with the lighting set to a 12 h dark:12 h light cycle. Lobsters were fed frozen shrimps and fish twice a week. The CO_2 -control system was operative in late September 2011, and the hatching was postponed accordingly. In early September the temperature was slowly increased to 18 °C, to induce hatching that commenced 28 September 2011. Of a total of 14 females, only four had eggs that hatched during the experimental period. The sizes of the ovigerous females were 91, 110, 113 and 135 mm carapace length (CL; measured as the distance from the posterior rim of the eye socket to the posterior edge of the carapace). The females hatched 15 160 larvae in total.

2.3 Larval rearing and sampling

Each of the individual tanks with the ovigerous females had an overflow through a 20 mm water hose leading to separate containers equipped with a filter to retain the hatched larvae. Larvae normally hatch during late night/dawn and were collected from the outflow containers each morning, counted and transferred in equal numbers to the 40 L upstream incubators (plankton Kreisler; Hughes et al., 1974). However, all females were not hatching at the same time, thus we chose to run the experiments with $p\text{CO}_2$ treatments at 10 °C before commencing at 18 °C. The incubators were supplied with




Fig. 1. Overview of the experimental setup during the (a) larval (two incubators as parallels in each tank unit) and in the (b) juvenile phase with trays with individual compartments (several trays in each tank unit).

11 L seawater per minute. The larvae were fed daily with frozen *Artemia* sp. Larvae hatching over a period of 3 days were mixed in the same incubator. Larvae with larger difference in age were not mixed due to increased risk of cannibalism. Maximum density for each incubator was set to 1000 larvae, or 25 larvae per litre. Each treatment, i.e. temperature and CO_2 had two replicates (as shown in Fig. 1). Every third day the incubators were treated with Chloramid-T to control growth of the bacterium *Leucothrix mucor*, as previously experienced in other lobster-rearing systems (A.-L. Agnalt, personal communication, 2011.). The larvae were staged 1 to 4, according to Sars (1875) and Herrick (1909). Care was taken to look for intermediate larval stages, as this has been observed in American lobster (*H. americanus*), especially between stage 2 and 3 and between stage 3 and 4 (Templeman, 1936; Wells and Sprague, 1976; Charmantier and Aiken, 1987). Intermediate larval stages have also been observed in hybrids, i.e. offspring from female American lobster and male European lobster (Agnalt, unpublished data).

A total of 10 larvae at each development stage (1 to 4) from each incubator were sampled for measurement of CL and dry weight. At each stage, larvae had been exposed for a mini-

imum of three days before sampling. All measurements of CL were recorded using a dissecting microscope. Dry weights of individual larvae were recorded after 3 days of drying in Termaks dry oven at 60 °C, and recorded to the closest microgram (mg) using a Mettler Toledo scale (AG204 Delta Range). As many of the samples as possible were processed while the larvae were still alive. When the time schedule was too tight the larvae were frozen individually to be processed at a later stage. To utilize the system fully, whenever an incubator was terminated, and provided there were still newly hatched larvae available, another production line was started. Two batches of larvae production was made, with consecutive sampling, in incubator 2a, 2b, 3a, 14b, 15a and 15b (Fig. 1). Unfortunately, during storage the freezer containing many of the frozen samples of the larval stages broke. For these samples, CL and dry weight were recorded for as many larvae as possible. For the experiments run at 10 °C, 185 out of 216 sampled larvae have size recordings, and correspondingly at 18 °C, 224 out of 323 sampled larvae.




Fig. 2. Variation in pH in the larval phase run in ambient, medium $p\text{CO}_2$ and high $p\text{CO}_2$ at (a) 10°C and (b) 18°C in 2011. Ambient is raw water from 90 m depth. Note that the 10°C experiment started before 18°C.

2.4 Long-term exposure of juveniles at five months

While still pelagic, stage 4 larvae were collected one by one and transferred to trays consisting of single-cell compartments, made of black PVC plastic. Each tray consisted of 30 to 40 individual compartments with perforated bottoms (1 mm × 1 mm holes) to ensure water flow. Three to four trays were placed together in 400 L units (87 × 87 × 53 cm). Each unit was given water quality according to ambient, medium $p\text{CO}_2$ or high $p\text{CO}_2$ treatment. Water flow was set to 18 L per minute. The lobster juveniles were fed commercially produced pellets (2 mm), patented by Norwegian Lobster Farm (<http://www.norwegian-lobster-farm.com/no>) and produced by Nofima (<http://www.nofima.no/>). Whenever a juvenile moulted, the old exoskeleton was not removed from the compartment. On 22 March, CL were recorded for each of the surviving juveniles.

2.5 Continued monitoring of surviving juveniles in ambient water until 1 year old

From five months the juveniles were all kept in ambient water, i.e. the water quality in the raw water intake, at 14°C from 22 March to 25 October 2012. The purpose was to verify if the deformities observed were retained or lost through moulting when kept in water of higher pH (between 7.95 and 7.96). The lobster juveniles were fed commercially produced pellets (5 mm), patented by Norwegian Lobster Farm (<http://www.norwegian-lobster-farm.com/no>) and produced by Nofima (<http://www.nofima.no/>). At the end of the experiment, CL was recorded for each surviving juvenile.

2.6 Statistics

To determine whether parallels could be pooled for each $p\text{CO}_2$ treatment and each temperature we used Mann–Whitney U test. A Kruskal–Wallis test was used to determine if there were significant differences between lobsters undergoing the different $p\text{CO}_2$ and temperature treatments.




Fig. 3. Variation in pH in the juvenile phase in ambient, medium $p\text{CO}_2$ and high $p\text{CO}_2$ until 22 March 2012 in the experiments run at 18°C. Ambient is raw water from 90 m depth. Note that only juveniles in 18°C were monitored, as non larvae reached a viable stage 4 in 10°C.

One-way ANOVA was used to analyse length frequency of the juveniles raised in the different $p\text{CO}_2$ treatments.

3 Results

3.1 Growth

Growth was very slow at 10°C, irrespective of $p\text{CO}_2$ level, and after 5 weeks none of the larvae had moulted into stage 4. No larvae reached stage 3 in the medium $p\text{CO}_2$ treatment. Eventually three larvae reached stage 4, in high $p\text{CO}_2$ treatment only, but died within the following two days. At 18°C, development from stage 1 to 4 lasted from 14 to 16 days independent of $p\text{CO}_2$ treatment as predicted under optimal conditions. Of the 409 larvae investigated, none were found in intermediate stages.

At 10°C, carapace length at stage 1 and stage 3 did not differ significantly between $p\text{CO}_2$ treatments ($p > 0.5$, Fig. 4.). However, there were significant differences at stage 2 ($p < 0.05$), and the larvae produced in medium $p\text{CO}_2$ treatment were slightly smaller in CL compared with ambient and high $p\text{CO}_2$ treatment. At 18°C, CL was significantly different at stage 2 and stage 3 ($p < 0.05$, Fig. 4.). At stage 2, larvae reared in medium $p\text{CO}_2$ treatment were on average larger than the larvae reared in the other treatments, but at stage 3 they were smaller. Dry weight differed significantly between $p\text{CO}_2$ treatments at stage 1 and 2 at 10°C, and stage 2 and 3 at 18°C ($p < 0.05$, Fig. 4). Larvae raised in ambient water at 10°C were lighter in weight at stage 1 and heavier at stage 2 compared with the treated larvae. At 18°C, the stage 2 larvae raised in ambient water were slightly heavier than the other treatments but at stage 3 the high $p\text{CO}_2$ treatment larvae were heavier. In other words, there was no consistence and clear effect of $p\text{CO}_2$ treatment on either CL or dry weight.




Fig. 4. Mean carapace length (mm) and mean dry weight (mg) with one standard deviation at each larval stage of *Homarus gammarus* undergoing ambient, medium $p\text{CO}_2$ and high $p\text{CO}_2$ treatment run at 10 and 18 °C. Number of observations in each stage is given above the column. Note that at 10 °C only three larvae reached stage 4 (high $p\text{CO}_2$ treatment), but they only survived for two days. Ambient is raw water from 90 m depth. Star indicates significant differences based on Kruskal–Wallis non-parametric test at $p = 0.05$ level.

As only larvae raised at 18 °C successfully moulted into stage 4, this was the only temperature for which long-term exposure to $p\text{CO}_2$ could be monitored. Survival from stage 4 until 5 months age averaged 46 % in ambient, 17 % in medium and 61 % in high $p\text{CO}_2$ treatment. In total 148 juveniles survived. There were significant differences in CL as a result of $p\text{CO}_2$ treatment ($p < 0.05$, Fig. 5), but the sample size was however low at the medium $p\text{CO}_2$ treatment ($N = 16$), accounting for the difference since there were no significant differences between ambient and high $p\text{CO}_2$ treatment ($p > 0.1$, Fig. 5).

3.2 Deformities

Deformities, i.e. a difference in the shape of a body part or organ compared to the normal shape was found in the larvae (Table 2) and the juveniles (Table 3). The morphological abnormalities in the larvae were classified as curled carapace (Fig. 6), damages to the tail fan or that the rostrum was bent (Table 2). No larvae suffered multiple deformities. The most affected part was curled carapace, occurring in 59 % of the deformed larvae when combining all treatments. Bent rostrum was found in 27 % and damages to the tail fan in 14 % of all the deformed larvae. The deformities in the larvae were

Table 2. Classification of deformities found in larval stages 1 to 4 in European lobster (*Homarus gammarus*) exposed to elevated $p\text{CO}_2$ levels.

Category	Organ affected	Description
Curled carapace	Carapace	The carapace was curled at the edge, often forming ridges penetrating the side of the carapace.
Tail-fan damages	Uropod	Damages to parts of the tail fan, or even lacking one or both of the tail fans.
Bent rostrum	Rostrum	The rostrum was bent, as if not yet straighten out after hatching.

observed and described in live samples only. It was difficult to distinguish misshapes as described in Table 2 from artefacts due to the freezing process. Hence frozen samples were not included in the calculations below.

The proportion of larvae with deformities increased with increasing $p\text{CO}_2$ exposure, but was similar across the two temperatures 10 and 18 °C. In the medium treatment, 22 and 24 % were deformed at 10 and 18 °C, respectively. At the high treatment as much as 42 and 45 % were deformed, respectively. An overall 12 % of the larvae in ambient water raised in 18 °C developed deformities, though it is important to note that pH in ambient treatment was only slightly higher than in the medium $p\text{CO}_2$ treatment. At 10 °C, 5 % of the larvae ($N = 2$) were deformed, but these two specimens were actually sampled after 11 November when pH dropped to below 7.9. None of the larvae exposed to pH above 7.9 developed deformities. Curled carapace was the most common deformity (45 %) found in larvae raised in medium $p\text{CO}_2$ treatment at 10 °C, followed by damages in the tail fan (33 %) and bent rostrum (22 %). Concurrently, in the high $p\text{CO}_2$ treatment curled carapace was the only deformity. In the deformed larvae raised at 18 °C and in ambient treatment, curled carapace (33.3 %), damages to the tail fan (33.3 %) and bent rostrum (33.3 %) were equally distributed. Concurrently, in the medium treatment half of the deformed larvae had a bent rostrum, 38 % a curled carapace and 12 % were found with damages in the tail fan. Curled carapace was the only deformity found in the larvae undergoing high $p\text{CO}_2$ treatment.

Of the 148 juveniles that survived after five months, 41 were classified as morphologically deformed (see Table 2). Overall, 33 and 44 % of the juveniles in ambient and medium $p\text{CO}_2$ treatments, respectively, were deformed compared with 21 % in juveniles exposed to high $p\text{CO}_2$ (Fig. 8). In ambient and medium $p\text{CO}_2$ treatment, deformed claws were most often found (56 %), followed by stiff/twisted walking legs (39 %) and puffy carapace (39 %). When the juveniles

Table 3. Classification of deformities found in juvenile stages of European lobster (*Homarus gammarus*) exposed to elevated $p\text{CO}_2$ levels.

Category	Organ affected	Description
Puffy carapace	Carapace	Carapace puffy/swollen, or up-folded on one side, often leaving some parts of the gills exposed.
Stiff/twisted walking legs	2–5 pereopod	The joints were fused together as if the joints were over-calcified. The entire pereopod leg was like one stiff piece, sometimes “frozen” in an arbitrary/twisted position.
Misshaped claw	1st pereopod/cheliped	Various shapes of the cheliped, but most often twisted, deviating from normal.
Bent rostrum	Rostrum	The rostrum was bent, as if not yet erected after hatching.
Tail-fan damages	Uropod	Damages to parts of the tail fan, or even lacking one or both of the tail fans.
Twisted abdomen	Abdomen	Abnormal shape of the abdomen as if some of the segments were once broken and then grown back in the wrong shape.
Stiff antenna	2 antenna	Segments of the antenna were fused, as if the joints were over-calcified. Felt “stiff” when touching. Difficult to observe when animal was out of water.

from ambient and medium $p\text{CO}_2$ treatment had deformed claw(s) about 54 % had also developed stiff/twisted walking legs. In comparison, at high $p\text{CO}_2$, 71 % of the deformed juveniles had developed a puffy carapace. Of these, 24 % had also developed deformed claws. Overall, about half of the deformed juveniles from the ambient and medium $p\text{CO}_2$ treatment had developed two or three different abnormalities (Fig. 10). In comparison, 70 % of the deformed juveniles in the high treatment had multiple deformities.

At one year of age, 76 of the 148 juveniles had survived. Mortality was equally high for those juveniles that had 5 months’ exposure in ambient or in high $p\text{CO}_2$. Overall, 28 % of the juveniles were deformed. The most common occurring deformities (28 %) were puffy carapace with stiff/twisted walking leg, as illustrated in Fig. 10. Of the 40 deformed juveniles found 22 March 2012, only 12 had survived another seven months. Of these, six were still deformed (four of the juveniles even developed at least one additional deformity). In other words, 50 % of the survivors had managed to recuperate, most likely through moulting. However, of the 108 juveniles classified as normal at five months of age, 15 had developed deformities seven months later.

4 Discussion

Deformities were observed both in the larval and juvenile phase in European lobster when exposed to higher $p\text{CO}_2$ from hatching. The proportion of larvae with deformities increased with increasing $p\text{CO}_2$ exposure and was similar across the two temperatures 10 and 18 °C. In high exposure as much as 45 % of the larvae developed deformities. After five months of exposure, 44 and 21 % of the juveniles were deformed in medium and high $p\text{CO}_2$ treatment, respectively. Deformities in lobster larvae and juveniles have not previously been reported, either in European or American lobster, although the scientific community have long-term experience in husbandry of these two species (Gruffydd et al., 1975; Capuzzo and Lancaster, 1979; Latrouite and Lorec, 1991; Addison and Bannister, 1994; Uglem et al., 1995; Agnalt et al., 1999, 2004; Nicosia and Lavalli, 1999; Linnane et al., 2000; Jørstad et al., 2001; Wickins and Lee, 2002; Kristiansen et al., 2004; Jørstad et al., 2005; Agnalt, 2008; Arnold et al., 2009; Ries et al., 2009; Schmalenbach et al., 2009; Keppel et al., 2012). Wickins et al. (1995) did report moulting abnormalities in European lobster larvae in relation to testing different diets, but with no further description




Fig. 5. Carapace length (mm) frequency of juvenile *Homarus gammarus* raised in 18 °C, from newly hatched larvae until 5 months of age in (a) ambient water, (b) medium $p\text{CO}_2$ treatment and (c) high $p\text{CO}_2$ treatment. Size recordings were made 22 March 2012.

of what the abnormalities were. In aquaculture, hatchery-induced changes due to feed, tank design and/or substrate have been described for a number of fish and shellfish species (Olla et al., 1998; Svåsand et al., 1998; Tsukamoto et al., 1999; and references therein). In shellfish, most changes documented are morphological, e.g. lower shell strength in the great scallop (*Pecten maximus*) (Grefsrud and Strand, 2006) and queen conch (*Strombus gigas*) (Stoner and Davis, 1994), lack of spines in top shell (*Trochus niloticus*) (Purcell 2002) or lack of differentiation in the claws in lobster *Homarus* spp. (Govind and Pearce, 1986). Deformities of 40 to 58 % have also been found in wild populations of shrimps of the genus *Palaemon* in the Gironde estuary in France (Béguer et al., 2008, 2010). The deformities (wrinkled or bent carapace, bent rostrum and damages in the tail fan) were reported to affect adult individual mortality and egg production. The cause of these deformities was not identified, although stress, pollution or even elevated $p\text{CO}_2$ levels may be explanatory factors. In future studies it will be vital to differentiate de-



Fig. 6. A stage 3 *Homarus gammarus* larvae raised in an environment with pH lower than 7.9. The carapace is curled, leaving "curls" on the side (indicated with the arrow). This is classified as a deformity.

formities caused by ocean acidification from effects due to hatchery production or other environmental factors.

Why are we so sure that the deformities observed in this study were due to high exposure $p\text{CO}_2$? The Institute of Marine Research has since the early 1990s hatched and produced larvae and juvenile European lobster for various studies such as, e.g. stock enhancement (Agnalt et al., 1999, 2004; Agnalt 2008), fitness and genetic studies (Jørstad et al., 2001; Jørstad et al., 2005), carrying capacity (Agnalt, unpublished data) and conditioning juveniles for release purposes (Agnalt, 2013; Aspaas, 2012; Trengereid, 2012). We have only in a few occasions found that the first pair of peripods (claws) was misshaped but this trait was lost after one or two moults. None of these misshapes looked like the claw deformities found in the present study. Assessing deformities demands experience with observing larvae and juveniles, as subtle trait changes might be missed. In the present study, deformities in the larval phase were found in the carapace (termed "curled" in larvae and "puffy" in juveniles), tail fan or in the rostrum (termed "bent rostrum") (Tables 2 and 3). Inexperienced personnel might miss these traits. In this study, measurements were made by one person to ensure consistency and the individuals were classified blindly. Experience should be accounted for also in future studies, whenever possible.

Our study shows that deformities occur in lobster larvae raised in $p\text{CO}_2$ levels higher than 727 μatm , independently of temperatures being 10 or 18 °C. Deformities due to OA has been documented on the embryonic in some marine invertebrate species, often resulting in low or even no hatching success (Parker et al., 2009; Kawaguchi et al., 2010; Byrne, 2011), but also in the larval stages in a few species (Kurihara, 2008; Byrne et al., 2011, and references therein). Whiteley (2011) stated that the exoskeleton of planktonic decapod larvae is unmineralised and elevated $p\text{CO}_2$ should therefore




Fig. 7. Percentage of *Homarus gammarus* with deformities in the larval phase in the different temperature, 10 and 18 °C, and pCO₂ treatments (ambient, medium and high).

not affect larval conditions. However, Arnold et al. (2009) analysed the calcium and magnesium concentrations per surface area of the carapace of *H. gammarus* larvae. They found significant reductions at elevated pCO₂ of 1200 μatm in stage 4 larvae; calcium decreased from 0.24 to 0.13 μg mm⁻² and magnesium decreased from 0.019 to 0.012 μg mm⁻². Calcium and magnesium seem to be important mineral components in the exoskeleton of the lobster larvae. However, Mg-CaCO₃ is more soluble than pure calcite or aragonite (Andersson et al., 2008) and since Ω_{Aragonite} was at the lowest 1.02 in this study, elevated pCO₂ could definitely have an effect on e.g. formation of the new exoskeleton. A thinning of the shell has been reported in blue mussels *Mytilus edulis* and Pacific Oyster *Crassostrea gigas* (Gazeau et al., 2007; Melzner et al., 2011). A thinner shell in lobster larvae may not be strong enough to keep its shape, especially when covering organs like the gills where water flow continuously and applies pressure on the shell. This may explain the “puffy” carapace observed in the present study.

In the high pCO₂ treatment carapace was the most affected organ, both in larvae and in the juveniles. In the juveniles, the deformities generally affected carapace (puffy or swollen, thus often leaving part of the gills exposed), walking legs (stiff), claws (twisted), abdomen (stiff joints), tail fan and even antenna (stiff). The latter is of vital importance for lobster communication, including finding a partner (Johnson and Atema, 2005). Fewer deformities were found in juveniles exposed to high pCO₂ compared with medium exposure. However, those affected had also developed multiple deformities, i.e. more severe damages. We found that damages to the tail fan could not be repaired through moulting, while walking legs and “puffy” carapace would become normal after several moultings. The exoskeleton in sub-adult and adult lobster can be divided into three layers: epi-, exo- and endocuticle (e.g. Sachs et al., 2006; Bosselmann et al., 2007; Romano et al., 2007; Al-Sawalmih et al., 2008; Sachs et al., 2008; Fabritius et al., 2009). The exoskeleton of *Homarus* sp. consists of chitin, protein, calcium, magnesium, phos-




Fig. 8. The percentage of juveniles with deformities. Light grey bar represents juveniles surviving five months of pCO₂ exposure (ambient, medium and high) and darker grey bar the juveniles after another seven months in ambient water only. Total number of juveniles in each category is given above each column.

phate and a few other compounds (Ba, Mn, Sr), although variable in composition in the different layers (Al-Sawalmih et al., 2008; Kunkel et al., 2012). The epicuticle is particularly rich in calcium, magnesium and phosphate. The outermost exocuticle consists of a thin calcite-containing layer while the rest is fully mineralized, mostly with amorphous calcium carbonate that is highly soluble and acts as a transient source of calcium. The endocuticle consists of crystalline calcite. Moulting is a complex process (Greenaway, 1985; Dillaman et al., 2005; Politi et al., 2010), and for instance, reabsorption of calcium occurs from the old exoskeleton before it is shed. The new exoskeleton is uncalcified and rapidly needs to harden by deposition of calcium carbonate. In other words, elevated pCO₂ can affect various cuticle layers and various stages of the moulting cycle. Thus, the deformities may have been caused by irregularities in the depositing of Mg-CaCO₃ in the shell. Too high deposits, a possible compensation for low hemolymph pH (acidosis), may have resulted in the stiff walking legs observed in our study. Whereas too little depositing of CaCO₃ in the shell may result in the incomplete formation of certain structures in the exoskeleton. This could be related to depleted energy resources (to maintain homeostasis) required for converting HCO₃⁻ to CO₃²⁺ and therefore the depositing of enough CaCO₃.

No consistent and clear effect of pCO₂ treatment was found in this study on neither carapace length, or on dry weight. This is coherent with observations with the same species by Arnold et al. (2009). Neither was moulting frequency affected by increased acidification in this study. However, Keppel et al. (2012) found decreased carapace length with increased pCO₂ in the closely related *H. americanus*, and the moulting cycle was prolonged with about 2 days in




Fig. 9. The percentage of lobster juveniles with single or multiple deformities when exposed to elevated levels of $p\text{CO}_2$ (ambient, medium and high) for five months.

larvae raised at $\text{pH} = 7.7$ compared with $\text{pH} = 8.1$. It seems that the growth and moulting cycle is affected in *H. americanus* (Keppel et al., 2012), while larvae and juveniles of *H. gammarus* maintain growth rates at the cost of mineralization of the exoskeleton.

A major concern with the present study was the very low pH with corresponding high $p\text{CO}_2$ (average $692 \pm 26 \mu\text{atm}$) found in ambient water taken from a locality in Masfjorden at 90 m depth. Compared to pH in ambient water at IMR's research stations Austevoll ($\text{pH} 7.98$, Andersen et al., 2013) and Parisvatn ($\text{pH} 8.11$; Agnalt, personal communication) located in outer coastal areas in the same region, the water in Masfjorden seems to have quite a large natural variation and may drop to levels well below global average pH of 8.05 (Fabry et al., 2008). European lobster is not normally found at the depths of 90 m at this locality in Masfjorden, but is found in the fjord system at shallower depths. Knowledge about the pH in the Norwegian fjord systems and how it fluctuates between seasons and years is scarce. However, measurements from other locations in the Northern Hemisphere shows that pH in coastal areas can be highly variable and as low as 7.7–7.6. Barton et al. (2012) showed that pH in Oregon coastal water varied from 7.6 to 8.2 in the early summer of 2009. Newton et al. (2012) reported a pH of 7.72 at 100 m depth in Puget Sound and the Strait of Juan de Fuca, on the East Coast of the USA during February 2008. Even in the “closer to home” North Sea, there are already indications that the level of pH is variable and decreasing (Olsen et al., 2006; Blackford and Gilbert, 2007; Bellerby et al., 2005). Based on these studies, we assume that fjord systems also are affected by the ongoing ocean acidification and may explain the low ambient conditions in Masfjorden. However, more effort is needed to verify this hypothesis.




Fig. 10. *Homarus gammarus* juvenile, one year of age, exposed to $\text{pH} < 8.1$ since hatch displaying deformities as puffy carapace (circle), stiff/twisted walking legs (arrow) and lacking antennae (square).

5 Conclusions and future work

There were no clear effects of $p\text{CO}_2$ treatment, on either carapace length or dry weight in *H. gammarus* larvae. However, the high ratio of larvae and juveniles with deformed exoskeletons strongly indicates a negative effect of elevated $p\text{CO}_2$ on European lobster from hatching to one year of age. Some of the deformities may affect the ability to find food and partners (walking legs, claw and antenna), respiration (carapace), and the ability to swim (tail-fan damages). Thus further studies on behaviour and respiration are needed. Studies have already commenced to elucidate mineralization of the exoskeleton in lobster juveniles exposed to various levels of $p\text{CO}_2$ and elevated temperature.

Acknowledgements. We thank the staff at the Institute of Marine Research, Matre Research Station, for helpful assistance during the project period. We also thank the two anonymous referees and Daniela Schmidt for their valuable comments. This study was supported by the Institute of Marine Research through the project 13193-01, Ocean Acidification-Lobster.

Edited by: D. Schmidt

References

- Addison, J. T. and Bannister, R. C. A.: Re-stocking and enhancement of clawed lobster stocks: a review, *Crustaceana*, 67, 131–155, 1994.
- Agnalt, A.-L.: Stock enhancement of European lobster (*Homarus gammarus*) in Norway; Comparisons of reproduction, growth and movement between wild and cultured lobster, Scientiarium, University of Bergen, Norway, p. 140, 2008.
- Agnalt, A.-L., van der Meeren, G.I., Jørstad, K.E., Næss, H., Farestveit, E., Nøstvold, E., Svåsand, T., Korsåen, E., and Ydstebø, L.: Stock enhancement of European lobster (*Homarus gammarus*); a large-scale experiment off southwestern Norway (Kvitsøy), 401–419, in: Stock enhancement and sea ranching, edited by: Howell, B., Moksness, E., and Svåsand, T., Fishing News Books, Blackwell Science Ltd, Oxford, p. 606, 1999.
- Agnalt, A.-L., Jørstad, K.E., Kristiansen, T.S., Nøstvold, E., Farestveit, E., Næss, H., Paulsen, O.I., and Svåsand, T.: Enhancing the European lobster (*Homarus gammarus*) stock at Kvitsøy Islands; Perspectives of rebuilding Norwegian stocks, 415–426, in: Stock enhancement and sea ranching developments, edited by: Leber, K. M., Kitada, S., Blankenship, H. L., and Svåsand, T., pitfalls and opportunities, Blackwell Publishing Ltd, Oxford, p. 562, 2004.
- Agnalt, A.-L., Farestveit, E., Gundersen, K., Jørstad, K. E., and Kristiansen, T. S.: Population characteristics of the world's northernmost stock of European lobster (*Homarus gammarus*) in Tysfjord and Nordfolda, northern Norway, *New Zeal. J. Mar. Fresh.*, 43, 47–57, 2009.
- Al-Sawalmih, A., Li, C., Siegel, S., Fabritius, H., Yi, S., Raabe, D., Frazl, P., and Paris, O.: Microtexture and chitin/calcite orientation relationship in the mineralized exoskeleton of the American lobster, *Adv. Funct. Mater.*, 18, 3307–3314, 2008.
- Andersen, S., Grefsrud, E. S., and Harboe, T.: Effect of increased $p\text{CO}_2$ on early shell development in great scallop (*Pecten maximus* Lamarck) larvae. Submitted to *Biogeosciences*, special issue: The ocean in a high CO world III, 2013.
- Andersson, A. J., Mackenzie, F. T., and Bates, N. R.: Life on the margin: implications of ocean acidification on Mg-calcite, high latitude and cold-water marine calcifiers, *Mar. Ecol. Prog. Ser.*, 373, 265–273, 2008.
- Arnold, K. E., Findlay, H. S., Spicer, J. I., Daniels, C. L., and Boothroyd, D.: Effect of CO_2 -related acidification on aspects of the larval development of the European lobster, *Homarus gammarus* (L.), *Biogeosciences*, 6, 1747–1754, doi:10.5194/bg-6-1747-2009, 2009.
- Aspaas, S.: Behavior of hatchery-produced European lobster (*Homarus gammarus*), comparing conditioned and naïve juveniles, Master of Science in Aquaculture Biology, University of Bergen, Norway, 1–49, 2012.
- Barton, A., Hales, B., Waldbusser, G. G., Langdon, C., and Feely, R. A.: The Pacific oyster, *Crassostrea gigas*, shows negative correlation to naturally elevated carbon dioxide levels: Implications for near-term ocean acidification effects, *Limnol. Oceanogr.*, 57, 698–710, 2012.
- Béguier, M., Pasquaud, S., Boët, P., and Girardin, M.: First description of heavy skeletal deformations in Palaemon shrimp populations of European estuaries: the case of the Gironde (France), *Hydrobiologia*, 607, 225–229, 2008.
- Béguier, M., Feuillassier, L., Elie, P., Boët, P., and Girardin, M.: Exoskeletal deformities in Palaemonidae: Are they a threat to survival?, *Mar. Environ. Res.*, 69, 109–117, 2010.
- Bellerby, R. G. J., Olsen, A., Furevik, T., and Anderson, L.G.: Response of the surface ocean CO_2 system in the Nordic Seas and northern North Atlantic to climate change, *Geophys. Monogr. Ser.*, 158, 189–197, 2005.
- Blackford, J. C. and Gilbert, F. J.: pH variability and CO_2 induced acidification in the North Sea, *J. Mar. Syst.*, 64, 229–241, 2007.
- Blunden, J. and D. S. Arndt, Eds.: State of the Climate in 2012, *Bull. Amer. Meteor. Soc.*, 94, S1–S238, 2013.
- Bosselmann, F., Romano, P., Fabritius, H., Raabe, D., and Epple, M.: The composition of the exoskeleton of two crustacean: The American lobster *Homarus americanus* and the edible crab *Cancer pagurus*, *Thermochimica Acta*, 463, 65–68, 2007.
- Byrne, M.: Impact of ocean warming and ocean acidification on marine invertebrates life history stages: vulnerability and potential for persistence in a changing ocean, *Oceanography and Mar. Biol. Ann. Review*, 49, 1–42, 2011.
- Byrne, M., Ho, M. A., Wong, E., Soars, N., Selvakumaraswamy, P., Sheppard Brennan, H., Dworjanyn, S. A., and Davis, A. R.: Unshelled abalone and corrupted urchins, development of marine calcifiers in a changing ocean, *Proc. R. Soc. Ser. B*, 278, 2376–2383, 2011.
- Caldeira, K. and Wickett M. E.: Ocean model predictions of chemistry changes from carbon dioxide emissions to the atmosphere and ocean, *J. Geophys. Res.*, 110, C09S04, doi:10.1029/2004JC002671, 2005.
- Capuzzo, J. M. and Lancaster, B. A.: Some physiological and biochemical considerations of larval development of larval development in the American lobster, *Homarus americanus* Milne Edwards, *J. Exp. Mar. Biol. Ecol.*, 40, 53–62, 1979.
- Charmantier, G. and Aiken, D. E.: Intermediate larval and postlarval stages of *Homarus americanus* H. Milne Edwards, 1837 (Crustacea: Decapoda), *J. Crust. Biol.*, 7, 525–535, 1987.
- Dickson, A. G., Sabine, C. L. and Christian, J. R. (Eds.): Guide to best practices for ocean CO_2 measurements, PICES Special Publication, 3, 1–191, 2007.
- Dillaman, R., Hequembourg, S., and Gay, M.: Early pattern of calcification in the dorsal carapace of the blue crab, *Callinectes sapidus*, *J. Morphol.*, 263, 356–374, 2005.
- Fabritius, H.-O., Sachs, C., Romano, P. T., and Raabe, D.: Influence of structural principles on the mechanics of a biological fiber-based composite material with hierarchical organization: The exoskeleton of the lobster *Homarus americanus*, *Adv. Mater.*, 21, 391–400, 2009.
- Fabry, V. J., Seibel, B. A., Feely, R. A., and Orr, J. C.: Impacts of ocean acidification on marine fauna and ecosystem processes, *ICES J. Mar. Sci.*, 65, 414–432, 2008.
- Factor, J. R. (Ed): Biology of the Lobster *Homarus americanus*, Toronto, Academic press, 1995.
- Feely, R. A., Doney, S. C., and Cooley, S. R.: Ocean Acidification: Present Conditions and Future Changes in a High- CO_2 World, *Oceanography*, 22, 36–47, 2009.
- Gazeau, F., Quiblier, C., Jansen, J. M., Gattuso, J.-P., Middelburg, J. J., and Heip, C. H. R.: Impact of elevated CO_2 on shellfish calcification, *Geophysical Res. Lett.*, 34, 1–5, 2007.

- Govind, C. K., and Pearce, J.: Differential reflex activity determines claw and closer muscle asymmetry in developing lobsters, *Science*, 233, 254–256, 1986.
- Greenaway, P.: Calcium balance and moulting in the crustacea, *Biol. Rev.*, 60, 425–454, 1985.
- Grefsrud, E. S., and Strand, Ø.: Comparison of shell strength in wild and cultured scallops (*Pecten maximus*), *Aquaculture*, 251, 306–313, 2006.
- Gruffydd, L. L. D., Rieser, R. A., and Machin, D.: A comparison of growth and temperature tolerance in the larvae of the lobster *Homarus gammarus* (L.) and *Homarus americanus* H. Milne Edwards (Decapoda, Nephropidae), *Crustaceana*, 28, 23–32, 1975.
- IPCC: Climate change 2007: The physical science basis. In Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, UK, 2007.
- Herrick, F. H.: Natural history of the American lobster, *Bull. US Bur. Fish.*, 29, 149–408, 1909.
- Hughes, J. T., Shleser, R. A., and Tchobanoglous, G.: A rearing tank for lobster larvae and other aquatic species, *Progve. Fish. Cult.*, 36, 129–132, 1974.
- Johnsen, M. E. and Atema, J.: The olfactory pathway for individual recognition in the American lobster *Homarus americanus*, *J. Exp. Biol.*, 208, 2865–2872, 2005.
- Jørstad, K. E., Nøstvold, E., Kristiansen, T. S., and Agnalt, A.-L.: High survival and growth of European lobster juveniles (*Homarus gammarus*), reared communally with natural bottom substrate, *Mar. Freshw. Res.*, 52, 1431–1438, 2001.
- Jørstad, K. E., Prodöhl, P., Kristiansen, T. S., Hughes, M., Farestveit, E., Taggart, J. B., Agnalt, A.-L., and Ferguson, A.: Communal larval rearing European lobster (*Homarus gammarus*): Family identification by microsatellite DNA profiling and offspring fitness comparisons, *Aquaculture*, 247, 257–285, 2005.
- Kawaguchi, S., Kurhara, H., King, R., Hale, L., Berli, T., Robinson, J.P., Ishida, A., Wakita, M., Virtue, P., Nicol, S., and Ishimatsu, A.: Will krill fare well under southern ocean acidification?, *Biol. Lett.*, 5 November, 2010.
- Keppel, E. A., Scrosati, R. A., and Courtenay, A. C.: Ocean acidification decreases growth and development in American lobster (*Homarus americanus*) larvae, *J. Northw. Atl. Fish. Sci.*, 44, 61–66, 2012.
- Kristiansen, T. S., Drengstig, A., Bergheim, A., Drengstig, T., Kollsgård, I., Svendsen, R., Nøstvold, E., Farestveit, E., and Aardal, L.: Development of methods for intensive farming of European lobster in recirculated seawater. Results from experiments conducted at Kvitsøy lobster hatchery from 2000 to 2004, *Fisken og havet*, 6, 1–52, 2004.
- Kunkel, J. G., Nagel, W., and Jercinovic, M. J.: Mineral fine structure of the American lobster cuticle, *J. Shellfish. Res.*, 31, 515–526, 2012.
- Kurihara, H.: Effects of CO₂-driven ocean acidification on the early developmental stages of invertebrates, *Mar. Ecol. Prog. Ser.*, 373, 275–284, doi:10.3354/meps07802, 2008.
- Latrouite, D. and Lorec, J.: L'expérience française de forçage du recrutement du homard européen (*Homarus gammarus*): resultants préliminaires, *ICES Mar. Sci. Symp.*, 192, 93–98, 1991.
- Linnane, A., Ball, B., Munday, B., and Mercer, J. P.: A long-term mesocosm study on the settlement and survival of juvenile European lobster *Homarus gammarus* L. in four natural substrata, *J. Exp. Mar. Biol. Ecol.*, 249, 51–64, 2000.
- Linnane, A., Brendan, B., Mercer, J. P., Browne, R., van der Meeren, G.I., Ringvold, H., Bannister, C., Mazzoni, D., and Munday, B.: Searching for the early benthic phase (EBP) of the European lobster: a trans-European study of cobble fauna, *Hydrobiologia*, 465, 63–72, 2001.
- Melzner, F., Stange, P., Trübenbach, K., Thomsen, J., Casties, I., Panknin, U., Gorb, S. N., and Gutowska, A.: Food supply and seawater pCO₂ impact calcification and internal shell dissolution in the blue mussel *Mytilus edulis*, *Plos one*, 6, 1–8, 2011.
- Newton, J. A., Feely, R. A., Alin, S. R., and Krembs, C.: Ocean acidification in Pudget Sound and the Strait of Juan de Fuca, 29–56, in: Scientific summary of ocean acidification in Washington State marine waters, edited by: Feely, T. A., Klinger, T., Newton, J. A., and Chadsey, M., Washington State Blue Ribbon Panel on Ocean Acidification, NOAA OAR Special Report, p. 159, 2012.
- Nicosia, F. and Lavalli, K.: Homarid lobster hatcheries: their history and role in research, management and aquaculture, *Mar. Fish. Rev.*, 61, 1–57, 1999.
- Olsen, A., Omar, A. M., Bellerby, R. G. J., Johannessen, T., Ninnemann, U., Brown, K. R., Olsson, K. A., Olafsson, J., Nondal, G., Kivimäe, C., Kringstad, S., Neill, C., and Olafsdottir, S.: Magnitude and origin of the anthropogenic CO₂ increase and 13C Suess effect in the Nordic seas since 1981, *Global Biogeochem. Cy.*, 20, GB3027, doi:10.1029/2005GB002669, 2006.
- Olla, B. L., Davis, M. W., and Ryer, C. H.: Behavioral deficits in hatchery-reared fish: potential effects on survival following release, *Aqua. Fish. Manag.*, 25, 19–34, 1994.
- Orr, J. C., Fabry, V. J., Aumont, O., Bopp, L., Doney, S. C., Feely, R. A., Gnanadesikan, A., Gruber, N., Ishida, A., Joos, F., Key, R. M., Lindsay, K., Maier-Reimer, E., Matear, R., Monfray, P., Mouchet, A., Najjar, R. G., Plattner, G. K., Rodgers, K. B., Sabine, C. L., Sarmiento, J. L., Schlitzer, R., Slater, R. D., Totterdell, I. J., Weirig, M. F., Yamanaka, Y., and Yool, A.: Anthropogenic ocean acidification over the twenty-first century and its impact on calcifying organisms, *Nature*, 437, 681–686, 2005.
- Parker, L. M., Rosa, P. M., and O'Conner, W. A.: The effect of ocean acidification and temperature on the fertilization and embryonic development of the Sydney rock oyster *Saccostrea glomerata* (Gould 1850), *Glob. Change Biol.*, 15, 2123–2136, 2009.
- Politik, Y., Batchelor, D. R. I., Zaslansky, P., Chmelka, B. F., Weaver, J. C., Sagi, I., Weiner, S., and Addadi, L.: Role of magnesium ion in the stabilization of biogenic amorphous calcium carbonate: A structure – function investigation, *Chem. Mater*, 22, 161–166, 2010.
- Pörtner, H. O., Langenbuch, M., and Reipschlag, A.: Biological impact of elevated ocean CO₂ concentrations: Lessons from animal physiology and earth history, *J. Oceanogr.*, 60, 705–718, 2004.
- Purcell, S. W.: Cultured vs wild juvenile *trochus*: Disparate shell morphologies send caution for seeding, *SPC Troch. Inf. Bull.*, 9, 6–8, 2002.
- Ries, J. B., Cohen, A. L., and McCorkle, D. C.: Marine calcifiers exhibit mixed responses to CO₂-induced ocean acidification, *Geology*, 37, 1131–1134, 2009.
- Romano, P., Fabritius, H., and Raabe, D.: The exoskeleton of the lobster *Homarus americanus* as an example of a smart

- anisotropic biological material, *Acta Biomaterialia*, 3, 301–309, 2007.
- Sachs, C., Fabritius, H., and Raabe, D.: Experimental investigation of the elastic-plastic deformation of mineralized lobster cuticle by digital image correlation, *J. Struct. Biol.*, 155, 409–425, 2006.
- Sachs, C., Fabritius, H., and Raabe, D.: Influence of microstructure on deformation anisotropy of mineralized cuticle from the lobster *Homarus americanus*, *J. Struct. Biol.*, 161, 120–132, 2008.
- Sars, G. O.: Om hummerens postembryonale udvikling (About the lobsters post-embryo development), *Christiania Videnskabs-Selskabs Forhandling*, Brøgger's Boktrykkeri, Christiania, 1–27, 1875.
- Schmalenbach, I. and F. Buchholz: Effects of temperature on the moulting and locomotory activity of hatchery-reared juvenile lobsters (*Homarus gammarus*) in Helgoland (North Sea), *Mar. Biol. Res.*, 9, 19–26, 2013.
- Schmalenbach, I., Buchholz, F., Franke, H.-D. F., and Saborowski, R.: Improvement of rearing conditions for juvenile lobsters (*Homarus gammarus*) by co-culturing with juvenile isopods (*Idotea emarginata*), *Aquaculture*, 289, 297–303, 2009.
- Skjelvan, I., Falck, E., Rey, F., and Kringstad, S. B.: Inorganic carbon time series at Ocean Weather Station M in the Norwegian Sea, *Biogeosciences*, 5, 549–560, doi:10.5194/bg-5-549-2008, 2008.
- Spicer, J. I., Raffo, A., and Widdicombe, S.: Influence of CO₂-related seawater acidification on extracellular acid-base balance in the velvet swimming crab *Necora puber*, *Mar. Biol.*, 151, 1117–1125, 2007.
- Stoner, A. W. and Davis, M.: Experimental outplanting of juvenile queen conch, *Strombus gigas*: comparison of wild and hatchery-reared stocks, *Fish. Bull.*, 92, 390–411, 1994.
- Svåsand, T., Skilbrei, O. T., van der Meeren, G. I., and Holm, M.: Review of morphological and behavioural differences between reared and wild individuals: implications for sea-ranching of Atlantic salmon, *Salmo salar* L., Atlantic cod, *Gadus morhua* L., and European lobster, *Homarus gammarus* L., *Fish. Manag. Ecol.*, 5, 473–490, 1998.
- Templeman, W.: Fourth stage larvae of *Homarus americanus* intermediate in form between normal third and fourth stages, *J. Biol. Board Can.*, 2, 349–354, 1936.
- Trengereid, H.: Shelter seeking and competitive ability behaviour in hatchery reared juvenile European lobster (*Homarus gammarus*) exposed to predator odors, Master of Science in aquaculture biology, University of Bergen, Norway, 1–83, 2012.
- Truchot, J. P.: Mechanisms of extracellular acid-base regulation as temperature changes in decapod crustaceans, *Respir. Physiol.*, 33, 161–176, 1978.
- Tsukamoto, K., Kuwada, H., Uchida, K., Masuda, R., and Sakaura, Y.: Fish quality and stocking effectiveness: behavioural approach, 205–218, in: Stock enhancement and sea ranching, edited by: Howell, B., Moksness, E., and Svåsand, T., Fishing News Books, Blackwell Science Ltd, Oxford, p. 606, 1999.
- Uglem, I., Grimsmen, S., Hold, M., Svåsand, T., and Korsøen, E.: Havbeite med hummer, yngelproduksjon (stock enhancement with lobster, juvenile production, final report PUSH), Sluttrapport PUSH, 1–30, 1995.
- Waddy, S. L., Aiken, D. E. and de Kleijn, D. P. V. (Eds): Control of Growth and reproduction. Chapter 10. In: *Biology of the lobster Homarus americanus* J. B. Factor, 1995.
- Walther, K., Anger, K., and Pörtner, H. O.: Effects of ocean acidification and warming on the larval development of the spider crab *Hyas araneus* from different latitudes (54 degrees vs. 79 degrees N), *Mar. Ecol-Prog. Ser.*, 417, 159–170, 2010.
- Wells, P. G. and Sprague, J. B.: Effects of crude oil on American lobster (*Homarus americanus*) larvae in the laboratory, *J. Fish. Res. Board Can.*, 33, 1604–1614, 1976.
- Whiteley, N. M.: Physiological and ecological responses of crustaceans to ocean acidification, *Mar. Ecol. Prog. Ser.*, 430, 257–271, 2011.
- Wickins, J. F. and Lee, O. C.: Crustacean farming, ranching and culture, Blackwell Science, London, p. 446, 2002.