

This paper not to be cited without prior reference to the Council

International Council for
the Exploration of the Sea

Statutory Meeting CM 1987/F: 2
Mariculture Committee

Ref.: Marine Environmental
Quality Committee

Report of the *ad hoc* Study Group on
"Environmental Impact of Mariculture"

by

¹Harald Rosenthal, ²Donald Weston,
³Richard Gowen and ⁴Edward Black
(Editorial Committee)

¹ Biologische Anstalt Helgoland,
Zentrale Hamburg, Notkestrasse 3,
2000 Hamburg 52,
Federal Republik of Germany

²School of Oceanography, WB-10
University of Washington,
Seattle, WA. 98195, USA

³ Department of Biological Science,
The University of Stirling,
Stirling FK9 4LA, Scotland

⁴Ministry of Agriculture and Fisheries
Marine Resource Section,
780 Blanshard Street, Victoria,
British Columbia, Canada, V8V 1X5

Ad hoc Study Group Membership:

Hans Ackefors	Zoologiska Institutionen, University of Stockholm, PO. Box 6801,113-86 Stockholm, Sweden
Jan Aure	Institute of Marine Research, P.O.Box 1870/72, Nordnes, 5024 Bergen, Norway
A.C. Drinkwaard	Netherlands Institute for Fishery Investigations, P.O.Box 68, 1970-AB Ijmuiden, The Netherlands
L.A. Dushkina	VNIRO, 7 Verkhne Krasnoselskaya, Moscow, B-140,USSR
Arne Ervik	Institute of Marine Research, P.O.Box 1870/72, Nordnes, 5024 Bergen, Norway
Maurice Héral	IFREMER, B.P.133, 17390 La Tremblade, France
Timo Mäkinen	Laukaa Fish Culture Research Station, 31360 Valkola, Finland
M.A. Sampayo	INIP, Avenida Brasilia, 1400 Lisbon, Portugal
David J. Wildish	Department of Fisheries and Oceans, Biological Station, St. Andrews, New Brunswick, Canada, E0G 2X0
Markku Pursiainen	Finnish Game and Fisheries Research Institute, POBax 193, Helsinki, Finland, 00131 (Observer EIFAC Working Party on Fish Farm Effluents)

Table of Contents

Abstract.....	3
Introduction.....	3
Environmental issues.....	6
- Bacteria.....	6
- Marine Algae.....	7
- Sedimentation and Benthos.....	9
- Chemicals in Aquaculture.....	11
- Water quality.....	15
- Site selection.....	18
- Predictive Models of Mariculture Impacts.....	20
- Environmental Regulations.....	21
- Need for Improved Feeds and Feeding Strategies.....	28
- Beneficial effects of Mariculture.....	28
Recommendations.....	29
Appendix 1: Country Reports.....	31
- Canada.....	31
- Denmark.....	34
- Federal Republic of Germany.....	36
- Finland.....	37
- France.....	39
- Norway.....	41
- The Netherlands.....	44 x
- Portugal.....	46
- Sweden.....	47
- United Kingdom.....	50
- United States.....	51
- USSR.....	52
Appendix 2: Agenda, Ad hoc Study Group, Hamburg, Apr 7-9,1987.....	53
Appendix 3: Executive summary.....	55
Appendix 4: Literature Consulted.....	57

x Note, pages 44 and 45 to come later.

Abstract

Following the adopted Resolution CR 1985/2:37 (73rd Statutory Meeting of ICES), CR 1986/2:36 (74th Statutory Meeting) an *ad hoc* Study Group on "Environmental Impacts of Mariculture" was established in order to: (a) delineate the dimensions of the problem, and (b) recommend a course of action which will lead to a development of criteria and to a standard system of monitoring and reporting." Membership to this ad hoc ICES Study Group was assigned during 1986 with experts nominated from 11 member countries. The Study Group worked by correspondence and met in Hamburg (Federal Republic of Germany) between April 7 and 9, 1987 to prepare the report. The environmental issues addressed include (a) our present understanding of the effects of mariculture on natural microbial communities and on the possible spread of pathogens, (b) the possible changes in the natural populations of marine algae (phytoplankton and macroalgae), (c) the influence of sedimentation on the benthos, (d) the use of chemicals in mariculture, (e) site selection criteria to minimize environmental effects, and (f) the state of the art in developing predictive models of mariculture impacts. Further chapters discuss environmental regulations (as applied to mariculture situations) in various member countries, the need for improved feeds and feeding strategies, and the possible beneficial effects of mariculture. Finally, a number of recommendations are formulated, related to research needs to assess and minimize environmental impacts. Considering the increasing environmental issues that are emerging with the expansion of the mariculture industry, the establishment of an ICES Working Group on "Environmental Impact of Mariculture" is proposed and the terms of references for such a Working Group are presented. Country reports are presented in Appendix 1 and an extended literature list is included in Appendix 4.

Introduction

Aquaculture in marine and brackish waters has shown a recent, rapid expansion in most of the ICES member countries. The introduction of new technologies, the expansion of suitable areas for cultivation, the improvements in feed technology, and the demand for high-quality protein products have opened new opportunities for the mariculturist. In many regions, mariculture has presently been identified as the only growth sector within fisheries. Several predictive investigations estimate a world aquaculture production of about 25 million tonnes by the end of this century, which requires an annual growth rate of over 12 %. Most of this increase is likely to occur in brackish and marine farming.

Progress in aquaculture development, however, will always be accompanied by problems and constraints that occur with or even through the expansion of the industry. Mariculture systems are structured around the use of natural waters and often of the natural food chain. As such, they are an integral part of the environment, and consideration must be given to their wider environmental impact. Aquaculture effluents from conventional farming systems were, in the past, considered to be "clean" and "natural" and the possibility that aquaculture may affect the environment has largely been overlooked.

Like any other industry, aquaculture has the potential to generate pollutants which are continuously released into the natural environment. Ecological concerns can no longer be ignored and have become a risk factor for the industry itself. The aquaculture industry and natural resource managers are becoming increasingly aware that the heavy demand for new sites, and the expansion of existing operations, may soon require more stringent environmental controls and limitations on the industry. To deal with the rapid growth of the mariculture industry, means that public administrators deciding not to regulate growth of such an industry must take risks. They must prioritize and minimize the amount of time spent evaluating and completing tasks. This includes evaluation of the potential environmental consequences of growth.

There must be a limit to the number of marine farms (water-based systems) which can be accommodated in a given area, and due allowance must be given for other sectors of the community sharing and competing for the water and coastal resources. This is true for competitive or joint uses of various branches of the mariculture industry, operating in the same area (e.g. interactions between fish farms and nearby mollusc cultures).

Lack of understanding of the interactions between mariculture and the marine environment can be divided into three major areas which are crucial to the future development of mariculture:

- (1) Water quality criteria of the water resource which is potentially suitable for mariculture,
- (2) Water quality requirements within farming systems,
- (3) The external impact of mariculture on marine ecosystems, including effects which affect the mariculture operation itself.

Improved understanding in these areas could help to determine the most appropriate information that governments could, in the future, expect the mariculture industry to provide in relation to "environmental impact statements". This seems to be of particular importance, because the aquaculture industry itself will be the first to suffer any adverse effects of environmental degradation. It would also be useful if strategies could be developed to minimize waste outputs from aquafarms while at the same time seeking efficient means of utilizing locally enhanced nutrient inputs in a mixture of extensive and intensive culture systems.

For a number of years the "European Inland Fisheries Advisory Commission" (EIFAC) has addressed environmental issues related to freshwater fish-farm effluents and stressed the need to reduce their negative effects. In 1980 an EIFAC Working Party on "Fish Farm Effluents" was established (Recommendation 80/19; FAO Fisheries Report No.248). The most important tasks of this Working Party are: (1) to provide guidelines for the control of suspended solids originating in fish farms, (2) prepare guidelines to reduce inorganic and organic phosphorus loadings, and (3) to review potential problems arising from the use of therapeutic agents in aquaculture. An outline of the present situation in EIFAC countries has been published in a Workshop Report on fish-farm effluents (EIFAC Tech. Paper No. 41, 1984).

Following the adopted ICES Resolutions C.Res. 1985/2:37 (73rd Statutory Meeting) and C.Res. 1986/2:36 (74th Statutory Meeting) an *ad hoc* Study Group on the "Environmental Impacts of Mariculture" was formed in order to (a) delineate the dimensions of the problem, and (b) recommend a course of action which will lead to the development of criteria and to a standard system of monitoring and reporting.

Membership to this *ad hoc* ICES Study Group was assigned during 1986 with experts nominated from a number of ICES countries. An editorial meeting with three Study Group members took place in December 1986 to compile and briefly review the available literature and unpublished data reports and prepare them for detailed considerations at the Study Group meeting. Country reports are provided in Appendix 1. The Study Group met in Hamburg (April 7 to 9, 1987; Appendix 2) and worked also by correspondence. An executive summary is attached as Appendix 3 and an extended literature list is given in Appendix 4.

The Study Group discussed the terms of references and felt that at present it would be difficult to develop a course of action that would lead to a standard system of monitoring and reporting. It was decided, however, to present available information from various

member countries on possible approaches to study and evaluate environmental issues of mariculture and practical measures to minimize its impact. The present document does not attempt to provide a comprehensive review of the published information available to the Group but addresses a selected number of issues to outline key areas of concern and to identify research needs.

The Study Group noted inconsistencies in the use of various aquacultural terms and felt the need for proper definitions of terms frequently used by aquaculturists, scientists and administrators. These inconsistencies, if not removed, will lead to continued misinterpretation with the consequent establishment of inadequate regulatory requirements. Although impossible to develop appropriate definitions of terms during the Study Group meeting, it was felt necessary to agree on the rule of common use for several terms for the purpose of this meeting. The following definitions were adopted:

The **carrying capacity** of a defined area refers in ecology to the potential maximum production a species or population can maintain in relation to available food resources within the area. It is necessary to distinguish this from the **holding capacity** of a defined area in which the potential maximum production is limited by a non-trophic resource. An example would be salmonid cage culture in an enclosed bay which is limited by oxygen availability - the holding capacity being the maximum production which can be supported without dissolved oxygen limiting production. **Assimilative capacity** in this example would be the property of the bay to maintain a dissolved oxygen concentration above a defined limiting level. **Hypernutrification**: This term has been defined by ICES (ICES,C.M. 1984 (E:12) as any substantial and measurable increase in the concentration of a dissolved nutrient. **Eutrophication**: Any increase in primary production by phytoplankton as a result of hypernutrification.

In addition, there are a number of other confusing definitions in the aquaculture literature which are mainly related to system performance and water quality criteria. Most of these terms have recently been defined in an EIFAC Report on "Terminology, formats and units of measurements" as they relate to flow-through and recirculation systems (EIFAC Tech. Paper 49:1-100, 1986). Other definitions can be found in the "Glossary on Aquaculture Terminology" (ICES, C.M. 1986/F: 34). Additional clarifications on frequently misused terms in aquaculture is required, particularly "Person Equivalent Values" (PEV) and "oxygen availability" (often derived from weight to volume ratios without considering partial pressure).

The present document constitutes to report of the *ad hoc* Study Group whose composition was the following: Harald Rosenthal (Chairman, Federal Republic of Germany), Hans Ackefors (Sweden), Jan Aure (Norway), Edward Black (Canada), A.C. Drinkwaard (The Netherlands), L.A. Dushkina (USSR), Arne Ervick (Norway), Richard Gowen (Scotland), Maurice Héral (France), Timo Mäkinen (Finland), M.A. Sampayo (Portugal), Donald Weston (USA), and David Wildish (Canada). Markku Pursiainen (Finland) participated in the Study Group Meeting as observer of the EIFAC-Working Party on "Fish Farm Effluents".

The final report presented here includes numerous comments and suggestions provided by Study Group members and by others. Those who submitted written comments and suggestions but did not belong to the *ad hoc* Study Group were: J. Brown (Dep. Biol. Sci., Simon Fraser Univ., Burnaby, Canada), N. Dankers (Rijksinstituut voor Natuurbeheer, Texel, The Netherlands), L. Håkanson (University of Umeå, Sweden), G.H. Hall (Windermere Laboratory, UK), C. Nielsen (Nykøbing Falster, Denmark), and Ø. Vethe (Institute for Georesources and Pollution Research, Norway). It should be emphasized, however, that the editorial committee (Rosenthal, Black, Gowen, Weston) accept responsibility for the views expressed in the present report.

Environmental Issues

Bacteria

Our understanding of the effect of mariculture on natural microbial communities is extremely limited, despite the fact that several marine microbes are potentially pathogenic to the cultured organisms, the indigenous biota or man. Most microbial work done in connection with mariculture has been directed towards identification of those organisms producing disease in cultured finfish and towards the development of therapeutic methods to prevent their spread. In comparison, much less work has been done in the area of microbial ecology, and specifically, identifying how culture operations modify the environment so as to change the number and type of bacteria.

The majority of bacteria in both freshwater and marine coastal and estuarine waters are heterotrophic, and the number of heterotrophs is largely determined by the concentration of organic matter (Fletcher, 1979). Thus, the input of organic matter associated with mariculture, and particularly the enrichment of bottom sediments, could reasonably be expected to lead to an increased number of cultivable bacteria in the vicinity of culture sites and to result into selective enhancement of specific bacterial groups.

The effect of culture activities on coliform, and specifically, faecal coliform bacteria is of particular interest because of the importance of this group in water quality monitoring. Finnish authorities have found elevated concentrations of coliform and faecal streptococci bacteria in effluents from freshwater trout ponds (Haavisto, 1974). Subsequent investigations (Niemi and Taipainen, 1982) found that the fish intestines and the sediments of the mud-bottom ponds can serve as multiplication sites for Enterobacteriaceae (*E. coli*, *Entero-* and *Citrobacter*), *Aeromonas* and faecal streptococci which enter the culture systems through the influent water and/or feed. The total number of indicator bacteria in the effluents was low, but detectable in the receiving waters. If oysters were cultivated in the effluent of fish farms they accumulated the enterobacteriaceae, the faecal streptococci, and also *Aeromonas* (Coeurdacier, Héral et al., 1983). Increased total coliform concentrations have also been reported in receiving waters near several freshwater hatcheries in the western United States (Hinshaw, 1973), although a large freshwater trout farm in Norway caused no change in faecal coliform numbers in receiving water (Bergheim & Selmer-Olsen, 1978). Near-surface waters near marine salmon net-cages in Sechart Inlet, British Columbia had a higher total coliform concentration than comparable reference areas, although the increase was not statistically significant (E. Black and B.L. Carswell, unpubl. data). We are in need of a clearer differentiation between faecal and total coliform, because the counts obtained with methods usually employed to evaluate sewage effluents and waste waters originating from human activities, may not adequately reflect the species and type composition encountered in fish farm effluents. The development of sound hygienic criteria, specifically adapted to the marine environment is urgently needed.

The most extensive work on microbial dynamics of fish farms is that of Austin and Allen-Austin (Allen, et al., 1983; Austin, 1982; 1983; 1985; Austin and Allen-Austin, 1985). Freshwater trout rearing facilities have been shown to have little effect on the bacteria populations of the receiving waters. In most instances bacteria numbers (CFU, = Colony Forming Units) from the effluent from these facilities are comparable or less than numbers in the influent. After investigation of 6 farms, in only 8 out of 29 comparisons was the number of bacteria in the effluent substantially higher (two-fold or more) than the number in the influent (Allen, et al., 1983; Austin, 1985; Austin and Allen-Austin, 1985). There was also little effect of culture on bacterial composition, with the taxa of the effluent generally typical of unpolluted freshwater. The most significant effect of culture was the selection for antibiotic resistant strains during

chemotherapy. The use of oxolinic acid, potentiated sulphonamide and oxytetracycline all increased the proportion of resistant bacteria in the effluent (Austin, 1985). This phenomenon was short-lived, however, for after 9 days or less following cessation of treatment, the proportion of antibiotic resistant bacteria in the effluent returned to pre-treatment levels.

Austin concluded that unlike freshwater finfish culture, marine culture may result in a net contribution of certain types of bacteria into the receiving water (Austin, 1982; 1983; Austin and Allen-Austin, 1985), although it should be recognized that this conclusion is based on studies at a single farm site and may be more representative of facility management rather than a general pattern. In each of four sampling periods, the number of bacteria in the effluent from a coastal turbot rearing facility was substantially greater than the number in the influent, with the increase ranging from 3 to 50-fold, mainly indicating growth of naturally occurring bacteria. There were also compositional changes in the bacteria including a decrease in the proportion of *Acinetobacter calcoaceticus*, *Hyphomicrobium*, *Hyphomonas*, *Micrococcus*, *Photobacterium*, *Pseudomonas*, *Staphylococcus* and *Vibrio* sp. and a commensurate increase in *Bacillus* spp., coryneforms, Enterobacteriaceae and *Prosthecomicrobium* spp. in the effluent. The Gram-positive bacteria are normally associated with sediments, and the Enterobacteriaceae with faecal pollution.

It is not known whether changes in bacterial composition or numbers in the vicinity of a mariculture pose an increased risk of disease in indigenous biota. Several investigators have raised the possibility of disease transmission from cultured to wild fish, and questioned whether fish pathogens in the effluent from mariculture facilities might infect wild stocks in the vicinity (Austin and Allen-Austin, 1985; Odum, 1974). There are cases of mariculture introducing a disease organism into an area where it had not previously occurred, eventually leading to the infection of wild fish (Rosenthal, 1980). There are, however, no documented cases of mariculture leading to increased incidence of disease in wild fish by a pathogen already present in the environment, but demonstrating no clinical symptoms. Many fish diseases are associated with some form of stress, including vibriosis (Colwell and Grimes, 1984), enteric redmouth (Austin and Allen-Austin, 1985; Bullock and Snieszko, 1975), and fin rot (Minchew & Yarbrough, 1977; Ziskowski & Murchelano, 1975). Deterioration in water quality, poor husbandry or other stresses imposed on fish in the culture environment can increase their susceptibility to diseases caused by opportunistic pathogens (Egidius, 1984; Sindermann, 1984), but appearance of the disease in culture, or even occurrence of the bacterial pathogens in the effluent, does not necessarily pose a risk to healthy, unstressed wild fish.

Studies on bacterial growth and species composition on molluscan culture sites are few. Recently, a Swedish study found no significant differences in bacterial counts and nutrients at sampling point outside and inside a mussel farm, which was located in a well protected inlet with strong tidal currents (up to 10 cm/sec) (Hagström & Larsson, 1985). Since this study was undertaken in April when water temperature is relatively low, the situation may be different during the most productive summer period.

Marine Algae

The Study Group recognizes that mariculture activity could bring about changes in the natural population of marine algae (phytoplankton and macroalgae) in the vicinity of the farm or at a distance where flushing action is vigorous. These effects include stimulation of primary production, changes in the species composition of the phytoplankton, reduction of phytoplankton standing crop, senescence of phytoplankton blooms with subsequent localized areas of low dissolved oxygen, and enhancement of macrophyte growth. These effects have implications for public health, natural populations of marine organisms, and the viability of mariculture industries.

Finfish Culture

By-products of fish metabolism and feed leachates represent a source of nutrients for phytoplankton growth (Gowen & Bradbury, 1987) and could, given suitable hydrographic conditions, stimulate primary production of phytoplankton if nutrients are controlling growth at the time. The input of soluble waste into the water column will alter the natural composition of macro- and micronutrients and it has been suggested that this can bring about changes in the composition of the phytoplankton (Takahashi & Fukazawa, 1982). Organic waste from fish farms might also play a role in stimulating the growth of specific algal species. For example, Nishimura (1982) has shown that, fish farm by-products (fish flesh and faeces) enhanced the growth of at least one red-tide forming dinoflagellate. Biotin has been implicated in the effects of *Gyrodinium aureolum* (Turner *et al.*, in press) and vitamin B₁₂ is a growth requirement for the toxic microflagellate *Prymnesium parvum*. Biotin and vitamin B₁₂ are constituents of fish feed but their fate in the marine environment is poorly understood. Recent British studies showed that Vitamin B₁₂ in fish feed made farmed fish more susceptible to *Gymnodinium* attack. *Gymnodinium* is as such not toxic to fish but affects mainly the oxygen transfer across gills through sloughing of gill membranes.

There are three mechanisms by which phytoplankton blooms might become a problem for mariculturists : (1) Depletion of dissolved oxygen (DO) in and around culture facilities. During dense blooms, oxygen consumption by algae during the night might reduce the concentration of DO to a level at which cultivated fish are asphyxiated. Reduction in DO during the senescence and anaerobic microbial degradation of large phytoplankton blooms might also have similar effects (Poole *et al.*, 1978). (2) Physical damage to sensitive tissues. Blooms of chain-forming species of *Chaetoceros* are known to have caused physical damage to the gills of farmed fish (Bell, 1961). It seems likely, however, that many species of diatoms which have long spines and some of the large armoured dinoflagellates could have similar effects. (3) Formation of algal toxins and subsequent uptake and accumulation by cultured organisms, or direct toxic effects on the cultured organisms. Shellfish toxicity (PSP and DSP) is a serious problem in many ICES member countries. Toxic algal blooms are known to have killed wild fish (A. White, unpublished data) and have been implicated in mass mortalities of farmed fish in European waters (Doyle, *et al.*, in press; Jones *et al.*, 1983) and shellfish (Buestel *et al.*, 1986). These problems are the subject of an ICES Working Group on Exceptional Algal Blooms and is not discussed in this report.

The formation of anoxic sediments beneath mariculture operations could provide suitable conditions for the accumulation of dinoflagellate cysts by preventing excretion. Dispersal of this sediment, by natural causes or for husbandry reasons, could result in the release of cysts into the water column and, under suitable hydrographic conditions, may result in the development of a bloom. Some coastal regions might be areas of natural cyst accumulations, and mariculture in these areas might be inappropriate.

The possible impact of mariculture on macroalgae is not well understood. The importation of exotic species, particularly shellfish, caused the introduction of new algae which can be considered as pests (e.g. the case of *Sargassum muticum* on the French and British coasts; see documents of the ICES Working Group on Transfers and Introductions). Their growth tend to replace some native algae. Development of mariculture (fish and shellfish) increase the release of nutrients (such as ammonia and phosphorus) and provides often additional substrate for attachment (e.g. cages, nets, racks, etc.). As a consequence, a large growth of green macroalgae can cause increased fouling on various substrates. The widespread dispersal of their spores can induce "green tides" near the culture installations, particularly in enclosed bays, in marshes and in ponds. These algae act in competition for nutrients with phytoplankton stocks, and this can have implications for the cultivation of molluscs. "Green tides" can also cause large decreases in dissolved oxygen.

Shellfish Culture

Extensive culture of bivalves may affect phytoplankton levels in one of two ways. First, bivalve culture will decrease the amount of phytoplankton. Imai (1977) clearly demonstrated that the culture of 50,000 to 90,000 oysters reduced concentrating seston (dominated by phytoplankton) by between 76 to 95%. Furthermore there is laboratory evidence that the filtering efficiency of bivalves (mussels) can be so great as to effect the growth of other mussels downstream (Wildish & Kristmanson 1985). Attempt have been made to estimate the seston requirements for growth of mussels (Wildish & Kristmanson, 1979; Incze & Lutz, 1980; Tenore *et al.*, 1982; Rosenberg & Loo, 1983) and Japanese oysters (Héral *et al.*, 1983; Sutherland & Roland, 1986).

Secondly, the stimulation of primary production by filter feeding bivalves has also been predicted. Campos and Marino (1982), Kaspar *et al.* (1985), and Tenore and Gonzalez (1975) have suggested that phytoplankton productivity would be stimulated by an increase in the rate of nutrient cycling. A number of investigations have demonstrated that recycled organic and mineral nitrogen excreted by oysters and clams is utilized by several species of phytoplankton (Roberts & Vincendeau, 1987; Lannergren, in by Rosenberg & Loo, 1983). Arakawa (1973) was able to correlate the frequency of phytoplankton blooms in Hiroshima Bay with historical trends in oyster production in those waters. However, field evidence of increased primary production in the vicinity of shellfish culture is lacking (Hagstrom & Larsson, 1982).

Sedimentation and Benthos

Mariculture structures can influence sedimentation simply by modification of current velocity and direction. Like any other structure placed in a moving fluid, mariculture structures may alter the flow regime creating zones of both erosion and deposition. In areas of intensive culture, the effects of bathymetry or even coastline topography can be significant. Hanging and stick shellfish culture in the Philippines has increased sedimentation to the point that progradation of coastlines has occurred in some areas (Davis, 1956). Some of the most well-studied examples are the mollusc culture grounds of France (Ottman and Sornin, 1985; Sornin, 1979; Sornin, 1981). Oyster farmers using metal racks in the intertidal zone have been confronted with sediment accumulation beneath and between culture racks. Concurrent with elevation of the sediment surface, erosion around the rack legs has accelerated sinking of the structures. Similar problems have been faced by mussel culturists using the bouchot technique. The numerous poles used to support the mussels reduced current velocity by up to 50%, thus accelerating the deposition of fine sediments. It is estimated that 30% of the oyster and mussel farms of France face problems of active sedimentation, forcing occasional relocation and abandonment of the old beds (Sornin, 1979).

In addition to sedimentation attributable to changes in water flow, all forms of mariculture result in the production of organic-rich particulate wastes. Faeces (and pseudofaeces in the case of bivalves) typically comprise the majority of the particulate wastes. Given the intensity of culture achievable, the amount of particulate material produced can be considerable. A typical oyster raft in Hiroshima Bay, for example, holds 420,000 oysters. Over a nine month culture period such a raft will produce 16 metric tons of faeces and pseudofaeces (Arakawa, *et al.*, 1971). With about 1000 rafts in operation in Hiroshima Bay (Arakawa, *et al.*, 1973) the oyster culture will have a major impact on the sediment in the bay. The production of pseudofaeces has also been studied in connection with mussel culture in the Netherlands. For example (Misdorp *et al.*, (1984) estimated the rate of sedimentation caused by mussel culture as 10mm year⁻¹.

Finfish culture will not only generate faecal waste, but for those operations in which feed is provided, feed which is not ingested by the fish also contributes to the total particulate load. In salmonid culture the amount of unutilized feed can range from 1% to 30% (VKI, 1976; Penczak, *et al.*, 1982; Braaten, *et al.*, 1983; Gowen, *et al.*, 1985;). Faecal production for salmonids is typically between 25 and 30% of the ingested feed (Butz and Vens-Cappell, 1981; 1982). Solid waste from salmonid farms will vary, depending on culture techniques and husbandry practices, but it has been estimated that the production of 1kg of salmon or trout will generate between 0.5 and 0.7kg of particulate waste (Weston, 1986a; Gowen and Bradbury 1987). The fate of particulate wastes will depend upon their settling velocities and hydrodynamic conditions in the area, but generally a large proportion of the wastes, particularly uneaten food will be deposited in the immediate vicinity of the culture site (Gowen *et al.*, 1985; Weston, 1986a). In low turbulent environments wastes accumulate typically as a soft, flocculent layer up to 40cm or more in thickness (Ervik *et al.* 1985). The deposition of organic wastes from fin and shellfish culture can result in physical and chemical changes in the substrate including:

- Increases in organic carbon (Dahlback & Gunnarson, 1981; Hall & Holby, 1986);
- Increased sediment oxygen consumption rates (Hall & Holby, 1986; Kaspar, *et al.*, 1985; Pamatmat, *et al.*, 1973);
- Decreased sediment redox potentials (Dahlback & Gunnarson, 1981; Brown *et al.*, 1987);
- Generation of hydrogen sulphide and methane (Cho, *et al.*, 1982; Hall & Holby, 1986; Liefbrig, 1985);
- Increases in organic and inorganic nitrogen content (Kaspar, *et al.*, 1985);
- Increases in phosphorus (Enell & Lof, 1983);
- Increase in silicon (Hall & Holby, 1986);
- Increase in calcium, copper and zinc (Aulio & Häkkinä, 1986)

The accumulation of organic-rich sediments and consequent physical/chemical changes in the vicinity of the culture site can have significant adverse consequences for the culturist. For example culture of shellfish in shallow embayments in Japan has led to nutrient and organic enrichment of the bottom sediments and consequent declines in culture productivity (Uyeno, *et al.*, 1970; Arakawa, *et al.*, 1971 Takagi, *et al.*, 1980). Clam (*Anadara sp.*) growers in Japan have adopted a strategy of bed rotation. After harvest a period of 1.5 years is required before the beds are reseeded. During this time the area is repeatedly trawled to accelerate remineralization (Terashima, *et al.*, 1972).

Finfish culturists have also been confronted with fish mortality and declines in productivity associated with the accumulation of particulate waste. The accumulation of enriched sediments may be harmful to the fish either because of outgassing of hydrogen sulphide, or the depletion of dissolved oxygen in the overlying waters. Furthermore, deterioration of water quality associated with enriched sediments may increase susceptibility to diseases (Arizono, 1979). Some Norwegian salmon farmers have found it necessary to disperse the accumulated sediments with propellers, or rotate cages among several sites, thereby allowing time for recovery of the sediment (Braaten, *et al.*, 1983).

In addition to the effects on the cultured organisms themselves, the accumulation of organic-rich sediments has pronounced effects on the benthic macro-invertebrates. In

cases of high organic input the sediment might become azoic but such an effect is usually limited to the sea bed directly beneath the culture structures (Hakilella, 1982, 1984; Junppanen, 1986; Brown *et al.*, 1987). At a greater distance generally less than 30m (Weston, 1986a) the effects are manifested by a proliferation of opportunistic species and a loss of many species intolerant of the physical/chemical consequences of organic enrichment (Kitamori, 1977; Tenore, *et al.*, 1982; Ervik, *et al.*, 1985; Kaspar, *et al.*, 1985; Lopez-Jamar, 1985; Brown *et al.*, 1987). The spatial and temporal effects of mariculture on benthic communities are comparable to those reported from other sources of organic enrichment (Pearson & Rosenberg, 1978; Poole *et al.* 1978).

A good example of the effects of mariculture on the benthos is provided by studies of a Swedish mussel site where the deposition of faeces and pseudofaeces created several centimeters of sediment each year (Mattson & Linden, 1981). Within six months after the start of culture, brittle stars had disappeared and species originally dominant in the unimpacted community decreased in number and finally disappeared after 15 months. Opportunistic species became established in the culture area concurrently with the decline of the original fauna. Within six months, large populations of *Capitella capitata* were established, and the species later reached densities as high as 20,000 individuals/m². Other opportunistic polychaetes (*Scolecopsis fuliginosa* and *Microphthalmus szcelkowi*) appeared after one year of culture. Total abundance and biomass decreased initially, but then fluctuated widely depending on densities of the opportunistic polychaetes.

Mattson and Linden (1983) also monitored the recovery of the benthos after removal of a mussel longline that had been in production for three years. Six months after removal the bottom was still covered by 20-40 cm of mussel shells and sulphide-rich sediments. The benthos was numerically dominated by opportunistic species indicative of disturbance. Monitoring continued for a year and a half after mussel removal, during which only very limited macrobenthic recovery was observed. In relation to salmonid culture, Gowen (unpubl. data) found a decline in macrobenthic species richness and biomass three months after initiation of culture. Following removal of the cages, sedimentary conditions showed improvement after three months, but the macrofaunal community showed evidence of substantial alteration relative to reference stations even eight months after cessation of culture.

Chemicals in Aquaculture

A wide variety of chemicals are used in mariculture which, if unrecognized or misused, represent a potential threat to the health of the cultured organism, the indigenous biota or the human consumer. An example of the range of chemicals available to culturists is provided by Schnick, *et al.* (1985) (Table 1). Twenty-four chemicals are registered or approved for use in food fish culture in the United States. The diversity of chemicals available to culturists is far greater in other countries where chemical licensing procedures are less rigorous. Several chemicals which are in use elsewhere have not been registered in the United States and efforts to do so have been terminated because of their potential toxicity, and concerns about human safety. For example furazolidone, nitofurazone, carofur, chloramphenicol and silvex have been dropped from consideration because they are known or suspected carcinogens. In other countries some of these chemicals are widely used. On the other hand, the use of TBT has not yet been banned in the United States while other countries have already restricted its application.

The chemicals in use can generally be divided into three groups, and it is in terms of this grouping that their usage is discussed: (1) Biocides and biostats which are deliberately introduced into the culture system, with the intent of eliminating predators or protecting the health of the cultured organisms (e.g. therapeutants, pesticides); (2) Chemicals which are introduced in the construction materials. They are not used with the intent to affect the health of the cultured organisms, and should there be an effect, it

may often go unrecognized. (3) Hormones used to alter reproductive viability, sex or growth rates.

The following discussion of chemical usage in mariculture is intended to illustrate the types and magnitude of chemical usage, and therefore the need to understand of the environmental consequences. It should be noted that with the possible exception of Tributyltin (TBT), there is no evidence that the chemicals in general use in mariculture pose a threat to the environment and/or human health. However our understanding of the environmental chemistry and toxicology of these chemicals is too limited to dismiss the issue without further consideration and the need for additional research is clearly apparent.

Biocides and Biostats

There is a long history of chemical usage in mariculture to control predators. For example, in the United States, lime was used as early as the 1930s to control oyster predators (Loosanoff, 1961). Trichloroethylene and dichlorobenzene have also been used to control oyster predators (Loosanoff et al., 1960). The pesticide Sevin is currently used in the northwestern United States to control burrowing shrimp in oyster beds. Copper sulfate has been used by the Japanese to control oyster predators (Koganezawa, 1978), but was discontinued when it was shown to adversely affect oysters.

In areas of intensive culture development the amounts of these chemicals used can be considerable. In the case of antibiotics for example, the 1984 antibiotic consumption by the Norwegian marine salmon culture industry included 6223Kg oxytetracycline, 7820Kg Tribissen R, 5500Kg Nitrofurazolidone and 9Kg sulphamerizine (Midtlyng, 1985). Considering the total production of salmon in Norway, however, the application of chemicals amounts to only 430g per tonne produced or half of this amount when taking the standing biomass into account. In recent years the use of antibiotics in Norway has been reduced and it should be noted that although there is less information from other countries the use of these chemicals in other countries is probably similar to Norway. The use of chemicals in aquaculture in the United Kingdom is summarized in Table 1a based on a survey of 148 fish farms (Solbe, 1982). Ninety-nine of the farms reported some chemical use with malchite green and formalin being used most extensively. Although the United Kingdom survey was based on freshwater facilities, most of the chemicals could be used in land-based marine culture systems.

The use of the chemicals in this group is widespread and intensive despite the fact that little is known about the environmental fate and effects of most of these substances. Environmental issues associated with use of these chemicals have often been raised (Pedersen, 1982; Solbé, 1982; Anonymous, 1983; Beveridge, 1984; Midtlyng, 1985), but rarely resolved. To date most attempts to consider potential environmental effects have relied on supposition based on dilution (Beveridge, 1984), persistence (Ackefors & Sodergren, 1985) or water solubility (Weston, 1986). With one exception field data are limited. Austin (1985) monitored antibiotic resistance of bacteria isolated from trout farm effluents. During treatment with oxytetracycline 90% of the bacteria strains examined showed antibiotic resistance. Within nine days after cessation of treatment, however, all resistance had been lost. Austin (1985) suggested that antibiotic resistance may be a short-term phenomenon. However, since the samples were obtained from effluent from a flow-through system, it is possible that there was a dilution effect rather than loss of resistance in the same bacterial population over time. Antibiotic resistance has been shown to be stimulated by antibiotic use in fish culture (Aoki, *et al.*, 1980) but more work on the subject is urgently needed. The development of antibiotic resistance is obviously of significance to the culturist, but it is only one of the environmental issues for which there is at best limited data. Other issues of equal concern include persistence of antibiotics and other mariculture chemicals, bioaccumulation potential, and toxic effects on indigenous biota.

Chemicals introduced via construction materials

Their presence is either unknown to the culturist (e.g., plastics additives) or they have been intentionally introduced but with the expectation that their use will have no effect on the cultured organisms (e.g., antifoulants). Plastics contain a wide variety of additives including stabilizers (fatty acid salts), pigments (chromates, cadmium sulfate), antioxidants (hindered phenols), UV absorbers (benzophenones), flame retardants (organophosphates), fungicides and disinfectants (Zitko, 1986). Many of these compounds are toxic to aquatic life, although some protection is provided by their low water solubility, slow rate of leaching, and dilution. Nevertheless, mortalities in mariculture have resulted from toxicants leaching from plastic construction material and the environmental effects of these toxicants beyond the confines of the culture facilities remain largely unresolved.

Antifouling compounds provide classic examples of chemicals associated with construction materials which can have profound effects on both the cultured animal and the indigenous biota. For example Tributyltin (TBT) compounds have been the subject of recent intensive study. TBT was first introduced as an antifouling paint in the mid-1960s, and was widely utilized because of many attractive properties (high toxicity to fouling organisms, low toxicity to man, lack of colour thus imposing no restrictions on paint colour, and ability to provide several years protection without pretreatment). The first indication of the effects of TBT on non-target organisms were noted in Arcachon Bay, France where 10% of the country's oyster production is carried out in close proximity to a large number of pleasure craft. In the mid-1970s oyster growers began to report poor spat collection, larval abnormalities and shell malformation. Subsequent work implicated TBT as the causative agent (Alzieu, *et al.*, 1980; 1982; Alzieu and Héral, 1984; Alzieu and Portmann, 1984; His and Roberts, 1983).

TBT has since been found in heavily utilized harbours throughout the world, and has repeatedly been shown to cause reproductive failure or growth abnormalities in molluscs, and has a high toxicity to many other forms of marine life (Cardwell and Sheldon, 1986; Cleary and Stebbing, 1985; Grovhoug, *et al.*, 1986; Maguire, 1986; Paul and Davies, 1986; Stang and Sellgman, 1986; Thain, 1986). TBT has been used extensively as an antifoulant on cages used for salmonid culture, but recent evidence indicates that the compound can cause mortality of the cultured fish and accumulate in the tissues, thus providing a route for human consumption (Short and Thrower, 1986a; 1986b, 1987). The rapidly accumulating evidence of adverse environmental effects of TBT has led to reductions in its use in mariculture facilities, restrictions on its use in France, United Kingdom, Federal Republic of Germany and Switzerland, and an ongoing re-evaluation of the compound in the United States.

Hormones

The final group of chemicals in aquaculture are hormones which are used to alter sex, reproductive viability or growth of the cultured organisms. The work of Goudie *et al.* (1986) and Johnstone *et al.* (1983) have reviewed the elimination from fish of a hormone commonly used to control sexual differentiation (Methyltestosterone). Both studies concluded that at those levels of use necessary to control sexual differentiation this hormone posed no potential human health hazard.

Much too little is known on bacterial degradation of hormones, biocides, biostats and antibiotics in the marine environment and studies in this field are urgently needed.

Table 1 : Chemicals registered or approved by the U.S. Food and Drug Administration for use in food fish culture¹ (Adapted from Schnick, et al., 1985)

Product	Use
THERAPEUTANTS	
Acetic acid	Parsiticide
Formalin	Parasiticide and fungicide
Romet 30 (sulfadimethoxine and orthomeprim)	Bactericide
Salt	Osmoregulatory enhancer
Sulfamerizine	Bactericide
Oxytetracycline (Terramycin)	Bactericide
ANESTHETICS	
Carbonic acid	Anesthetic
MS 222 (tricaline methane-sulfonate)	Anesthetic and sedative
Sodium bicarbonate	Anesthetic
DISINFECTANTS	
Calcium hypochorite	Disinfectant, algicide and bactericide
WATER TREATMENT	
Fluorescein sodium	Dye
Lime (calcium hydroxide, oxide or carbonate)	Pond sterilant
Potassium permanganate	Oxidizer and detoxifier
Rhodamine B and WT	Dye
Copper sulfate	Algicide and herbicide
Copper, elemental	Algicide and herbicide
2,4-D	Herbicide
Diquat dibromide	Algicide and herbicide
Endothall	Algicide and herbicide
Simazine	Algicide and herbicide
Clean-Flo (aluminum sulfate, calcium sulfate and boric acid)	Algicide and herbicide
Glyphosate	Herbicide
Potassium ricinoleate	Algicide
Xylene	Herbicide

¹The FDA often registers or approves a compound under a trade name. For example there are eight approved or registered algicides/herbicides containing copper sulfate. A single entry chemical name has been provided in the table for these cases.

Table 1a : Chemical usage in freshwater fish farms in the United Kingdom
(Adapted from Solbé, 1982)

	No. of farms (out of 148 surveyed)	Total amount used per year (sum of all farms)	Average Effluent Concentration (mg/l)
Malachite Green	89	176 kg	0.61
Formalin	56	1230 L	15.2
Hyamine	23	77 L	0.55
Oxytertracycline/Terramycin	16	0.7 kg	
Chloramine T	12	30 kg	
Sodium chloride	12		
Copper sulfate	11		
Tribrissen/Sulphamerizine/Vesadin	10		
Betacide/Marinol/Roccal	9		
Iodophores/Vescodyne/Pevedine/ Vanadine/ Polyvinylpyrrolidone-Iodine complex	9		
Furazolidone/Neftin	8	38 kg	
Buffodine	5		
Methylene Blue	4		
MS 222	4		
Potassium permanganate	3		
Acintrasole	2		
Acriflavine/Pioflavine-Hemisulphate	2		
Dipterex	2		
Enheptine	2		
Slaked/Quick lime	2		
Sodium hydroxide	1		
Sodium hypochlorite	1		

Water Quality

Information on the effect of marine fish farms on water quality in and around culture facilities is insufficient to allow for a detailed evaluation. For freshwater situations, the European Inland Fisheries Advisory Commission (EIFAC) of FAO organized a workshop (Alabaster, 1982) to assess the extent of the problem in freshwater aquaculture in central Europe (Rosenthal, 1983). An analysis of the information gathered through a questionnaire, for which replies were received from 15 member countries (Alabaster, 1982), revealed that fisheries downstream from farms are not adversely affected unless the total flow of recipient waters is less than 5 L per second for each tonne of annual fish production. There was a general tendency for pH to drop slightly when water passed through the farming system (average 0.2 pH units at 1L/sec/t annual production; no change at 4 L/sec/t annual production or higher flow rates). At higher dilutions fresh water farm effluents are still associated with increased eutrophication. Inorganic phosphorus was considered to be the major contributor to the eutrophication effect observed in receiving waters (Sumari, 1982). Oxygen levels in the outlet of farms decreased generally at an average of 1.6 mg/L for an average flow of 12.6 L/sec/tonne annual production. This figure is based exclusively on data obtained from salmonid farms.

Most of the water quality criteria examined in the EIFAC study showed an increase in concentrations (Table 2) when comparing inlet and outlet values. However, the effluent quality data available do not provide a consistent figure. BOD load (Biological Oxygen Demand) increased only with an increase of suspended solids.

Table 2 : Net increase in constituents of water passing through fish farms (Kg/day/t annual production) according to an analysis of an EIFAC questionnaire. SS= suspended solids; BOD= biological oxygen demand; COD= chemical oxygen demand; NH3-N = unionized ammonia-nitrogen; ND = no data available (modified after Alabaster, 1982).

Country	SS	BOD	COD	NH3-N	oxidized nitrogen	total phosphorus
Cyprus	1.3	1.4	ND	0.4	0	0,16
Denmark	1.4	2.1	1.0	0.2	0.1	0.06
Finland	1.5	1.5	ND	0.1	0.5	0.08
FR Germany	ND	ND	ND	0.5	1.5	0.27
Italy	11.0	1.5	ND	0.4	ND	0.31
Norway	3.5	2.7	31.0	0.3	0.9	0.14
United Kingdom (after Solbé, 1982)	4.1	2.4	ND	0.1	0.5	0.03
United Kingdom (after Purdom, 1982)	9.4	ND	ND	0.1	0.1	0.10

Selected water quality aspects

In this report several water quality issues have already been addressed in connection with other environmental considerations. Therefore, only a few aspects are presented and discussed here, placing the main emphasis on oxygen depletion at cage farm sites, the use and misuse of the unit "Person Equivalent" as a means of fish farm waste water evaluation and on the concentration of metabolic nitrogen (such as total ammonia) at water-based farm sites. Table 3 summarizes data from several studies available during the Study Group meeting. They do not provide a comprehensive overview but demonstrate that the net loading for various water quality parameters differs substantially for the same species, and is dependant on site specific conditions, feeding strategy, feed composition and other operational criteria. The dependence of the release of total suspended solids from fish farms on feed conversion efficiency, and on growth rate has clearly been demonstrated in fresh water aquaculture (Knösche, 1971, Table 3). While information on waste output is available for freshwater species and salmonids cultured in marine waters, data are lacking for potential mariculture species such as halibut, turbot, cod and sablefish.

Table 3 : Estimated pollutional load derived from observations at various commercial intensive fish farms using pond and tank system.

System	Species	size	feeding rate (%) bodyweight	feed type	Net loading	Reference
Tanks and ponds	Brown trout (biomass 2,260kg)	2.2g - 100g	1.3 - 17.5	dry pellets	11.5g COD/kg fish/24h	Bergheim et al. (1982)
				manual + automatic	2.7g BOD ₇ /kg fish/24h 0.05g Total P/kg fish/24h 0.9g SS/kg fish/24 h	
Tanks and ponds	Brown trout (biomass 7,320 kg)	1.0 g - 25.0 g + brood stock	3-6 %	dry + wet pellets, manual	75.3 gCOD/kg/24 h 83.3 gBOD ₇ /kg/24h 0.43g Total P/kg/24h 0.24g PO ₄ -P/kg/24h 1.4-3.8 g Total-N/kg/24h	Bergheim et al. (1982)

Table 3 (continued)

ponds	Brown trout, rainbow trout (biomass 2.690kg)	0.2 - 500 g	0.55- 4.5 %	dry pellets manual	17.0 gCOD/kg/24h 7.1 gSS/kg/24h 0.45 g Total-N/kg/24h 0.08 g Total-P/kg/24h 0.05 g PO ₄ -P/kg/24h	Bergheim et al. (1982)
tanks and ponds	Brown trout (biomass 5,970g)	1-550g	0.5 - 16 %	dry, automatic	3.1 g COD/kg/24h 1.6 g BOD ₇ /kg/24h 1.2 g SS/kg/24 h 0.13 g Total-N/kg/24h 0.03 g Total-P/kg/24h 0.05 g P ₀ ₄ -P/kg/24h	Bergheim et al. (1982)
ponds	rainbow trout	35-150g 500-2000g	?	dry pellets	0.4-0.8 gTotal-N/kg/24h 0.05 g Total-P/kg/24h 1.6-4.6 g BOD7/kg/24h	Bergheim + Selmer-Olsen(1978)
ponds	rainbow trout	2.0 -300g	?	dry pellets, wet feed	0.5-1.4 g Total-N/kg/24 h 0.13-0.18 g Total-P/kg/24h 1.9-5.7 g BOD5/kg/24h	Markham (1978)
Tanks	lobsters	100-200g	?	?	1.43-2.10 g COD/kg/24 h	Tchobanoglous Shlesher (1974)

Table : Relationship between suspended solids loading and feed conversion efficiency in intensive carp culture. SS = suspended solids (dry weight), FC = Feed conversion factor, dry to wet weight basis. (modified after Knörsche, 1971).

FC	Tonnes SS per tonne weight gain	Suspended solids produced (g SS/kg Fish/day)	
		Feeding rate	
		2%/day	3%/day
1.0	0,68	13.6	20.4
1.2	0.86	14.3	21.5
1.5	1.13	15.1	22.6
2.0	1.58	15.8	23.7

Oxygen depletion

Oxygen availability at cage farm sites is a problem of immediate concern to the farmer. The best documented example for oxygen depletion in and around net cages is that of the Japanese large-scale yellowtail farming operation in the Usui Bay (Kyushu Island). In Usui Bay, farm operations cover an area of approximately 10 ha, employing about 250 to 350 cages, and producing annually about 750 tonnes of fish. Although tidal currents are relatively strong (3 to 4 knots, maximum), oxygen depletion occurred within the cages,

especially during slack tide, and in recent years farmers were forced to reduce the number of cages in the inner part of the Bay. Dissolved oxygen concentrations, in the central part of the cage-covered area, decreased by 40-60 ppb during a single pass of the water mass through a single cage (stocking density $6 \text{ kg} \cdot \text{m}^{-2}$). Between 0.3 and $0.5 \text{ mg} \times \text{L}^{-1}$ was consumed when stocking density was as high as $22 \text{ kg} \times \text{m}^{-2}$. There was also a substantial difference in levels of oxygen depletion between those cages installed close to the inshore and those further offshore in deeper waters. Figure 1 depicts isolines of oxygen concentrations measured in and around the cage culture facilities (samples taken approximately 1-2 m below water surface) prior to, during, and after feeding. It seems obvious that oxygen depletion and water exchange are largely related to both the shoreline structure and the arrangements of the cage units. Those cages located closer inshore and surrounded by other cage units are likely to be affected first by any oxygen depletion. Oxygen depletion may also be related to seasonal events, especially during the summer and fall, when warm water temperatures support faster growth rates and feeding levels are high, oxygen concentrations in center cages can be reduced to critical levels. The oxygen depleted area extends not only horizontally but also vertically, and this is shown in Figure 1a, which shows the vertical profiles in various farm operations in the same bay and at two different tidal situations. The impact on oxygen levels is largest in the yellowtail farms, followed by the Red Sea Bream cages, and least at the pearl oyster raft site. During several years of studies, the investigators (Kadowaki et al., 1978 a,b, 1980) also noted that the situation became gradually worse. This was partially attributed to the substantial amounts of oxygen-consuming substances accumulating at the sea bed under and around the cages.

Person Equivalent values (PE)

Fish farm waste water is not directly comparable to domestic sewage. The use of PE-values, if not specified for individual components, is therefore invalid. Most of the BOD produced in a fish farm is related to the suspended solid fraction which settles quickly. Soluble BOD loads have been estimated to reach $2.56 \text{ kg} / \text{tonne}/\text{day}$ in trout farms, while the settleable solid fraction amounts to $6 \text{ kg} / \text{tonne}/\text{day}$. Taking the average PE values as $12 \text{ g N}/\text{day}$, $2.5 \text{ g P}/\text{d}$ and $75 \text{ g Oxygen}/\text{day}$ (asBOD₇), land-based fish farms may produce between 21 and 60 PE loads per tonne of fish stocked in the system. Markmann (1978) reports for danish pond farms a BOD₅ load equal to 19 Person Equivalents, while the loading for N and P would equal 32 PEs. High loadings can be calculated for french trout farms, amounting to 0.2-05 PE per kg fish stocked (Fauré, 1977). These data show, that the comparison with untreated sewage is not valid, mainly because of the totally different C,P,N ratio and the significant difference in settleable and soluble wastes. From a management viewpoint it was felt unrealistic to use PE values as an index of the overall pollutional load in fish farming.

Site selection

In the context of this report site selection for the siting of mariculture operations is confined to a discussion of those criteria which might be used to minimize the environmental effects of an operation and does not include an assessment of site selection criteria to ensure optimal growth of the culture species. Neither does this section discuss site selection for the avoidance of harmful algal blooms, this topic is under review by the ICES Working Group on Exceptional Algal Blooms.

Mariculture operations interact with the water body within which they are located and the benthos over which they are sited. The balance of the impact, that is, which of the two is the main recipient of waste will obviously depend on the operation but it is the intensity




Fig. 1: Examples of the effects of feeding on oxygen depletion in large-scale cage culture facilities for yellowtail (*Seriola quinqueradiata*) and Red Sea Bream (*Pagrus major*) in the Usui fish farms: a = pre-feeding; b = during feeding; c = after feeding (by courtesy of Prof. Hirata and Dr. Kadowaki).

and scale of these interactions which will determine many of the broader ecological impacts. The following discussion is confined to those factors which limit the impact of a mariculture operation on the benthos and water quality. Some of these factors might be combined to formulate site selection protocols which would be of use in resource management but the formulation of such protocols needs careful consideration and is beyond the present scope of the Study Group.

Impact on the benthos

For finfish and some forms of shellfish culture (for example raft and longline culture of mussels) the same criteria for site selection can be applied. However, for bottom and near bottom culture of some shellfish, minimizing the impact on the benthos by site selection is more problematic. Careful husbandry practices can minimize the amounts of waste released by finfish farms, however the distribution and loadings of organic waste to the benthos will be dependent on a number of physical processes. In Norway, the possibility of shifting cages to alternative locations has been considered. However, nothing is known on the required recovery time of overexposed sites.

Sediment Type: Sediments can be divided into two broad groups, depositional and erosion. In general the former tend to be composed of fine particles and are indicative of regions where fines and organic detritus accumulate. In contrast erosional sediments are generally composed of coarse particles and indicate regions of the sea bed where there is active transport of fine particles. The identification of coastal regions with coarse grained sediments is one criteria for the selection of mariculture sites where the accumulation of organic waste would be reduced. However, grain size is also dependent on the source of the sediment material and a coarse sediment might be a reflection of the fact that there is not a local source of fines. In such cases, the sorting characteristics as measured by the sorting coefficient need to be identified (see in Holme and McIntyre, 1984: Methods for the study of marine benthos, IBP Handbook 16).

Bottom topography. Regions where sea-bed features promote accumulation of sediment should be avoided.

Water depth: The dispersion of faeces, pseudofaeces and waste feed is partly dependent on the depth beneath the operation. In general the greater the depth the greater the lateral transport of the waste particles. However, this is not always the case. In some areas (for example the deep basins of some fjords) current velocities might be low and result in limited lateral dispersion. Adequate depth under cage systems will also allow hydrogen-sulfide bubbles produced from anoxic sediment to be rendered harmless by the time they pass through the culture facility on the way to the surface (Ervik, pers. comm.).

Current velocity: Current velocities will determine the lateral dispersal of solid waste settling through the water. The higher the current velocity, the greater the lateral displacement of waste particles and hence the accumulation of organic waste will be minimal. Thus in general sites with high current velocities should be selected although it should be borne in mind that high current speeds might have adverse effects on finfish growth. In addition it should be noted that good tidal flow does not automatically lead to good water exchange and the retention time of water around the farm should be considered.

Existing benthic community: Coastal areas where benthic communities provide nursery grounds or natural beds for commercially fished species, and areas of marine conservation interest should be avoided.

Water Quality

Many of the criteria described above will have a bearing on the impact of mariculture operations on water quality. In general factors which promote good water exchange and rapid dilution of soluble waste materials are likely to be useful in site selection protocols.

Water movement: Good water flow through the culture structures will disperse soluble waste materials and reduce the possibility of localized oxygen minima. However, current patterns are also important. Soluble waste could accumulate in regions where there are eddies or gyres despite adequate water flow.

Retention time: The retention time (flush time or dilution period) of a semi enclosed water body has an important influence on the dilution of soluble waste. In poorly flushed locations soluble wastes are likely to accumulate and the demands of the mariculture operation on oxygen supply can result in localized minima which could affect the cultured organisms. It is in such poorly flushed locations that the ecological impacts of mariculture operations are likely to be greatest. In some coastal regions water in deep basins can persist for periods of time ranging from several months to years. In these locations the input of wastes(soluble and solid) from mariculture operations could enhance oxygen depletion and nutrient levels within the deep water. Upwelling of nutrient rich water into the euphotic zone could stimulate phytoplankton growth and the upwelling of oxygen depleted water could affect both the cultured organisms and indigenous species.

Water column stability: The stability of the water column can influence the volume of water into which soluble wastes are dispersed. For example in the case of mariculture dependent on floating structures the presence of a strong pycnocline immediately beneath the structures could result in limited dispersion of wastes. The effect of water column stability will depend on the duration of the pycnocline or thermocline, and the strength of such density discontinuities.

Natural levels of nutrients: It has been suggested that the release of soluble nutrients from mariculture operations could influence the ecology of phytoplankton populations within a water body, but only if nutrients are limiting phytoplankton growth at the time. The identification of coastal regions where phytoplankton growth is nutrient limited, particularly during the summer months might be a means of reducing the impact of soluble nutrients and possibly reducing the risk of mariculture operations stimulating the formation and persistence of harmful algal blooms.

Predictive Models of Mariculture Impacts

The Study Group recognized the importance of models for research into the interactions between mariculture and the marine environment and as tools for resource management. However to date the development and application of models in mariculture has been limited. Models can be divided into two broad groups, those dealing with sedimentation and those dealing with water quality.

Sedimentation Models

In attempting to gain an understanding of the effects of the deposition of organic waste from mariculture activities there have been two approaches. Firstly, the use of models for site selection, for example, Hakanson *et al.*, (1986) looked at physical and geomorphological features of coastlines in Sweden to identify areas of sediment deposition and erosion. Secondly, sedimentation models which predict the accumulation of sediment beneath a particular mariculture operation and thus allow an environmental impact assessment to be made. In their simplest form solid waste is assumed to settle as single independent particles with only a simple lateral component derived from water movement (Hagino, 1977). In relation to sedimentation beneath salmonid cage farms Gowen (unpubl.data) has taken this a stage further by distinguishing between faecal waste and uneaten feed pellets and relating deposition to variations in current speed and direction. However in both of these models the effects of upwelling, resuspension the maricultures themselves and the transport characteristics of flocs has been ignored. It may be possible to adapt models

which have been developed to determine the distribution of pollutants from other industries (for example pulp mill effluent Parker and Sibert (1973) and Sibert and Parker (1973)).

Water Quality Models

In relation to water quality it is necessary to distinguish between those models aimed at estimating the carrying capacity of a water body and models for assessing the holding capacity of a water body. In the case of shellfish culture mass balance conceptual models have been developed to quantify the flux of phosphorus and nitrogen between the water column, cultivated oysters and the sediment (Sornin *et al.*, 1986; Feuillet, unpubl.data). Models have also been developed to estimate the carrying capacity of a water body to optimize shellfish production (Roland, unpubl. data) or predict what environmental changes may cause (Smeal *et al.* 1986). These models also give an estimate of the impact of the culture operation on the food and hence the possible effects on natural populations.

Holding capacity models have been developed with the aim of minimizing adverse interactions between finfish culture operations and the water body within which they are located. Such models are generally based on estimating dilution of specific parameters (for example dissolved oxygen (Inoue *et al.*, 1966,1970; Sakamoto 1977; Kils, 1979; Pedersen, 1982). In their simplest form dilution models assume that the modelled parameter is diluted evenly in the water passing through a culture facility (Inoue *et al.*, 1966 ,1970; Incze & Lutz,1980; Beveridge, 1984). However these models are likely to find only limited application in coastal waters and those models which attempt to assess the impact on semi-enclosed water bodies and more open coastal waters are likely to be of more use. For example Beyer and Madsen (1974) used a box model approach to model the transport of phytoplankton, dissolved oxygen, phosphorus and nitrogen in a Danish fjord. Black (unpubl. data) has attempted to model the effect of salmonid farms on dissolved oxygen levels in a semi-enclosed water body. Existing models to predict the impact of pollutants from other industries might be of use in assessing some of the environmental impacts arising from mariculture operations (see Weston, 1986 and references cited therein).

Two types of models can be developed for simulating the carrying capacity of a bay for mollusc production. The first one provides a global approach based upon the evaluation of the dynamics of the cultivated species: biomass, production, growth rate (Héral *et al.*, 1986). The second model is an analytical model which can be sensitive to the variability of the environment. This model incorporates the flux of available food calculated with physical numerical models and the consumption, respiration, and assimilation of the molluscs is developed to predict Bivalve production (Bucher and Héral, unpubl. data).

Environmental Regulations

The available information on environmental regulations as applied to mariculture situations in ICES member countries is incomplete. The Study Group decided to compile the material available at the meeting under two aspects: (a) country specific information, and (b) environmental criteria.

Regulations in various countries

Canada

In Canada both the Federal and Provincial Agencies regulate mariculture industries- The Federal Department of Fisheries and Oceans and the Environment Canada regulate under the Fisheries Act and its requirement to protect the habitat necessary to maintain the wild fisheries and natural populations of fish and shellfish. The Provincial agencies are responsible for the actual allocation of water leases for culture purposes and also have a

role in habitat protection. At both levels of government there are, however, regional differences in the interpretation of what is required to accomplish these tasks.

On the west coast constraints on each application for a mariculture license are principally determined on the basis of site specific information. Regulating agencies require bivalve culture, principally of Pacific oysters, to locate where it will not conflict with traditional harvesting of wild bivalve populations and not within 125 meters of a wharf or finfish culture. The substratum under proposed finfish culture sites must be inspected to ensure no important fish habitat is being alienated. As an indicator of whether an area is likely to accumulate faeces and excess feed the local substrate type (fine muds, gravel, bedrock, etc.), is considered prior to permitting culture activities to proceed. It is also required that fish culture activities locate at a distance greater than 1 km from the mouth of salmon bearing streams and that fish culture leases be no closer together than 3 km.

The approach on the East coast is different. Production per farm has been limited to that amount which can be produced from 30,000 seed fish annually maintained on a lease of no more than 2 ha in size. This is expected to yield 75 to 100 tonnes of marketable fish per farm per year. Federal regulations on this coast require that weirs be sited no less than 305 meters from each other. The same separation distance is suggested for application to the siting of fish cage culture adjacent to weirs and is required as separation between fish farms.

France

Licences for private coastal aquaculture operations are issued by the State authorities. For oysters, mussels, and clams the licence is governed by a management plan. The plan describes the culture operation's stocking density, orientation of structures and the prevailing currents. The plan also describes those permits required for temporary and/or permanent installation of the structures. The density of the culture can be reviewed on a yearly basis, and depending on productivity new stocking densities may be imposed.

There are at present no regulations specifically for marine fish production, except that combined finfish and molluscan culture is not permitted.

Finland

In Finland there is no general guideline on site selection for net cage farms. At present, the access to a particular coastal area is the most important factor that determines site selection by interested parties. Licences are required for establishing farms over a certain size (which in practice covers all of the present operations), and these are granted by the water courts (Salonen 1986). Authorities (National Board of Waters) generally limit the yearly production of a farm to a level of 30 to 50 tonnes (Turun vesipiirin vesitoimisto).

Norway

Norwegian salmon cage farms are currently limited to a maximum cage volume of 8,000 m³ per facility. There is no upper limit to the stocking density per unit volume of cage.

The increasing production rate over the last few years, together with the unutilized production potential in existing farms, have resulted in a preliminary moratorium in the granting of new licenses.

The regulations set by the veterinarian authorities ask for a 1 km separation between fish farms in order to minimize the potential for the transfer of diseases or parasites between

sites. This regulation has been developed by arbitrary administrative decisions and is not based on any scientific data.

In recent years there has been a growing concern on the possibility of transfer of diseases and parasites from fish farms to wild salmonid stocks. There exists also a potential of genetic implications on local native salmon strains caused by intermingling of escaped farmed fish. The environmental authority has therefore established restricted areas for fish farming in fjord areas with natural salmon runs.

For new mariculture species (e.g., cod and halibut) there are currently no regulations as to the number of marine fish farms and on the minimum distance between individual farms. In the last year there has been an increase in the number of applications for marine fish farms licenses. Each license must be sanctioned by the appropriate authorities.

Portugal

Licenses for private coastal aquaculture operations are issued by the State Authorities after hearing a commission in which several Departments are presented.

Sweden

There are 3 to 4 acts which regulate aquaculture in Sweden. The County Administration examines applications with the provisions of the Environmental Protection Act and the Nature Conservation Act. The Swedish National Board of Fisheries examines applications in accordance with the provisions of the Fisheries Act. The latter act is concerned with the dissemination of diseases and the ecological consequences for the indigenous fauna if new species or foreign populations are introduced (Ackefors, 1983, Anonymous, 1983, Ackefors & Grip, 1985).

With regard to aquaculture, the basic concept in the Environmental Protection Act is to evaluate the optimal or most suitable site and to restrict the environmental impact when considering available technology. Considerations should also be given to economical facts for the farmer.

To establish a farming operations, applications for licencing are divided into two categories :

- (1) Farmers who intend to keep more than 10 tonnes of fish in their operations must send in their applications to the authorities. The application must first pass the Environmental Protection Act before a licence can be considered.
- (2) Farmers who intend to operate at a size between 0.5 to 10.0 tonnes of fish are only obliged to inform the authorities about the planned operations.

The application from the farmers must also be evaluated in accordance with the Nature Conservancy Act. This Act regulates the protection of the seashore. In principle, areas at a distance of up to 100 m offshore and a land area of the same width from the shoreline are protected against commercial use or settlement.

The mussel operations are also divided into two categories. A plant which covers an area of at least 20,000 m² (including space for moorings) must be examined according to the Environmental Protection Act. Farmers who intend to utilize an area between 5,000 and 20,000 m² are only obliged to inform the authorities about the planned operation.

Processing plants are also covered by the Environmental Protection Act if more than 200 tonnes of fish and shellfish are processed per year.

Applications for aquaculture can also be made through the Water Rights Act. Under specific circumstances, this Act is applicable to landbased operations, in case the use of ground water or surface water is intended.

The application for aquaculture licences must be sent to both the County Administration and the Regional Fishery Office. These authorities send the application as referral to several authorities for detailed evaluation. Applications for big production units are usually handled through the Environmental Protection Board. This agency can reject decisions already made by the County Administration.

The aquaculture operations are included in the Environmental Protection Act because of their possible adverse effects on water quality. Discharging organic matter may also cause oxygen depletion in the water. The nutrient load in the coastal waters of the Baltic Sea, mainly from inputs other than mariculture, is already causing algal blooms. The ultimate effect of these is the depletion of oxygen in bottom waters.

The Swedish philosophy for regulating the aquaculture industry is based on the concept that the phosphorus and nitrogen load should be minimized to a level, which does not change the nutrient status of the receiving water. In freshwater the phosphorus content is checked against the Vollenweider diagramme to estimate the possible allowable amount of aquaculture production, which can be accepted with regard to the water volume and the water flow. Although the Vollenweider model cannot be applied in coastal waters, the concept presently employed is almost identical. A nutrient budget is normally made in order to evaluate the extra loading of nutrients through fish farming: However, we are still in need of more detailed knowledge on the hydrodynamics within the water bodies of concern, the amount of nutrient discharged into coastal waters as well as the transformation of nutrients into different parts of the ecosystem.

Water quality in farming areas is controlled by the Swedish Food Administration. In principle, the same rules are applied as for bathing waters with regard to the number of coliform bacteria. The farming is not permitted in "blacklisted waters". Aquaculture is prohibited in waters where wild fish contain high concentrations of environmental pollutants such as mercury (or other heavy metals) and chlorinated hydrocarbons. However, it is obvious from the present experience in such waters that cultured fish which do not depend on natural food but grow on a properly fed diet, do accumulate little of these contaminants.

The Swedish Food Administration has introduced rules concerning withdrawal time, when antibiotics and other therapeutics are used in aquaculture. The slaughter of fish is not permitted for a certain period of time after treatment. The time limit depends also on water temperature and on the specific properties of antibiotic.

United Kingdom

At the present time there are no regulations which require fish farms (existing and potential) to assess or monitor the ecological effects of their operation. The Crown Estate Commissioners who administer the sea-bed for the Crown are responsible for the issuing of leases to potential fishfarmers. The Commissioners have a consultative procedure before a lease is granted. This procedure involves obtaining statements concerning potential sites from a number of agencies including the British Nature Conservancy Council, but at present this consultative procedure does not include any assessment of the potential ecological impact.

Regulations regarding various environmental criteria

Sedimentation

In most types of mariculture, there are no regulations intended to prevent the accumulation of faeces, pseudofaeces or fine sediments in the area of the culture site. It is generally the growers' responsibility to rotate sites, till the sediment, or take whatever other steps may be necessary to maintain the suitability of the culture grounds. Finfish net-pen culture is the only form of mariculture in which governmental regulations exist to mitigate sediment accumulation.

French authorities have enforced regulations that determine the orientation of raft and tray structures for oyster culture in relation to the prevailing currents. The intent is to minimize sedimentation underneath and in between structures. The duration and time when structures have to be removed is also regulated, especially to utilize the strong winter currents to carry away accumulated sediment.

The potential for feed and faeces accumulation is generally an issue of concern in review of applications, although this review is typically qualitative without defined criteria. In countries where criteria do exist, they generally take the form of a required water depth for pen siting and a lateral separation from habitats or resources of special concern or which are especially susceptible to sedimentation impacts. British Columbia requires that net-pens be located in water depths of at least 10 m, be no closer than 125 m from a shellfish bed, and may not be located near eelgrass beds, herring spawning areas or other habitats of special significance. New Zealand requires that the cages be sited in water depths of at least 12 m, and that the bottom of the cages be at least 4 m from the seafloor. In the United States, the state of Washington utilizes a sliding depth scale which takes into account current velocity and facility production (see Figure 2). A distance of 6 to 18 m is required beneath the cages, with the greatest distance being required for large operations in areas of weak currents. Although not formally adopted as law in Washington, applications for salmon net-pen culture are evaluated with respect to these guidelines (Table 4).

Water quality

Land-based mariculture operations in which effluent is conveyed through a pipe or similar confining structure are subject to the most intensive environmental regulation. Discharge limits may be imposed for such parameters as Biological Oxygen Demand (BOD)




Figure 2 : Minimum depth and current guidelines for salmon net-pen siting in Washington State: Class I = production \leq 9 t/yr; Class II = 9-45 t/yr; Class III = $>$ 45 t/yr.

or total settleable solids, and the operator may be required to install settling basins or other devices to meet these limits. Most mariculture operations, however, are in open water where discharges of nutrients or BOD are less easily measured or regulated. Regulations, if any, take the form of density controls (e.g., number of operations within a given area, stocking density limits, or distance between operations) rather than pollutant removal. In Japan, standards are established by the individual prefectures to control the intensity of culture so as to prevent deterioration of the culture grounds. In Kagoshima Prefecture, for example, the hamachi culture grounds have been divided into several subareas determined by the minimum annual dissolved oxygen concentration. In Zone A, where oxygen concentrations outside the cages reach as low as $5.5 \text{ mg}\cdot\text{l}^{-1}$, stocking densities can not exceed $11 \text{ kg}\cdot\text{m}^{-3}$ and culture intensity can not exceed $40 \text{ tonnes}\cdot\text{hectare}^{-1}$. In areas with lower dissolved oxygen concentrations, stocking densities must be reduced to $7 \text{ kg}\cdot\text{m}^{-3}$ or less and culture intensity can not exceed $20 \text{ tonnes}\cdot\text{hectare}^{-1}$ (K. Fukusho, National Aquaculture Research Institute, pers. comm.).

Farm size restrictions

The Maritime Provinces of Canada have limited the density of salmon cage culture to no more than 24 cages and a maximum of 2 hectares per lease. In the United States, the state of Washington has placed a limit of 1 million pounds annual production per square nautical mile on salmon cage culture in Puget Sound. Furthermore, recommended limits have been established for the maximum amount of production in specific embayments. These limits are determined by calculation of the natural flux of nitrogen in the embayment with each tidal cycle, and then allowing no more than a 1 % increase in this flux attributable to cage culture.

Disease transmission

A required separation between culture operations is commonly employed both to reduce water quality impacts and to minimize the potential for disease transmission between culture facilities or between cultured and wild animals. The required separation of salmon net-pen culture facilities ranges from 500 m in Norway to 1 km in several Canadian provinces. British Columbia and Norway have also regulations requiring a minimum distance between net-pen facilities and major salmon spawning streams. The distance has been established at 3 km in British Columbia and Norway.

No regulation exists in any of the member country that controls the separation of net cage fish culture and shellfish culture units, except in British Columbia.

Transplantations and Importation

The importation of an exotic species or disease organism poses the greatest environmental risk of mariculture, for unlike other environmental effects, the consequences may be widespread and irreversible. Therefore, the importation of live animals or their reproductive products is subject to a greater degree of regulatory scrutiny than any other aspects of mariculture. Before an exotic species is approved for importation, most countries require an evaluation of the potential environmental effects of the introduction. This will include an assessment of the potential for interbreeding or competition with native species, the likelihood of the imported species established a self-sustaining reproducing population, and the potential for accidental introduction of pests or pathogens. If introduction of the species is approved, each individual shipment will generally require a permit from the responsible government agency. The permit will carry with it certain conditions which may include restrictions on the country of origin, visual and/or

Table 4: Environmental survey requirements for salmon cage culture in Washington state

	Site Characterization Survey	Baseline Survey	Annual Monitoring
Class I Facilities (<9 t/yr)	<ul style="list-style-type: none"> • Recommended consultation with state and local authorities • Bathymetric survey • Hydrographic survey -current velocity and direction • Diver survey 	<ul style="list-style-type: none"> • None 	<ul style="list-style-type: none"> •None
Class II Facilities (9- 45 t/yr)	<ul style="list-style-type: none"> • Recommended consultation with state and local authorities • Bathymetric survey • Hydrographic survey -Current velocity and direction • Diver Survey 	<ul style="list-style-type: none"> • None 	<ul style="list-style-type: none"> • Benthic survey - Diver survey
Class III Facilities (>45 t/yr)	<ul style="list-style-type: none"> • Recommended consultation with state and local authorities • Bathymetric survey • Hydrographic survey -current velocity and direction -Droque tracking -Vertical hydrographic profiling • Diver survey 	<ul style="list-style-type: none"> • Sediment chemistry sampling • Benthic infauna sampling 	<ul style="list-style-type: none"> • Benthic survey - Diver survey - Sediment chem - Benthic infauna • Water quality sampling • Current velocity direction

pathological inspections, periods of quarantine, or disinfection requirements. The number and type of permit restrictions vary widely from country to country. Among the most restrictive regulations are those of New Zealand. The government allows no importation of live, fresh or frozen fish because of concern over the introduction of disease. Only processed seafoods (e.g., canned goods) are allowed into the country. While there are regulatory pathways through which one may apply for special exemptions, the stringent requirements (e.g., two years holding time in a quarantined facility) generally render the importation of live animals for culture impractical.

The Study Group also refers to the ICES/EIFAC "Code of Practice on the Introduction and Transfer of non-indigenous species" and strongly recommends the strict application of the Code in member countries. It is alarming to see that despite the efforts to minimize transfers the spread of disease agents has rapidly increased.

Bacteria

There are generally no regulatory restrictions controlling the release of bacteria from mariculture facilities, other than those discussed above pertaining to the importation of bacterial pathogens. There are, however, regulations intended to protect mariculture operations from bacteria originating from other sources. Regulations of this nature are most prevalent in shellfish culture. Contamination of shellfish grounds by human bacterial and viral pathogens of faecal origin is a major concern and a common problem throughout the world. Most shellfish producing countries have adopted systems of monitoring and certification of shellfish culture grounds. Since detection and identification of all potential pathogens is difficult, standards are based upon the abundance of faecal coliform bacteria. In those culture grounds which exceed faecal coliform standards, harvest may be prohibited or the grower may be required to relay the shellfish elsewhere for depuration.

Needs for improved feeds and feeding strategies

The reduction of loading by any mechanical treatment method seems to be insufficient in mariculture. The limiting factor for primary production in the sea is in general nitrogen, and only in certain circumstances phosphorus. The nitrogen not retained in fish is mostly in soluble form and not in the settleable solids. This means that the use of sludge collecting devices in net cage farming is not beneficial. This was also observed in practical experiments (Leminen et al. 1986, Kehitysaluehasto OY, 1987).

The conclusion is that the only reasonable way to reduce the loading without restricting the production amounts or relocation is the developing of feeds and feeding.

In this context the development of feeds means in this connection lowering the nitrogen content in relation to metabolizable energy. The aim is to make the protein efficiency ratio as good as possible. This is an important new viewpoint which needs to be taken into account in research on fish nutrition.

The better nutritional knowledge means better feed quality and thus also better feed conversion efficiency. The other option to improve feed conversion coefficients and reduce pollutional loads is to avoid overfeeding (see e.g. Åsgård et al 1986). The optimization of feeding is one good way to reduce not only nitrogen loading but also the accumulation of sludge underneath farms. The most important technical problem relates to the inaccuracy of determining the actual biomass in a fish cage. Most farmers estimate the standing stock on the basis of very unreliable figures. The development of more exact techniques to identify growth rates and total biomass without disturbing the fish too much, are urgently needed. One example on the possibilities to reduce feed losses through the development of feeding technique has recently been experimentally introduced by Norway, employing a floating pellet that is released near the bottom of the net cage.

The development of adequate feeds that allow not only for a high conversion efficiency but also lower the amount of feed wasted is also an important economic issue for the industry.

Beneficial effects of mariculture

Mariculture activities can have beneficial environmental effects. Fisheries can also profit from mariculture through enhanced production. One particular contribution to natural production is known as sea ranching. In the Baltic Sea for example, more than four million salmon smolts and about one million sea trout smolts are released annually, mostly by Finnish and Swedish hatcheries. This has led to a remarkable development in professional salmon and recreational sea trout fisheries in the Baltic Sea area where most salmonid spawning runs have been destroyed by various human activities. A significant salmon or sea

trout fishery could not exist without the support of an extensive ranching programme. It can also be noted that many fish stocks, especially salmonids would have completely disappeared from their natural area without regular stocking from hatchery produced fish. Recent development in the Pacific region (Canada and Japan) and also around Iceland indicate that appropriate ongrowing and release strategies (so-called time-size release window) will improve the efficiency of this type of fishery supportive mariculture. Attention has to be payed, however, to the problems of increased fishing pressure on natural populations fished together with the enhanced stocks.

In areas of dense bivalve culture, the development of a polyculture system, which provides nutrients to the extensive shellfish farming could be considered as a possible beneficial site effect in a properly managed system combining intensive and extensive operations.

Recommendations

The *ad hoc* Study Group on the Environmental Impact of Mariculture makes the following recommendations:

Research and Development

Work should be encouraged on:

- a critical evaluation of the wider use of chemicals (e.g., disinfectants, antibiotics, algicides, antifoulants) in the developing mariculture industry. An inventory should be established to include information on the types and quantities of chemicals used, the amounts used per unit of fish and shellfish produced, and national regulations controlling their usage.
- chemicals used in mariculture to determine their effects on marine organisms, persistence in the marine environment, degradation processes and products, bio-accumulation potential, stimulation of antibiotic resistance, and public health risks.
- reducing environmental nutrient loadings through the development of feeds having better conversion efficiency, protein efficiency ratios (PER-value), and physical consistencies. Studies should also be undertaken to assess the effects of husbandry practices on feed wastage.
- the effect of mariculture on bacteria including stimulatory effects on indigenous forms (e.g. *Vibrio*) and the introduction of bacteria (e.g. *Staphylococcus*) via the feed. The potential for shellfish to serve as vectors for the transmission of diseases or as reservoirs for pathogens must be evaluated. Bacterial monitoring for public health protection needs refinement to differentiate between the various sources and types of bacteria.
- sources and levels of loadings affecting critical water quality parameters, with the aim of evaluating the effects of aquaculture on spatial and temporal variation in the availability of oxygen, hypernutrification, toxicity of ammonia and hydrogen sulphide, and particle loadings to the water column.
- waste production from emerging mariculture industries utilizing species such as halibut, turbot, cod, and sable fish.
- the effects of soluble inorganic and organic waste from mariculture on primary production and the formation of harmful algal blooms.
- the potential for enriched sediments beneath mariculture operations to serve as reservoirs for cysts of harmful algae.
- the effects of organic enrichment on the benthos, with particular emphasis on quantifying the relationship between carbon flux and consequent biological and chemical effects.

- to identify criteria for site selection to minimize alteration of benthic communities.
- to determine the "recovery time" of deteriorated cage farm sites which had been left and to develop methods to shorten this period.
- methods to deal with accumulated organic sediment including procedures for deciding when the removal of sediments is necessary, appropriate removal procedures, and environmentally-sound methods of disposal.
- effects of mariculture on indigenous fauna. This work should include: (1) effects of culture activities on species composition and abundances; (2) site selection criteria to minimize impact on critical habitats such as nursery grounds or spawning areas; and (3) potential for alteration of the native gene pool by release/escape of cultured organisms.
- effect of mariculture-related structures (e.g., cages, pens, breakwaters, artificial sills) on the hydrographic regime in coastal embayments. This work should include studies to identify the effects of different configurations and sizes of culture units on the hydrodynamics, sedimentation, and water chemistry of tidal and non-tidal areas.
- development and testing of sedimentation models which predict the fate of solid wastes from mariculture facilities, and water quality models which can be used for the determination on carrying capacity for shellfish growing and the holding capacity for fish cage culture in natural water bodies. The models should be useful to resource managers.
- sediment-water column exchange of oxygen and nutrients for input to models of nutrient and oxygen balances.

Establishment of a Working Group on

Environmental Impact of Mariculture

Considering the increasing environmental issues that are emerging with the expansion of the mariculture industry, the *ad hoc* Study Group proposes that at the 1987 Statutory Meeting the Mariculture Committee recommends to the Council the establishment of an ICES Working Group on "Environmental Impact of Mariculture". If approved by the Council, the Working Group should meet initially in the spring of 1988. The terms of reference of the Working Group should be:

- to review ongoing research projects on environmental issues related to mariculture, facilitate transfer of information in order to avoid duplication of efforts in member countries, and identify research priorities. This is necessary because of the rapid and unco-ordinated development of research programmes; the results of which often appear in internal or local reports, inaccessible to the scientific community working in this field.
- to evaluate techniques and monitoring strategies, and to prepare a technical report on appropriate monitoring protocols.
- to determine criteria for site selection which minimize environmental impacts, and to produce guidelines which could be used by member countries in resource management.
- to compile a directory of chemicals currently used in mariculture and to seek the advice of appropriate organizations on the possible environmental consequences of these chemicals, with the view to providing advice to member countries on the use of chemicals in mariculture.
- to collaborate closely with the EIFAC (European Inland Fisheries Advisory Commission) Working Party on "Fish Farm Effluents". This collaboration should be facilitated by an exchange of observers between Working Groups and, if necessary, by joint meetings.

Appendix 1: Country Reports

Information on the overall trends in mariculture development and on environmental issues related to aquaculture were available only from those ICES member countries who participated in the Ad-hoc Study Group meeting. Additional information was received by correspondence from The Netherlands and Denmark. The presented information provides a brief overview and is certainly in need of expansion, especially from ICES member countries not mentioned in this report. Aquaculture regulations existing in various ICES member countries are summarized in a separate chapter in the body of the report (see pp. 21-28).

Canada

Mariculture has only become a successful industry in Canada during the last 15 years. The physiographic and hydrographic conditions associated with each area provide unique culture conditions for the production of each species. Rapidly developing sectors of the mariculture industry are the British Columbia (B.C.) and New Brunswick (N.B.) salmonid industries, and the Prince Edward Island (P.E.I.)/Nova Scotia (N.S.) blue mussel industries (Table 5). Some idea of the rapid growth of each industry is illustrated in the following tables. Environmental problems associated with the developing Canadian industry have had little investigation. Some research is underway.

Table 5: Number of sites and tonnes(brackets) produced in Mariculture in Canada in 1986

Species	B.C.	Quebec	Newfoundland	P.E.I.	N.B.	N.S.
Salmonids*	82 (500)	1(10)	3(19)	12(9)	27(679)	26 (79)
Oysters	336 (3933)**	8(50)	0	0	0	0
	3 (<1)***					
Mussels	2 (35)	4(50)	2(14)	50 (1250)	2(3)	65(545)

* *Salmo salar*, *S. gairdneri* in Atlantic Canada; in British Columbia mainly species of the genus *Oncorhynchus*. ** *Crassostrea gigas*; *** *Ostrea edulis*; mussels = *Mytilis edulis*.

Atlantic Canada

Growth of mariculture in the Atlantic provinces has been dramatic (Table 6). While many types of culture have shown growth at least one has disappeared. Bluefin tuna culture has failed due to lack of available seed stock.

Table 6: Growth of Mariculture Production (tonnes) in Atlantic Canada

	1980	1981	1982	1983	1984	1985	1986
Salmonids	123	114	153	274	272	284	408
Mussels	36	82	174	432	876	886	1777

On the Atlantic coast the most dramatic increase in production has been in the extensive culture of mussels. The intensive culture of salmon which is perceived as having the greatest potential for environmental impacts. New Brunswick produces the largest portion of Atlantic cultured salmonids, 260 of a total of 408 tonnes

in 1986. A five year production plan has been developed for salmonid mariculture (Anonymous, Department of Fisheries and Oceans, M.S., November 1985) and guidelines have been created for the physical separation of fish farms in order to minimize the potential for the spread of pathogens and pollution from the farm sites.

The potential for increased environmental impacts from salmon farming in this area is limited by its potential for growth. Competition for suitable sites in the Bay of Fundy limits the growth of the Atlantic salmonid industry to areas where the water gets no colder than -0.7°C . Further competition with the traditional herring weir fishery and a shortfall in smolt production also constrain the industry.

The impacts due to mussel culture however can be expected to grow. In this industry seed supply appears adequate and sites are available. It appears that only market forces will determine the limits for this industry and market surpluses are not projected, even at the present rate of growth, until 1990.

Pacific Canada

- Oysters (*Crassostrea gigas*)

In 1986 the British Columbia oyster industry was comprised of 336 active tenures held by 209 growers who produced about 1050 tonnes. The 1986 values are part of a general trend and represent a 7% growth in the number of farmers and an 11% increase in the hectares cultivated relative to 1985.

In the context of obtaining the maximum benefit from the utilization of their natural resources and in order to set a fair value on the rent for usage of crown foreshore, British Columbia has developed a "Habitat Suitability Index" (HSI). Though still requiring some improvement the present HSI is useful in evaluating the suitability of coastal areas for the culture of *Crassostrea gigas*.

The HSI model used for the mariculture of the Pacific Oyster is an adaptation of a methodology utilized by the U.S. Fish and Wildlife Service to document the habitat requirements for a particular species. The model was constructed from information available in the literature. Using biophysical data the model rated the aquaculture potential of a coastal area on a relative scale. The model was subsequently tested at 10 locations over a 14 month period. The output of the model was found to be highly correlated to the growth of the oysters in the field however, density dependent effects are not part of the model's considerations.

- Salmonids

In British Columbia the very rapid growth of the salmon farming industry can be seen in the increase in the total number of applications processed, the number of leases assigned and the number of eggs taken each year to support the new industry. Because the industry is in a period of rapid expansion the growth evident in both the number of active farm sites and in the number of eggs supplied to the industry has not yet been fully expressed as biomass on the farm site or as tonnage of marketed product. As most of these mariculture products require several years of growth before they are marketed it will be several years before we see the full environmental costs or financial benefit derived from the industry's development to date.

Table 7: Growth of the Salmon Farming Industry and Interest in Farm Sites in British Columbia, Canada

Year	1983	1984	1985	1986
Total number of applications	3*	23	124	501
Number of active Farm Sites	8-10	10	35	82
Tonnes Produced	128	107	120	500**
Supply of Eggs (in millions)	2.8	9.6	25	40

*Applications for marine fish farms were not recorded prior to 1984.

**estimated.

Constraints to future development of this portion of the Canadian industry do not appear to be egg supply, or site availability. Feed and financing are the more likely bottle necks in industry's development.

Concerns over potential environmental impacts of mariculture, particularly of cage culture of salmon are topical. There have been numerous public meetings and articles in the popular literature where this has been discussed. In response to these and other concerns in October of 1986 the provincial government created a one man commission of inquiry to investigate salmon farming in B.C. Among the recommendations of the commission were the strong recommendations for more monitoring of and research into the effects of the cage culture on the environment. There has been quick response on the part of government agencies. New environmental monitoring and siting requirements are now being implemented and financing for research is expected to increase. Inventory maps of prime finfish culture areas are also being created for sections of the British Columbia coast. As many of the criteria for the siting requirements and for identification of good areas for finfish culture are dependent on good water flushing, the use of this information by the industry will help to minimize environmental impacts of finfish farming in coastal waters.

Research (Atlantic)

- There is regular annual monitoring of benthic conditions in the L'Etang Inlet Bay of Fundy where twenty salmonid sites are located (Wildish *et al.* 1986).
- A physical oceanographic study of L'Etang is planned.
- There has been initiation of a seasonal study of phytoplankton species diversity and density in areas with mariculture.
- A benthic study of the impacts of salmon mariculture in Dark Harbour has been completed.

Research (Pacific)

- A study is under way on the rate of uptake of tributyltin by oysters near marine net cages treated with tin antifoulant paints, on methods of removing tributyltin paints from netting and on depuration in Pacific salmon.
- A field study of possible transmission of sulphamerizine from salmon feed to nearby oysters is in progress.
- A computer model of the water quality response of an fjord to salmon farming is being developed.

- There is an ongoing monitoring of the water quality in Sechart Inlet, and area of rapid mariculture development in a fjord with limited water exchange over a shallow sill.
- Monitoring of bivalve feed availability in a fjord with extensive oyster culture is under way.
- A system to warn mariculturists of the occurrence of potentially harmful phytoplankton blooms has been developed.
- Monitoring of bacterial water quality for growth of oysters is routinely carried out.




Figure 3: Number of salmon farms and/or applications and proposals in British Columbia for the years 1986 to 1988 (by regions).

Denmark (K.Nielsen)

Production trends

Mariculture in Denmark is dominated by trout production. Shellfish (blue mussels and oysters) are of minor importance. Trout are raised in cages and in land-based flow-through systems. In 1985 the number of cage farms and plants on land were 20 and 7, respectively (Hoffman, 1986). After cage farming of trout has been introduced in danish coastal waters during the late 70s, the increase in mariculture production has been due mainly to the expansion of these few farms. The total trout production in marine and brackish waters increased from 100 t in 1978 to over 3,800 t in 1986 (Figure 4). The annual aquaculture production in freshwater amounts to about 22,000 to 22,400 tonnes in 1986 (Anonymous, 1986, Henrik Leth, pers. comm.).

Large trout of 0.5 to 1.0 kg wet weight are placed in net cages in early spring. During the summer and autumn most of the fish will reach a weight between 3.0 and 5.0 kg. A few cage farms only are located in relatively open sea situations, while most of the farm sites are placed in bays and fjords, which offer relative good protection against wind and wave action. The production of trout in both marine and fresh waters is limited by the environmental authorities and, in 1987, there is a one year freeze for all new applications and permits will not be granted before 1988.

The production of blue mussel on hanging long line systems was 700 to 1,000 tonnes in 1985 and 1986, but only about 25 tonnes of this production were harvested during the same time period. Experiments with extensive cultures of blue mussels have been carried out, but there is yet no industrial use of this system. In 1985 about 60,000 to 70,000 tonnes of "wild" mussels were harvested. The Pacific oyster (*Crassostrea gigas*) is grown on long lines and in cages. In 1985 and 1986 the oyster production reached about 20 tonnes. This production was mainly achieved by one farm at one site (Hoffmann, 1986, Per Sand Christensen, pers. Comm.).

Research and Monitoring

During the one year ban on all new trout cage farm licences, environmental impact studies have been implemented at existing and proposed farm sites. After completion of these studies it is intended to prepare a national plan for cage farms. Some counties in Denmark have already developed such plans. For example, in the county of Storstrøms amtskommune, such a plan has been developed on the basis of the calculated nutrient loading produced by the farms and released into the environment. The content of total nitrogen and total phosphorus is measured in fish and feeds and a budget is computed assuming a feed conversion coefficient of no more than 2.0. An overall fish production of 100 tonnes of trout would then result in a loss of 12 tonnes of nitrogen and 2 tonnes of phosphorus (Table 5). On the political level it was decided that the maximum permissible nutrient loading produced by fish farms has to be expressed as a given percentage of the total load a coastal area is receiving from land based sources (Storstrøm amtskommune, 1985).

This simple method with arbitrary limits has been adopted because of the difficulties involved in detecting biological effects in marine ecosystems originating from a single nutrient source at subcritical levels. Mathematical models have not been used due to their great uncertainty of predicting responses in complicated ecological systems. Furthermore, it is felt more relevant to calculate inputs (nutrient loads) from fish farms, when the need arises to control and reduce the total nutrient loading discharged from danish coasts into the sea.

Cage farms in Denmark are often controlled by measuring the content of N, P and organic matter in the sediment around the farms. Measurements from 7 farms in 1986 located at 5 to 15 m depth show generally no distinct increase in these parameters between May and September. These results can be explained by rapid currents, relatively low nutrient content in deposited material and fast degradation of faecal pellets. In sheltered areas, however, the growth of macroalgae seem to be stimulated and sulphur bacteria cover the sea bed around cage sites (From & Rasmussen, 1984, Storstrøm amtskommune, in preparation).

In Denmark, the monitoring programme for toxic algae has been expanded in 1987, especially in areas where blue mussels are harvested intensively. Harvesting of mussels has yet not been prohibited as a result of toxic algal blooms in any area and season. The tendency towards hyper-nutrication may in the future well promote conditions for toxic algal blooms along danish coasts and may, therefore, be considered as a future area of concern to mariculture development.

Table 8. Nutrient load from 100 tons net production assuming 200 tons fodder consumption.

	Fodder (tons)	Fish (tons)	Loss (tons)
Total nitrogen	14,9	2,9	12,0
Total phosphor	2,4	0,4	2,0




Figure 4. Production of rainbow trout in marine and brackish water.

Federal Republic of Germany (Harald Rosenthal)

Production Trends

Conventional pond farming (trout = 15,000 tonnes, carp = 8,400 tonnes, 1982) contributes the major portion of fish culture products. The number of freshwater farms has drastically increased between 1971 and 1981 (by 58%) while their average size decreased, indicating that more and more farmers share the limited water resources. Marine finfish farming is presently limited to net cage farms in Power plant effluents of the Kiel Bight (Baltic coast), to land-based hatchery operations for turbot production, to brackish-water culture of eels in heated effluents (Emden, North Sea coast) and to a few small-scale oyster farms (total production in 1986 about 2 tonnes). Substantial growth was experienced in extensive mussel farming, although the number of companies operating in the inner German Bight has not changed and will most likely not expand in the near future. In the past, annual yields of blue mussels fluctuated between 11,000 and 17,000 tonnes, and this situation remained constant for more than a decade. A major increase occurred in 1982, leading to a harvest of 27,000 tonnes, which increased further to 32,000 tonnes in 1983 and an exceptional level of 51,000 tonnes in 1984. Since then the production figures have leveled around 30,000 tonnes.

Research

Research activities related to environmental concerns are carried out mainly in freshwater situations. In general, environmental issues in mariculture are restricted to water quality within farming systems, on waste heat utilization and on oxygen budgets and plankton blooms in Baltic coastal waters .

Regulations for mariculture applications are not yet harmonized in various States. Usually, interested applicants will have to apply to the local fisheries authorities. A number of other agencies (transport, water resources, fisheries, environmental Protection Agencies, Island and Coastal Protection offices) will be involved to grant a licence. If a licence for mussel farming, for example, is issued, the farmer has to use the lease, otherwise he loses his licence immediately. There are no restrictions as to the size of the operation.

In freshwater, effluent standards have been set for land-based intensive farming systems (flow-through operations), limiting the COD and the suspended solid inputs into the receiving waters.

Finland (Timo Mäkinen, Markku Pursiainen)

Production Trends

Aquaculture has grown rapidly in Finland over the last decade. Figure 5 shows number of farms and total production during the early 1980s. Fish production of brackish water net cage farms has tripled during the 80's. At present the approximately 7,000 tonnes of rainbow trout constitutes 70 per cent of the total aquaculture yield of the country. In the past it took three growing seasons in freshwaters to produce trout of 1.5 to 2.5 kg weight. Because of the utilization of heated effluents to produce fish ready for stocking into cages, this time in freshwater is being shortened by one year and fish has to spent only the first spring in the freshwater farm (average weight about 5 g). Most of the production in brackishwater cages is derived from sites in the southwestern archipelago (between Hango peninsula SE and the city of Pori N, a 200 km long stretch of the coast with 125 fish farms, producing 6,000 tonnes annually). About 30 farms with around 2,000 tonnes annual output are located around the county of Åland. Future production increases are not likely to be as rapid (see Mäkinen & Pursiainen 1987).

Finland also produces salmon fingerlings for stocking into the Baltic. The number of smolts released annually exceeds 2.5 million individuals.




Figure 5: Development of finfish culture in Finland during the early 1980s. Column 1 = total production; column 2 = brackishwater cage culture

Research and Monitoring Activities

Monitoring aquaculture operations is often requested by the water court, and is done by the farmers themselves. The activity covers the registration of (a) the amount of feed used, (b) the total fish stock on site, and (c) the annual production (Salonen, 1983). Monitoring water quality usually does not show any obvious negative effect in the vicinity of the fish farms (Halonen 1985). In order to gain some understanding on the effect of these farms in oligotrophic fresh and coastal brackish waters several research projects were initiated, studying the immediate - and most important - the long-term biological effects around the net cage farms. Most of the research projects include all trophic levels in their investigation: from primary production to fish community structure (e.g. Jumpanen, 1986; Hallikainen, 1986; Koivisto, 1987).

Research projects aimed at reducing the pollutional load from coastal net cage farms tend to apply methods already tested on land-based fish farms (e.g. Mäkinen & Naukkarinen, 1982; Mäkinen 1984, 1985). On the basis of the Wollenweider-Dillon model on acceptable loading levels decisions are obtained whether measures to reduce the overall load from farms in a particular area have to be implemented. Because feed composition and wastage contribute substantially to the pollution load derived from fish farms, the development of appropriate feeds and efficient feeding technologies are subjects of high priority which can lead to an immediate reduction of the problem. As an example, the phosphorus content of the feed may largely influence the amount of phosphorus released into the environment (Figure 6). To some extent, the removal of sludge accumulated under cage farms has been tested as an option to minimize environmental impact. So far, these studies have provided rather poor results (Leminen *et al.*, 1986; Kehitysaluerahasto, OY., 1987).




Figure 6 : Estimated phosphorus loading per kg fish produced in relation to phosphorus content of feed and feed conversion efficiency (modified after Mäkinen, 1985).

France (Maurice Héral)

Production trends

The primary species utilized in oyster culture in France is the Japanese oyster (*Crassostrea gigas*). The annual production is usually above 100,000 metric tonnes (wet weight), reaching a maximum of 130,000 tonnes. Production of the European flat oyster (*Ostrea edulis*) remains low due to the epizootic disease outbreaks of *Martelia* and *Bonamia*. The annual production of mussels (*Mytilus edulis*, and *M. galloprovincialis*) has recently reached almost 60,000 tonnes. This has been possible mainly because of the development of long-line culture in new areas. Culture of the Manila-clam (*Ruditapes phillipinum*), is an emerging industry with a yield of 500 tonnes in 1986.

The production of rainbow trout (*Salmo gairdneri*) has reached 27,000 tonnes annually, however, 98% of this figure originates from freshwater production. In 1985, marine cage farming of rainbow trout reached 500 tonnes. Coho salmon (*Oncorhynchus kisutch*) production reached 60 tonnes during the same year. Production is limited principally by the difficulties of cultivating this species in the warm summer water. The cultivation of sea bream (*Sparus aurata*) and sea bass (*Dicentrarchus labrax*) has passed the experimental phase and has recently entered commercial production (100 tonnes). The culture of shrimps (*Penaeus japonicus*) has also recently started on a commercial scale and reached an annual output of 50 tonnes.

Determination of the environmental impact of mariculture has focussed mainly on studies related to molluscan aquaculture (e.g. oysters, mussels, clams). The following aspects have been investigated:

- Development of a dynamic model for oyster carrying capacity based on historical data. The results so far show that the bay studied has a production limit which remains constant even if the total biomass of the stock increases.
- A trophic approach to determine oyster carrying capacity of a restricted area (bay, embayment), combining a physical model of the flux of particulate organic matter to show the distribution of the available food. Organic matter is consumed and transferred in the production of the molluscs in accordance with energetic laws. This model will be useful in the future to predict the impact of the various inputs (nutrients, sedimentation, currents) and/or environmental alterations on molluscan culture.
- A model of nitrogen and phosphorus fluxes under oyster and mussel culture units.
- A model of biodeposition from molluscs as a function of seston and particulate organic matter.

No research programmes are in progress, as related to environmental issues in marine fin fish culture. However, studies on the interaction between molluscan culture and intensive fish farming have shown that bacterial contamination of oysters can occur in the vicinity of fish culture. For example, after the occurrence of a furunculosis outbreak in rainbow trout (land-based tank culture), *Aeromonas* spp. have been found in cultivated oysters which were grown in the effluent of the fish farm. The problems of accumulating antibiotics and of the transfer of bacterial resistance to organisms which are pathogenic to human have been discussed.

Tributyltin (TBT), used as an antifoulant on pleasure boats, has been shown to accumulate in fish and can cause difficulties for adjacent oyster culture in the form of TBT accumulation in bivalve tissues, reduced shell growth and spat mortalities

Problems of the carrying capacity of a bay for mollusc culture

Molluscan food is mainly composed of natural phytoplankton and particulate organic material. For that reason the biomass of cultivated molluscs can surpass the carrying capacity of the bay for optimum growth causing not only a significant decrease of the growth rate (Figure 7) but also high mortality rates in the stocks. Such events can occur particularly in shallow bays where the residence time of the water body is high so that the quantity of natural food is rapidly depleted through effective filter feeding of the molluscs, limiting their productions. This relationship has been established for the Bay of Marennes-Oleion which is the main European center of oyster production with a mean annual production of 40,000 tonnes (total wet weight). An historical approach has been realised for the cupped oysters, Portuguese and Japanese species. The first sign of overcrowding in an area is the decrease of the growth rate of the population which occurred already for both species.




Figure 7: Changes in growth rates of the Japanese oyster *Crassostrea gigas* in the Bay of Mrennes-Oleian, an area of increased aquaculture production, during the period 1972 to 1986.

It has been shown that the changes in growth rates over the years were not caused by internal biological factors, but by overcrowding. The survey also showed that growth depression was more the result of lack of nutrition which expressed itself in physiological mortalities during the winter and after the spawning period in summer.

The relation between the total biomass of the stock (B) and the production (P) shows that it is following the Von Bertalanffy function, where P_{∞} corresponds with the limit of the carrying capacity of the bay for this species. It has been demonstrated that the climate does not influence this scheme but induces some variability. The evolution of the annual turnover ratio (P/B) of the stock exhibits a negative exponential curve.




Figure 8: Relationship between oyster production and total stock biomass of the cupped oyster in the Bay of Marennes -Oleion




Figure 9: Principle decrease of the oyster production rate per unit biomass with increasing overcrowding (total stock biomass).

The economic benefits of oyster culture in the bay can be considered as a function of the curve in Figure 9. All the arguments show the necessity to reduce the total biomass of the stock in order to increase the efficiency of the aquaculture production on extensive molluscan farming grounds. The relationship is now used practically to regulate stock size and to maximize the production and the overall economic benefits of this operation.

Norway (Arne Ervik, Jan Aure)

Production trends

In Norway salmonid mariculture has developed into a significant industry. Its annual production reached about 50,000 tonnes in 1986, and a substantial further growth is predicted with a possible output of about 80,000 tonnes in 1989 (Table 9).

Table 9: Recent production figures in Norwegian Mariculture and the projected overall production for the year 1989.

Species	Production 1986 (tonnes)	Production 1989 (tonnes, predicted)
Rainbow trout	4 300	6 000
Atlantic Salmon	45 600	74 000
Arctic Char	2	2 - 50
Blue Mussel	200	1 800
Oysters (numbers)	0.1 million	5.0 million

New species are expected to enter the production in marine aquaculture in the near future. For the first time cod fry will be produced on a commercial scale in 1987 (2.0 - 3.5 million specimens), and results in experimental halibut culture are sufficiently encouraging to expect soon a commercially feasible operation. Other species of interest include turbot (a production potential of 12,000 to 20,000 tonnes has been estimated), arctic char, wolffish, mussels, scallops and oysters.

Figure 10 shows the number of of the salmon cage farms and their regional distribution in Norway in 1986. The majority of the farms can be found along the western and northern coasts. These regions are sparsely populated and anthropogenic nutrient inputs are insignificant.

Research and Monitoring Activities

Research:

Research on environmental aspects of mariculture has only recently been established. The first projects were partly descriptive and aimed at an overview on water quality variations in cage fish farms and on the effects of these farms on surrounding areas.

The present research activities are more specific, often experimental and aim at specific targets such as:

- **Optimisation of environmental conditions for fish farming.** These investigations include the following two aspects: (a) how does intensive mariculture influence water quality within the farm and in the near vicinity of the farms, and (b) how does the resulting water quality influence the farmed fish? Studies on water exchange rate, the importance of water depth, the effects of stocking density, overfeeding, and feed quality are being conducted. Further, the growth of fish, their health, and the quality of the final product are other factors that are considered. Finally, measures to control environmental factors are studied.

- **Influence of mariculture on the receiving water.** These studies include investigations on the transformation of organic compounds originating from fish farms and their environmental effects. Of central importance are flux rates between sediment and water phase, the pathways by which organic material is transformed, the trophic relations altered by the additional inputs of organics, the determination of the holding capacity of an area (e.g. local fjord), and on the development of models for calculating the holding capacity.


- **Effect of antibiotics on the environment.** Studies concentrate on possible pathways and their alteration, on resistant microorganisms, and on traces of antibiotics in free living consumer organisms.

- **Method for treating maricultural wastes.** Various methods to minimize waste output on farms are presently evaluated.

Monitoring

Monitoring of water quality is not compulsory and has been imposed on only a few sensitive and contentious areas. Some fish farmers voluntarily monitor their sites with methods available to them. Hydrographic conditions in basins with largely stagnant deep water masses, have been monitored. The studies included water quality, the composition of the sediments, and the macro-benthos community.

Figure 10: Regional distribution of salmon cage farm licences in Norway in 1986.



Portugal (M. A. Sampayo)

Production trends

Fish and shellfish cultivation in Portugal has a long history and includes mainly extensive culture methods. Marine fishes have been cultivated mainly in connection with salt production at the water entrance ponds serving salinas, in estuariene ponds and in coastal lagoons. Species cultured include *Sparus aurata*, *Dicentrarchus labrax*, *Solea* spec. and *Anguilla*. Since 1975, however, there are a few additional places that produce marine fish only and this type of aquaculture has recently become a growing industry. Intensive fish culture systems are presently employed only in freshwater aquaculture, using rainbow trout and eel as the prime species.

In shellfish production the species used are *Ruditapes decussata* (clam), *Crassostrea angulata* (oyster) and recently the culture of *Penaeus japonicus* (shrimp) has been introduced.

The 1986 estimates on the overall aquaculture production in Portugal amounts to the following figures:

Polyculture (Mariculture)	1,000 tonnes
Clams	7,000 tonnes
Other Bivalves	1,000 tonnes
Eels	100 tonnes
Rainbow trout	1,865 tonnes

Research and monitoring activities

Research on environmental aspects of mariculture have, in the past, been sporadic and have focussed mainly on water quality variations in fish ponds, shellfish beds and supporting estuaries and coastal lagoons.

The present research activities consider more specific aspects, include often an experimental part and aim at very defined key issues:

- Optimization of environmental conditions for shrimp farming, including nutrient evaluation, benthos studies and plankton growth.
- Identifying conditions in fish ponds and studying the influence of the intake water on system performance. Consideration is given to water exchange rates and their importance on environmental conditions in the vicinity of farm sites.
- Monitoring shellfish beds from a sanitary viewpoint.

The awareness among scientists, that a growing mariculture industry may face environmental problems, is increasing, calling for more detailed studies to identify the extent of the problem and to seek for solutions. There seems to exist circumstantial evidence, that a recent unusual localized plankton bloom (red tide) may have occurred in succession of repeated releases of nutrient rich waters from nearshore intensive pond culture systems (near Alvor Ria, Lagos Bay, Algarve coast, in 1983) and that this event was favoured at the time by calm weather conditions.

Since most of the present and future intensive fish farms in Portugal will operate as land-based systems, regulations on possible effluent treatment requirements for new farms and/or effluent standards are presently discussed.

Sweden (Hans Ackefors)

Production trends

The cultivation of fish and shellfish for the market has been very slow to develop, in part because of strict regulations based primarily on environmental considerations. Rainbow trout, salmon, arctic char, carp, eel and blue mussel are farmed for the commercial market. The 1985 production was as follow:

	Marine Production (tonnes)	Total Production (tonnes)
Rainbow trout	1,772	2,532
Salmon	81	81
Arctic char		5
Carp		1
Eel		47
Blue mussel	415	415
Total	2,268	3,081

The present production of blue mussels is about 3,000 tonnes. However, the harvesting of blue mussel has been banned many times during the two last years due to the high concentration of toxic substances (DSP and PSP) induced by algal blooms. The mussel farmers are using the long line method. Vertical polypropylene strips are suspended in the water from horisontal ropes. The normal growing season lasts 18 months.

The present annual production of rainbow trout is in the order of 4,000 tonnes. About 70% of the production originates from brackish or marine waters.

Sweden currently has an extensive hatchery programme for the conservation of its fishery. Between two and three million smolts of salmon and sea trout are cultivated annually and stocked in Swedish rivers. This is necessay because the natural spawning areas were destroyed when the hydropower plants were built. In addition, a number of other salmonid species are raised in hatcheries to enhance sportfisheries. Most of the marine fish production takes place in net cages and only a minor part is produced on land in flow-through systems. The biggest eel plant with an annual out-put of 100 tonnes per year uses a land-based technique. The eel facility employs a brackish water flow-through system and utilizes waste heat water from three sulphuric acid plants (Ackefors et al., 1986). Only one commercial eel plant is using a recirculating system.

Research/monitoring activities

Swedish authorities and scientists have been concerned with the loading of nutrients and organic matter from cage culture. In addition to that chemical therapy and antibiotics in feeds have been critical issues.

The topic of primary concern was originally phosphorus input, which can increase the primary production, and the input of organic matter which can cause depletion of oxygen. In recent years, the research emphasis has shifted to both phosphorus and nitrogen loading, and the effect of these substances in freshwater and marine waters.

Enell (in preparation) has investigated the variation of nitrogen and phosphorus in various feed types. He found that the phosphorus content varied from 14 to 20 mg P/g TS, while the nitrogen content varied from 55 to 100 mg N/g TS. The average content of phosphorus in the feed was 1.62 % of the dry matter and the content of nitrogen was 8.45 %. Wide variation in nitrogen and phosphorus was found not only between feeds provided by different manufacturers, but among batches from the same supplier. Enell (in preparation) also investigated the contents of phosphorus and nitrogen in the rainbow trout (the harvest). Enell found that the average values were 1.07 % for phosphorus and 8.06 % for nitrogen on a dry weight basis, although there was substantial variation during the year.

The amount of sedimentation under freshwater cage culture operations was investigated over a three year period in an eutrophic lake (Byasjön) and an oligotrophic lake (Skårsjön). The annual production of fish at each site was in the order of 20 to 40 tonnes. The amounts of sedimentation per day and m² were 20-40 g measured as dry substance and 0.4-0.6 g measured as phosphorus. At the reference stations the values were much lower although the sedimentation measured as dry substance was about 1/3 in the eutrophic lake. The redox potential in the sediment was only slightly lower relative to the reference station under the net cage in the eutrophic lake, while in the oligotrophic lake, the sediments had a far lower redox potential relative to the reference stations.

Enell & Löf (1983) investigated the phosphorus and nitrogen flow in an open net cage operation of rainbow trout. Ackefors & Södergren (1985) used their values to show a scenario with a nitrogen and phosphorus budget for a production unit of 50 tonnes rainbow trout. The calculated values were as follows.

A. Phosphorus

1. Production	50,000 kg
2. Feed coefficient (1.6)	
3. Feed amount	80,000 kg
4. Phosphorus in fodder (1.1 %)	880 kg
5. Phosphorus in the harvest (0.4 %)	200 kg (23 %)
6. Phosphorus dissolved in the water	97 kg (11%)
7. Phosphorus particulated (sediment)	580 kg (66 %)

B. Nitrogen

1. Production	50,000 kg
2. Feed coefficient (1.6)	
3. Feed amount	80,000 kg
4. Nitrogen in fodder (7.2 %)	5,760 kg
5. Nitrogen in the harvest (2.9 %)	1,450 kg (25 %)
6. Nitrogen dissolved in the water	3,570 kg (62 %)
7. Nitrogen bound to particles	780 kg (13 %)

These figures indicate that most of the phosphorus (66 %) in the feed accumulated in the bottom mud, while only 11 % was dissolved in the water. Only 13 % of the nitrogen accumulated in the bottom while 62 % occurred as dissolved nitrogen in the water.

A device to collect the feed wastes and faeces under the net cages was constructed by a Swedish company (Enell & Löf, 1984). Funnelshaped PVC-cloth was attached under the net cages and the waste was siphoned to a tank. The initial results were good. However, in practice this has procedure has not been a success.

The water quality around a cage fish farm in the Bothnian Sea was investigated by Müller-Haeckel (1986). Criteria studied were dissolved oxygen, nitrate, phosphate, particulate organic matter and the annual cycle of planktonic algae abundance. None of the investigated parameters showed significant differences between farm site and reference station.

From marine cage culture, Hall and Holby (1985, 1986) presented the first results from the Swedish west coast (Gullmar Fjord area, Bohuslän), where rainbow trout are raised in cages (15 cages, 25x35 m total area coverage) between April and December/January. The model farm produces a maximum biomass of about 40 tonnes of trout annually and cages are placed in waters of 18 to 21 m depth. Bottom sediments under the farm are always below the halocline (surface water 20-25 % S, bottom water 30-35 % S). The findings on sedimentation rate, benthic fluxes of nutrients, sediment oxygen uptake, gas ebullition and chemical composition of sediments and pore waters indicate that the top 12-15 cm of the sediments under the farm were strongly influenced by the farm, resulting into high sulphide concentrations in pore water and a high flux of sulphate, drastic enrichment of nutrients in the top layer (i.e. phosphate concentration up to 400 μM). Under the farm the organic carbon load was about 10 times the concentration in sediments near the farm, but 20 times higher than from control stations. Similar figures were found for nitrogen.

Studies on sedimentation rates using sediment traps showed that flux values ranged from 50 to 200 TPM (Total particulate matter; g/m^2 /day, dry weight) during the peak growing period (May - October), with an average value of about 115 TPM (control values for the fjord: 4-5 TPM). Phosphate was rapidly released from the sediment, being 400 times higher under the farm than 50 m outside the farm and still 40 times above values for the control station. In this study gas ebullition was measured for the first time under marine cage farms. During late summer and early fall gas ebullition rates averaged 7.9 L/m /day in September (Depth 19 m = 23 L/m/day at 1 atmosphere pressure). The gas consisted entirely of methane.

It is assumed that further studies in Sweden will provide site specific data on nutrient loads and on biological criteria identifying the areal extent of the impact.

Research has been done to find out the best site selection for aquaculture. One approach to this problem in a non-tidal area was taken by Håkansson (1985). This author constructed a model based on two key environmental factors: (1) the water retention time and (2) the bottom dynamic conditions. Together with co-workers (Håkansson et al., 1985) developed this method further to predict in a simple way a first approximation on the best location of aquacultural net cage operations.

United Kingdom (Richard Gowen)

Production Trends

The finfish industry in the United Kingdom was relatively small, with a first sale value of approximately L 44.4 million in 1984. At present this industry is expanding rapidly. Rainbow trout production has grown from 1,000 tonnes to over 12,000 tonnes in the last 10 years. Atlantic salmon output has risen from about 20 tonnes in 1974 to over 4,000 tonnes in 1984. In 1986 the production of salmon was 12,000t. The production of salmon in marine waters takes place almost wholly in Scotland. By 1990 the annual production is expected to exceed 20,000 tonnes.

Other species farmed include eels, turbot, and carp, with Dover sole on an experimental basis. The past production trends are outlined in Table 10. Shellfish production is also listed in Table 10.

Table 10: Production trends for fin fish farming in the United Kingdom.

Tonnes (live weight)

Year	1974	1980	1981	1982	1984	1986	1987
Atlantic Salmon	20	598	1,133	2,150	4,000	10,000	16,000
Rainbow Trout	1,100	5,707	7,001	6,700	12,500	12,000	12,000
Eels						50	50
Turbot						100	150
Dover Sole						20	20

Shellfish (1986 estimates): mussels= 500t, oysters= 300t, Scallops= 50t.

Current Research in Scotland

At present there are two projects aimed at assessing aspects of the ecological impact of fish farming.

1) Stirling University (Dept. of Biological Science) is conducting a study to assess the effect of fishfarming on the benthos and water column of Scottish fiordic sea lochs. The aim of this work is to develop guidelines or a model which would allow the potential impact of a fish farm (shell or finfish) in any location to be assessed (Gowen, *et al.* 1985; Gowen & Bradbury, 1987; Brown *et al.*, 1987).

2) The Department of Agriculture and Fisheries for Scotland (DAFS) has recently started a study to assess the level of hypernutrification resulting from salmon farming.

DAFS has undertaken studies on the effects of Tributyl tin and are about to undertake toxicity trials to evaluate the toxicity of Nuvan (used in the treatment of sea-lice).

United States (Don Weston)

Production trends

With the exception of extensive bivalve culture, commercial mariculture in the United States has been largely limited to a small number of salmon farms along the northeast and northwest coasts, and a few shrimp culture facilities on the Gulf Coast. Production trends are presented in Table 11. Private mariculture produces over 40% of the oyster yield, most of the catfish and crawfish, nearly all of the rainbow trout, and small quantities of several other species.

Research and monitoring activities

Little attention has been given to the environmental effects of mariculture. A great deal of work was done in the 1970s on water quality effects of freshwater culture (e.g., Hinshaw, 1973; Liao, 1970; Willoughby, et al., 1972), but comparable work in mariculture has been generally lacking, or confined to the gray literature. Odum (1974) represents one of the first attempts to review available information on the effects of mariculture, but scope of this review was seriously limited by the lack of data.

Within the past few years, a number of investigators have begun to consider the environmental consequences of continued industry growth. Some examples include:

- 1) The use of tributyltin has attracted much recent attention in the United States as it has around the world. Staff of the National Marine Fisheries Service have specifically studied the use of the compound as an antifoulant on salmon cages. They have found that the substance can be toxic to salmon and was responsible for the mortality of fish at one mariculture facility (Short and Thrower, 1986a). Residues of TBT were found in the flesh of market cage-reared salmon, thus providing a route of entry into the human diet (Short and Thrower, 1986b).
- 2) Much work has been done in the United States on diseases of cultured marine organisms (e.g., reviews of Fryer and Rohovec, 1984 and Sinderman, 1984). While most of this work has been directed towards improving the health of the cultured organisms, either by improved disease diagnosis or treatment, there has been some recognition of the environmental consequences of disease in mariculture (Sinderman, 1977).
- 3) Cage culture of salmon shows potential for rapid growth in the states of Maine and Washington. In Washington in particular, considerable interest has been shown on the environmental effects of cage culture including effects on the sediments and the benthic infauna, water quality, and native salmon stocks. This interest has spawned several studies at individual farms (Milner-Rensel Assoc., 1986; Pease, 1977) and a review of the literature on environmental effects of cage culture and suspended shellfish culture (Weston, 1986).
- 4) The work of Stanley Katz and associates at Rutgers University represents one of the few on-going efforts to examine the environmental effects of antibiotic usage in both in agriculture and aquaculture. This group has undertaken studies of antibiotic persistence in soils (Gavalchin, 1983) and surface waters (Schomburg-Barrett, 1982), and evaluated the potential for stimulation of antibiotic resistance in bacteria (Katz, unpubl.; Brady, et al., in prep.).

Table 11: Aquaculture Production Trends for the most important species in the United States. (compiled from various sources)

Species	1980	1982	1983	1984
Oysters	10,775	9,878	10,540	11,158
Crawfish	10,849	25,000	27,300	27,000
Channel Catfish	34,855	90,909	100,000	109,000
Pacific Salmon	3,455	11,587	9,400	16,644
Trout	21,836	21,818	22,000	22,700

Sowjet Union (L.A. Dushkina)

Production

Besides ocean ranching programmes for salmonids, mariculture production along the coasts of the USSR is relatively small when compared to freshwater production.

Research activities

In order to realize cage rearing of salmonids in the coastal zone of the White Sea, the Polar Research Institute undertook studies to implement this method. Trout for stocking the cages were received from a farm near the Imanda Lake, which uses heated water for rearing. The optimum weight of fish for stocking was determined and the effects of various stocking densities were evaluated in relation to various feeds. The results of the studies showed that 1-year-old rainbow trout of 100-140 g weight were of the best suited size class to be planted into cages for the summer growth in the White Sea. A cultivation period of 120 days during the summer yielded marketable fish weighing between 350 and 400g. Feed conversion efficiency reached a factor of 2.2. The use of automatic feeders decreased the conversion factor to 1.6-1.7. The best stocking density for cages measuring 3 x 4 x 2 m was determined with 10 kg m⁻², resulting a final density at harvest of 30 to 35 kg m⁻². The growth rate of trout fed an experimental diet was higher (3.2-3.6 g/day) than that of trout feeding on commercial standard diets (3.0-3.1 g/day).

In 1986, research was also conducted to determine the acceleration of growth of young Atlantic salmon in order to obtain early smolts which can be stocked into sea water. The cultivation of juveniles at 12 to 14^o C resulted into a weight increase of the 1-year old fish in the order of 5 to 10 g in comparison to 0.8 to 1.0 g in fish raised in non-heated water. Studies of the chloride cells in the gill epithelium showed that raising in heated water had no negativ effect on smoltification but enhanced the process by a factor of 2 to 3.

The effectiveness of artificial reefs were further studied in the Baltic, Azov, and Black Sea. Experimental and industrial trials proved that these reefs were effective in providing protection for juvenile fish, for reproduction of some hydrobionts and for elevating the biological productivity in coastal areas. Reef communities were studied in the Azov and Black Sea, especially the development of biocenotic patterns.

The biotechnology of growing mussels in the White and Black Sea was further developed, reaching densities of 50 to 60 tons per ha. The cultivation of mussels to marketable size takes 4 years in the White Sea and 1.5 year in the Black Sea.

The impact of mussels farms on adjacent environments has also been studied. It has been shown that the composition of sediments, bottom species composition and the biomass of the benthos changed in the vicinity of mussel farms.

Appendix 2:

Meeting of the
ICES Ad hoc Study Group on
"Environmental Impacts of
Mariculture"

April 7 to 9 , 1987
in Hamburg

at the

Biologische Anstalt Helgoland
Zentrale
Notkestrasse 31
2000 Hamburg 52
Federal Republic of Germany

Tentative Agenda

Tuesday, April 7

- 09:00 - 12:30 - Opening of the Meeting
- Adoption of the Agenda
- Election of Rapporteur
- Country by country status reports
- 12:30 - 14:00 - Lunch
- 14:00 - 18:00 - Discussion of a conceptional outline
for the Study Group Report
- Drafting Session: country reports
- 18:00 - adjourn

Wednesday, April 8

- 09:00 - 12:30 Polluttional aspects in coastal farming
- extent of the problem (regional considerations)
 - Quality of fish farm effluents
 - land-based systems
 - water-based systems
 - estimates on polluttional loads (nutrients) based on physiological (metabolic) data
 - nutritional aspects and waste load
 - treatment options (methods to minimize loads)
- 12:30 - 14:00 - lunch
- 14:00 - 18:00 - drafting session
- 18:00 - adjourn

Thursday, April 9

- 09:00 - 12:30 - Management aspects
- Environmental Impact Statement
 - Estimating carrying capacities of coastal areas
 - Modelling approaches
 - Habitat Suitibility Index
 - Aquaculture and Eutrophication
- Discussion of Conclusions and Recommendations
- 12:30 - 14:00 - Lunch
- 14:00 - 18:00 - Drafting session:
Conclusions and Recommendations
- 18:00 - adjourn

Appendix 3: Executive Summary

Following the adopted Resolution CR 1985/2:37 (73rd Statutory Meeting of ICES), CR 1986/2:36 (74th Statutory Meeting) an *ad hoc* Study Group on the "Environmental Impacts of Mariculture" was established in order to: (a) delineate the dimensions of the problem, and (b) recommend a course of action which will lead to a development of criteria and to a standard system of monitoring and reporting." Membership to this *ad hoc* ICES Study Group was assigned during 1986 with experts nominated from 11 member countries. The Study Group worked by correspondence and met in Hamburg (Federal Republic of Germany) between April 7 and 9, 1987 to prepare the report. Country reports on present production figures, anticipated yields and effective regulations were received from Canada, Denmark, Federal Republic of Germany, Finland, France, Norway, Portugal, Sweden, United Kingdom, and the United States. The environmental issues addressed include (a) our present understanding of the effects of mariculture on natural microbial communities and on the possible spread of pathogens, (b) the possible changes in the natural populations of marine algae (phytoplankton and macroalgae), (c) the influence of sedimentation on the benthos, (d) the use of chemicals in mariculture, (e) site selection criteria to minimize environmental effects, and (f) the state of the art in developing predictive models of mariculture impacts. Further chapters discuss environmental regulations (as applied to mariculture situations) in various member countries, the need for improved feeds and feeding strategies, and the possible beneficial effects of mariculture. Finally, a number of recommendations were formulated, as they relate to research needs and to the assessment and minimization of environmental impacts:

Recommendations:

Research and Development

Work should be encouraged on:

- the wider use of chemicals (e.g., disinfectants, antibiotics, algicides, antifoulants) in the developing mariculture industry. An inventory should be established to include information on the types and quantities of chemicals used, the amounts used per unit of fish and shellfish produced, and national regulations controlling their usage.
- chemicals used in mariculture to determine their effects on marine organisms, persistence in the marine environment, degradation processes and products, bioaccumulation potential, stimulation of antibiotic resistance, and public health risks.
- reducing environmental nutrient loadings through the development of feeds having better conversion efficiency, protein efficiency ratios (PER-value), and physical consistencies. Studies should also be undertaken to assess the effects of husbandry practices on feed wastage.
- the effect of mariculture on bacteria including stimulatory effects on indigenous forms (e.g. *Vibrio*) and the introduction of bacteria (e.g. *Staphylococcus*) via the feed. The potential for shellfish to serve as vectors for the transmission of diseases or as reservoirs for pathogens must be evaluated. Bacterial monitoring for public health protection needs refinement to differentiate between the various sources and types of bacteria.
- sources and levels of loadings affecting critical water quality parameters, with the aim of evaluating the effects of aquaculture on spatial and temporal variation in the availability of oxygen, hypereutrophication, toxicity of ammonia and hydrogen sulphide, and particle loadings to the water column.

- waste production from emerging mariculture industries utilizing species such as turbot, cod, and sable fish.
- the effects of soluble inorganic and organic waste from mariculture on primary production and the formation of harmful algal blooms.
- the potential for enriched sediments beneath mariculture operations to serve as reservoirs for cysts of harmful algae.
- the effects of organic enrichment on the benthos, with particular emphasis on quantifying the relationship between carbon flux and consequent biological and chemical effects.
- to identify criteria for site selection to minimize alteration of benthic communities.
- to determine the "recovery time" of deteriorated cage farm sites which had been left and to develop methods to shorten this period.
- methods to deal with accumulated organic sediment including procedures for deciding when the removal of sediments is necessary, appropriate removal procedures, and environmentally- sound methods of disposal.
- effects of mariculture on indigenous fauna. This work should include: (1) effects of culture activities on species composition and abundances; (2) site selection criteria to minimize impact on critical habitats such as nursery grounds or spawning areas; and (3) potential for alteration of the native gene pool by release/escape of cultured organisms.
- effect of mariculture-related structures (e.g., cages, pens, breakwaters, artificial sills) on the hydrographic regime in coastal embayments. This work should include studies to identify the effects of different configurations and sizes of culture units on the hydrodynamics, sedimentation, and water chemistry of tidal and non-tidal areas.
- development and testing of sedimentation models which predict the fate of solid wastes from mariculture facilities, and water quality models which can be used for the determination on carrying capacity for shellfish growing and the holding capacity for fish cage culture in natural water bodies. The models should be useful to resource managers.
- sediment-water column exchange of oxygen and nutrients for input to models of nutrient and oxygen balances.

Establishment of a Working Group on

Environmental Impact of Mariculture

Considering the increasing environmental issues that are emerging with the expansion of the mariculture industry, the *ad hoc* Study Group proposes that at the 1987 Statutory Meeting the Mariculture Committee recommends to the Council the establishment of an ICES Working Group on "Environmental Impact of Mariculture". If approved by the Council, the Working Group should meet initially in the spring of 1988. The terms of reference of the Working Group should be:

- to review ongoing research projects on environmental issues related to mariculture, facilitate transfer of information in order to avoid duplication of efforts in member countries, and identify research priorities. This is necessary because of the rapid and unco-ordinated development of research programmes; the results of which often appear in internal or local reports, inaccessible to the scientific community working in this field.

- to evaluate techniques and monitoring strategies, and to prepare a technical report on appropriate monitoring protocols.
- to determine criteria for site selection which minimize environmental impacts, and to produce guidelines which could be used by member countries in resource management.
- to compile a directory of chemicals currently used in mariculture and to seek the advice of appropriate organizations on the possible environmental consequences of these chemicals, with the view to providing advice to member countries on the use of chemicals in mariculture.
- to collaborate closely with the EIFAC (European Inland Fisheries Advisory Commission) Working Party on "Fish Farm Effluents". This collaboration should be facilitated by an exchange of observers between Working Groups and, if necessary, by joint meetings.

Appendix 4: Literature consulted

- Ackefors, H. 1986. The impact on the environment by cage farming in open water. *J. Aquaculture Trop.* 1: 25-33.
- Ackefors, H. 1983. Development of Aquaculture in Sweden. A collection of the summaries from the various reports made by the Swedish Steering Committee on Aquaculture and its Working Groups. Council for the Planning and Coordination of Research (FRN) Report No. 83:8.
- Ackefors, H., Murray, K., Rosenthal, H. 1984. A European view on the Asian approach to modern aquaculture development. *ICES C.M. 1984/F:3*, 1-38. (Mariculture Committee).
- Ackefors, H., Rosen, C.-G. 1979. Farming aquatic animals. The emergence of a world-wide industry with profound ecological consequences. *Ambio* 8(4): 132-143.
- Ackefors, H., Södergren, A. 1985. Swedish experience of the impact of Aquaculture on the Environment. *ICES/C.M. E:40/Sess.W: 1-7*. (Mariculture Committee)
- Adlsukresno, S. 1982. Criteria for the selection of suitable site for coastal fishfarms. In: Report of consultation/seminar on coastal fishpond engineering; 4-12 Aug. 1982, Surabaya, Indonesia. FAO/UNDP Manila (Philippines), pp. 67-71.
- Alabaster, J.S. 1982. Survey of fish-farm effluents in some EIFAC countries. *EIFAC Tech. Paper* 41: 5-20.
- Allen, D.A., Austin, B., Colwell, R.R. 1983. Numerical taxonomy of bacterial isolates associated with a freshwater fishery. *J. Gen. Microbiol.* 29(7): 2043-2062.
- Allendorf, F.W., Espeland, D.M., Scow, D.T., Phelps, S. 1980. Coexistence of native and introduced rainbow trout in the Kootenai River drainage. *Proc. Montana Acad. Sci.* 39: 28-36.
- Alzieu, C. 1981. Evaluation des risques dus à l'emploi des peintures antisalissures dans les zones conchylicoles. Anomalies de calcification. Thème Surveillance continue. Mécanismes d'actions des polluants. Addendum du rapport du 15 Juin 1981. Institut Scientifique et Technique des Pêches Maritimes, Nantes, le 15 Décembre 1981.
- Alzieu, C., Héral, M. 1984. Ecotoxicological effects of organotin compounds on oyster culture. *Ecotoxicological Testing for the Marine Environment Vol 2*: 187-196.
- Alzieu, C., Portmann, J.E. 1984. The effect of tributyl tin on the culture of *C. gigas* and other species. *Proc. Annual Shellfish Conf.* 15:1-15. (Pre-print).
- Alzieu, C., Thibaud, Y., Héral, M., Boutler, B. 1980. Evaluation des risques dus à l'emploi des peintures anti-salissures dans les zones conchylicoles. *Rev. Trav. Inst. Peche Marit.* 44(4): 301-348.
- Alzieu, C., Héral, M., Thibaud, Y., Dardignac, M.-J., Feuillet, Y. 1981. Influence des peintures antisalissures à base d'organostanniques sur la calcification de la coquille de l'huître *Crassostrea gigas*. *Rev. Trav. Inst. Peche marit.* 45(2): 101-116.
- Alzieu, C. 1986. TBT detrimental effects on oyster culture in France - Evolution since antifouling paint regulation. *Oceans'86 Conference Record, Organotin Symposium, Vol.4*: 1130-1134.
- Anderson, C.D., Dalley, R. 1986. Use of organotins in antifouling paints. *Oceans'86 Conference Record, Organotin Symposium, Vol.4*: 1108-1113.
- Anderson, D.M., Wall, D. 1978. The potential importance of benthic cysts of *Gonyaulax tanarensis* and *Gonyaulax excavata* in initiating toxic dinoflagellate blooms. *J. Phycol.* 14:224-235.
- Andren, L.E. 1975. Pollution and degradation of environment affecting aquaculture in Africa. *FAO/CIFA/75/SR* 12: 1-16.
- Anonymous, 1972. Recycled waste powers shellfish project. *Quick Frozen Foods* 35 (2): 64-65.

- Anonymous**, 1979. Operation clean-up. *Fish Farming Int.* 6(2): 32-34.
- Anonymous**, 1982. Får jag lov? Vattenbrukets juridik. (May I have permission? Legal concerns in aquaculture). Swedish Council for Planning and Coordination of Research (FRN)., 73 pp. [In Swedish].
- Anonymous**, 1983. Water quality surveyed in Clam Bay. Quaterly Report, Jan.-March 1983, pp.1-12. Northwest and Alaska Fisheries Center, National Marine Fisheries Service, National and Oceanic Atmospheric Administration, Seattle, Washington.
- Anonymous**, 1983. The environmental impact of aquaculture. Report No 83(5): 1-74. Swedish Council for Planning and Coordination of Research (FRN).
- Anonymous**, 1985. Akvakultur - Miljöproblem. *Nordforsk* 1985(2): 290 pp. [In Swedish].
- Anonymous**, 1986. Udviklingsprogram for Dansk Akvakultur. Fiskeriministeriet 1986, 201 pp.
- Aoki, T., Jo, Y., Egusa, S.** 1980. Frequent occurrence of drug resistant bacteria in Ayu (*Plecoglossus altivelis*) culture. *Fish. Path.* 15(1): 1-6.
- Arakawa, K.Y., Kusuki, Y., Kamigaki, M.** 1971. Studies on biodeposition in oyster beds. I. Economic density for oyster culture. *Venus, The Japanese J. Malacology*, Kyoto 30 (3): 113-128. [In Japanese].
- Arakawa, Y., Kubota, H.** 1974. Distribution and seasonal settlement of *Hydroides norvegica* (Gunnerus) (Polychaeta: Annelida) in Hiroshima Bay.
- Arakawa, K.Y.** 1973. Early life history of *Hydroides norvegicas* (Gunnerus) (Polychaeta: Annelida)
- Arakawa, K.Y.** 1973. Yoshokus kaki seisan no suii narabi ni kaisan seibutsu gunshu no seni kara mita. (Aspects of eutrophication in Hiroshima Bay viewed from transition of cultured yoster production and succession of marine biotic communities). *Nihon Kaiyo Gakkai-Shi* 11(2): 43-48. [In Japanese].
- Arakawa, K.Y.** 1973. Tolerances of fresh water, saturated salt water and sun-drying by *Hydroides norvegica* (Gunnerus) (Polychaeta: Annelida).
- Arakawa, K.Y.** 1973. Notes on damage to cultured oyster crop in the vicinity of Onodo, Hiroshima, caused by keen competition with blue mussel, *Mytilus edulis galloprovincialis* Lamarck.
- Arizono, M.** 1979. Disease control in mariculture, with special reference to yellowtail culture. In G. Yamamoto (ed.), *Proc. Seventh Japan-Soviet Joint Symposium on Aquaculture*, September 1978, Tokyo and Tsuruga, Japan, Tokai University. pp. 79-88.
- Arlathural, R., MacArthur, R.C., Krone, R. B.** 1977. Mathematical model of estuarial sediment transport. Tech. Rep. D-77-12. Waterways Experiment Station, Vicksburg, Mississippi.
- Åsgård, T., Storebakken, T., Austreng, E.** 1986. Reduction of self-pollution by correct feeding. *Acquacultura, Verona'86*, Abstract, Poster leaflet. 2pp.
- Aullo, K., Häkklilä, S., Mäkinen, A. & Puhakka, M.** 1985. Ecological monitoring of fish farming activities. Abstracts, 9th Baltic Marine Biologists Symposium, Turku 11.-15.6.1985, p 66.
- Aullo, K. & Häkklilä, S.** 1986. Composition of bottom deposits near a floating fish farming establishment in the SW coast of Finland. *Marine Pollution Bulletin* (manuscript).
- Aulstad, D., Gjødrem, T., Skjervold, H.** 1972. Genetic and environmental sources of variation in length and weight of rainbow trout (*Salmo gairdneri*). *J. Fish. Res. Bd. Canada* 29: 237-241.
- Austln, B.** 1982. Taxonomy of bacteria isolated from a coastal, marine fish-rearing unit. *J. Appl. Bact.* 53: 253-268.
- Austln, B.** 1983. Bacterial microflora associated with a coastal, marine fish-rearing unit. *J. mar. biol. Assoc. U.K.* 63: 585-592.

- Austin, B. 1985. Antibiotic pollution from fish farms: effects on aquatic microflora. *Microbiol. Sci.* 2 (4): 113-117.
- Austin, B., Aileen-Austin, D. 1985. Microbiol quality of water in intensive fish rearing. *J. appl. Bact.* 59 (Suppl.14): 207S-226S.
- Ayres, PaA., Barrow, G.I. 1978. The distribution of *Vibrio parahaemolyticus* in British coastal waters: rep. of a collaborative study 1975-6. *J. Hyg. Camb.* 80: 281-294.
- Balley, W.A. 1986. Assessing impacts of organotin paint use. *Oceans'86 Conference Record, Organotin Symposium, Vol.4:* 1101-1107.
- Barbay, J.R., Bradford, H.B. Jr., Roberts, N.C. 1984. The occurrence of halophilic vibrios in Louisiana coastal waters. In R.R. Colwell (ed.), *Vibrios in the Environment*. pp. 511-520. John Wiley and Sons, N.Y.
- Barltrop, J., Martin, D.F. 1984. A spectroscopic technique for evaluating sterol-aponin interactions and implications for management of *Ptychodiscus brevis* red tides. *Microbios* 41 (163), 23-29.
- Baross, J.A. 1973. Some influences of temperature, bacteriophage, and other ecological parameters on the distribution and taxonomy of marine vibrios Ph.D. thesis, University of Washington, Seattle. 451 pp.
- Baross, J.A., Liston, J. 1970. Occurrence of *Vibrio parahaemolyticus* and related hemolytic vibrios in the marine environments of Washington state. *Appl. Microbiol.* 20(2): 179-186.
- Barrie, D. 1979. Collection, preparation and identification of dinoflagellate resting cysts. pp. 443-453, In: D.L. Taylor and H.H. Sedinger (eds.), *Toxic Dinoflagellate Blooms*. Elsevier, North Holland, New York.
- Baumann, P., Baumann, L., Bang, S.S., Woolkalis, M.J. 1980. Reevaluation of the taxonomy of *Vibrio*, *Beneckeia* and *Photobacterium*: abolition of the genus *Beneckeia*. *Curr. Microbiol.* 4: 127-132.
- Bayer, J.E., Madsen, O.B.G. 1974. Computer simulations of combined ecological and Hydrodynamic processes in a sill Fjord. *Colloquia Math. Societatis Janos Bolyai* 12: 97-131.
- Bayne, B.L., Scullard, C. 1977. Rates of nitrogen excretion by species of *Mytilus* (Bivalvia: Mollusca). *J. mar. biol. Ass. U.K.* 57: 355-369.
- Bayne, B.L., Widdows, J. 1978. The physiological ecology of two populations of *Mytilus edulis* L. *Oecologia* (Berlin) 37: 137-162.
- Beaumont, A.R., Budd, M.D. 1984. High mortality of the larvae of the common mussel at low concentrations of tributyltin. *Mar. Pollut. Bull.* 15: 402-405.
- Bergheim, A., Selmer-Olsen, A.R. 1978. River pollution from a large trout farm in Norway. *Aquaculture* 14: 267-270.
- Bergheim, A., Sivertsen, A. 1981. Oxygen consuming properties of effluents from fish farms. *Aquaculture* 22(1/2): 185-187.
- Bergheim, A., Sivertsen, A., Selmer-Olsen, A.R. 1982. Estimated pollution loadings from Norwegian fish farms. I. Investigations 1978-1979. *Aquaculture* 28: 347-361.
- Beveridge, M. 1984. Cage and pen fish farming. *FAO Fisheries Technical Paper* 255. 131pp.
- Beveridge, M. Muir, J.F. 1982. An evaluation of proposed cage fish culture on Loch Lomand, an important reservoir in central Scotland. *Can. Water Resour. J.* 7 (2): 181-196.
- Beyer, J.E., Madson, O.B.G. 1974. Computer simulation of combined ecological and hydrodynamic processes in a sill fjord. *Progress in Operations Research*, Eger (Hungary), 1974, Vol. 12: 97-131.

- Bills, T.D., Marking, L.L., Mauck, W.L.** 1981. Polychlorinated biphenyl (Aroclor®1254) residues in rainbow trout: effects on sensitivity to nine fishery chemicals. *North Amer. J. Fish. Mngt.* 1: 200-208.
- Black, E.A.** 1986. Plankton kills fish. *The B.C. Mariculture Newsletter* 6(3):23.
- Black, E.A., Carswell, B.L.** 1986. The impact of salmon farming on the marine water quality. Fisheries Development Paper No. 11 (draft)
- Black, E.A., Rosenthal, H.** 1986. a survey of mariculture water quality. *British Columbia Maricult. Newsl.* 6(3): 18-22.
- Blake, P.A., Weaver, R.E., Hollis, D.G.** 1980. Diseases of humans (other than cholera) caused by vibrios. *Ann. Rev. Microbiol.* 34: 341-367.
- Braaten, B., Aure, J., Ervik, A., Boge, E.** 1983. Pollution problems in Norwegian fish farming. *ICES, C.M.* 1983/F:26: 1-11.
- Brandsma, M.G., Divoky, D.J.** 1976. Development of models for prediction of short-term fate of dredged material discharged in the estuarine environment. Prepared by Tetra-Tech, Inc., Pasadena, California for the US Army Corps of Engineers Waterways Experiment Station, Vicksburg, Mississippi.
- Bohl, M.** 1982. Production of freshwater fish in the Federal Republic of Germany in relation to the waste-water charges act. *EIFAC Tech. Paper* 41: 141-147.
- Bowden, G.** 1979. Low, politics and biology: aquaculture in the primordial ooze. In: Mann, R. (Ed.) *Exotic Species in Mariculture*. pp. 306-330. MIT Press, Cambridge Mass.
- Brady, M.S., Strobel, R.J., Katz, S.E.** in prep. In vitro analytical system for determining the ability of residue levels of antibiotics to select for resistance in bacteria.
- Brown, T.R., Gowen, R.J. Mc Lusky, D.S.** 1987. The effect of salmon farming on the benthos of a Scottish sea-loch. *J. Exp. Mar. Biol. Ecol.* (In press).
- Brown, C.M., Nash, C.E.** 1981. Aquaculture as a method for meeting hatchery discharge standards. *Proc. Bio-Engng Symp. Fish Culture, Spec. Publ. Am. Fish Soc.* pp. 183-189.
- Brown, S.W.** 1981. Water quality and off-flavor problems in commercial channel catfish culture in west-central Alabama. Auburn Univ., Off. Water Res. Tech., Washington. Rept. W82-04087.
- Bullock, G.L., Stuckey, H.M., Collis, D., Herman, R.L., Maestrone, G.** 1974. In vitro and in vivo efficacy of a potentiated sulfonamide in control of furunculosis in salmonids. *J. Fish. Res. Board Can.* 31: 75-82.
- Bullock, G.L., Snieszko, S.F.** 1975. Hagerman redmouth, a disease of the Enterobacteriaceae. U.S. Dept. of the Interior, Fish and Wildlife Service, Fish Disease Leaflet 42.
- Burrows, R.E.** 1972. Effects of accumulated excretory products on hatchery-reared salmonids. *Res. Report* 66: 1-11.
- Butman, C.A., Grant, W.D., Stolzenbach, K.D.** 1986. Predictions of sediment trap biases in turbulent flows: a theoretical analysis based on observations from the literature. *J. Mar. Res.* ---
- Butz, I., Vens-Cappell, B.** 1981. Organische Belastung des Wassers mit Stoffwechselprodukten von Forellen bei Verfütterung von Trockenfutter. [Organic load from the metabolic products of rainbow trout fed with dry food.]. *Fisch und Umwelt* 10: 103-125.
- Butz, I., Vens-Cappell, B.** 1982. Organic load from the metabolic products of rainbow trout fed with dry food. *EIFAC Tech. Paper* 41: 73-82.
- Campos, M.J., Marino, J.** 1982. Comparisons of phytoplankton species composition and abundance in the Rias Arosa and Muros (N.W. Spain). *Mar. Ecol.* 3(1): 1-12

- Cardwell, R.D., Sheldon, A.W. 1986. A risk assessment concerning the fate and effects of tributyltins in the aquatic environment. Oceans'86 Conference Record, Organotin Symposium, Vol.4: 1117-1129.
- Carlucci, A.F., Pramer, D. 1960. An evaluation of factors affecting the survival of *Escheria coli* in sea water. Appl. Microbiol. 8(4): 251-254.
- Carmark, E.C., Wiegand, R.C., Daley, R.J., Gray, C.B.J., Jasper, S., Phara, C.H. 1986. Mechanisms influencing the circulation and distribution of water masses in a medium residence-time lake. Limnol. Oceanogr. 31(2): 249-265.
- Carmichael, W.W. 1986. Algal Toxins. In: Callow, J.A. (Ed.). Advances in Botanical Research Vol. 12: 47-101, Academic Press.
- Carpenter, R.L., et.al. 1976. Aquaculture as an alternative wastewater treatment system. Philadelphia, Pa., Univ. Pennsylv. Press, 215-224. (Ref. in : Pollution Abstract 78: 05374).
- Cattell, S.A. 1969. Dinoflagellates and vitamin B12 in the Strait of Georgia, B.C. Ph.D. Thesis. University of British Columbia, Vancouver, B.C., 132 pp.
- Chew, K.K. 1984. Recent advances in the cultivation of molluscs in the Pacific United States and Canada. Aquaculture 39: 69-81.
- Cho, C.-H., Kim, Y. S. 1977. Microenvironment in oyster farm area. 1. On the eutrophication and raft density in Geoje Bay. Bull. Korean Fish. Soc. 10(4): 259-265.
- Cho, C.-H., Yang, H.-S., Park, K.-Y., Youm, M.-K. 1982. Study on bottom mud of shellfish farms in Jinhae Bay. Bull. Korean Fish. Soc. 15 (1): 35-41.
- Cho, L.-H., Park, K.-Y. 1983. Eutrophication of bottom mud in shellfish farms, the Goseong-Jaran Bay. Bull. Korean Fish. Soc. 16 (3): 260-264.
- Choubert, G., Luquet, P. 1982. Fixation and retention of canthaxanthin in rainbow trout. (Fixation et retention musculaire de la canthaxanthine par la truite arc-en-ciel). Ann. Zootech. 31(1): 1-10. [In French].
- Christie, N.D., Moldan, A. 1977. Effect of fish factory effluent on the benthic macrofauna of Saldanha Bay. Mar. Pollut. Bull. 8: 41-45.
- Clark, E.R., Herman, J.P., Forster, J.R.M. 1985. Production of metabolic and waste products by intensively farmed rainbow trout (*Salmo gairdneri*, Richardson). J. Fish. Biol. 27: 381-391.
- Cleary, J.J., Stebbing, A.R.D. 1985. Organotin and total tin in coastal waters of southwest England. Marine Pollut. Bull. 16: 350-355.
- Coëurdacier, J.M., Héral, M., Liange, M., Moreau, J., Prou, J., Razet, D., Williot, P. 1983. Rejets d'élevage de poissons en eau de mer: conséquences pour l'huître *Crassostrea gigas*. Rev. Trav. Inst. Peches Marit. 46(2): 121-140.
- Collins, I. 1983. A study on the environmental impact of particulate matter derived from a salmonid cage culture system on L. Fad, Isle of Bute, Scotland. B.Sc. Thesis, University of Stirling.
- Colwell, R.R., Liston, J. 1960. Microbiology of shellfish. Bacteriological study of the natural flora of Pacific oysters (*Crassostrea gigas*). Appl. Microbiol. 8: 104-109.
- Colwell, R.R., Grimes, D.J. 1984. Vibrio diseases of marine fish populations. Helgoländer Meeresunters. 37: 265-287.
- Colwell, R.R., West, P.A., Maneval, D., Remmers, E.F., Elliot, E.L., Carlson, N.E. 1984. Ecology of pathogenic vibrios in Chesapeake Bay. In: Colwell, R.R. (Ed.). Vibrios in the Environment. pp. 367-387. John Wiley and Sons, New York.

- Comi, G., Cantoni, C., Traaldi, C. 1981. Streptomyces in muds of trout-farming waters. (Streptomyces in fanghi di acque di allevamento di trote). Arch. Vet. Ital. 32(3-4): 21-24. [In Italian].
- Corliss, J.P. 1979. Accumulation and depletion of oxytetracycline in juvenile white shrimp (*Penaeus setiferus*). Aquaculture 16:1-6
- Crisetlg, G., Lucchini, M., Poletti, R. 1982. Heavy metals in cultured eels (*Anguilla anguilla*). (Metalli pesanti in anguille allevate.) Arch. Vet. Ital. 33(3-4): 50-55. [In Italian].
- Crockett, A.B., Wiersma, G.B., Tai, H., Mitchell, W. 1975. Pesticide and mercury residues in commercially grown catfish. Pestic. Monit. J. 8 (4): 235-240.
- Dahlbäck, B., Gunnarsson, L.A.H. 1981. Sedimentation and sulfate reduction under a mussel culture. Mar. Biol. 63 (3): 269-275.
- Dale, B. 1979. Collection, preparation, and identification of dinoflagellate resting cysts. In: Taylor, D.L., Seliger, H.H. (Ed.). *Toxic Dinoflagellate Blooms*. Proc. 2nd Int. Conf. Toxic Dinoflagellate Blooms. Development in Marine Biology Vol. 1: 443-452. Elsevier, North Holland/ New York.
- Danil'chenko, O.P., Stroganov, N.S. 1974. Evaluation of the effect of small concentrations of tripropyltin chloride on the mollusc, *Lymnaea stagnalis* L. based on a change of reproduction indices. Biol. Nauki 17(6): 130-134. [In Russian].
- Davies, I.M., McKie, J.C., Paul, J.D. 1986. Accumulation of tin and tributyltin from anti-fouling paint by cultivated scallops (*Pecten maximus* L.) and Pacific oyster (*Crassostrea gigas*). Aquaculture 55: 103-114.
- Davies, I.M., Paul, J.D. 1986a. Accumulation of copper and nickel from anti-fouling compounds during cultivation of scallops (*Pecten maximus*) and Pacific oysters (*Crassostrea gigas* Thun.). ICES, C.M. 1986/F:11: 1-15.
- Davies, I.M., Paul, J.D. 1986b. Accumulation of copper and nickel from anti-fouling compounds during cultivation of scallops (*Pecten maximus*) and Pacific oysters (*Crassostrea gigas* Thun.). Aquaculture 55:93-102.
- Davis, J.H. 1956. Influence of man upon coast lines. In W.L. Thomas (ed.), *Man's Role in Changing the Face of the Earth*. pp. 504-521. Uni. of Chicago Press, Chicago. 714 pp.
- Dean, J.C., Nielsen, L.A., Flick, G.J. 1983. Seafood processing wastes as fish meal substitutes in pelleted catfish feeds. VPI/SG, Blacksburg, VA 22pp. NTIS Order No. PB84- 104207.
- Derksen, G. 1980. Water quality and macroinvertebrate community structure of Bessette Creek, B.C.: An interior stream adjacent to a sewage lagoon. Environmental Protection Branch, Environmental Protection Service, Pacific Region, Regional Program Report 80-2, 32 pp.
- Deslous-Paoli, J.M., Héral, M. 1984. Transferts énergétiques entre la nourriture potentielle disponible dans l'eau d'un bassin ostréicole et l'huître *Crassostrea gigas* agée de un an. *Haliotis* 14: 79-90.
- Dixon, D.R., Prosser, H. 1986. An investigation of the genotoxic effects of an organotin antifouling compound (Bis-(Tributyltin) oxide) on the chromosomes of the edible mussel, *Mytilus edulis*. *Aquatic Toxicol.* 8: 185-195.
- Dooley, C.A., Taylor, V. 1983. Tributyltin compounds in the marine environment; uptake and sorption behaviour. NOSC Technical Report.
- Dooley, C.A., Vafa, G. 1986. Butyltin compounds and their measurements in oyster tissues. Oceans'86 Conference Record, Organotin Symposium, Vol.4: 1171-1176.
- Ducklow, H.W., Purdie, D.A., LeB Williams, P.J., Davies, J.M. 1986. Bacterioplankton: A sink for carbon in a coastal marine plankton community. *Science* 232: 865-867.

- Edwards, A., Edelsten, D.J. 1976. Marine fish cages - the physical environment. Proc. R. Soc. Edinburgh Sect. B 75: 208-221.
- Edwards, A., Edelsten, D.J. 1977. Deep water renewal of Loch Etive: a three basin Scottish fjord. Estuar. Coast. Mar. Sci. 5: 575-595.
- Egidius, E. 1984. Diseases of salmonids in aquaculture. Helgoländer Meeresunters. 37: 547-569.
- Eklund, J. 1986a. Rainbow trout farming polluting the Archipelagic Sea. Can. Transl. Fish. Aquat. Sci. No. 5251: 1-8. (Original Title: Kirjolohen kasvatus pilaa Saaristomerta. Suomen Luonoto 43(4): 24-27, 1984).
- Eklund, J. 1986b. Fish farm polluting the Archipelagic Sea. Can. Transl. Fish. Aquatic. Sci. 5269: 1-9. (Original Title: Fiskodlingarna fororenar Skargardshavet. Finlands natur 43(4): 21-23, 1984).
- Eley, R.L., Carroll, J.H., DeWoody, D. 1972. Effects of caged catfish culture on water quality and community metabolism of a lake. Proc. Oklahoma Acad. Sci. 52:10-15.
- Ellis, J.E., Tackett, D.L., Carter, R.R. 1978. Discharge of solids from fish ponds. Progr. Fish-Cult. 40 (4): 165-166.
- Enell, M. 1982. Changes in sediment dynamics caused by cage culture activities. Proc. 10th Nordic Symposium on Sediments, pp. 72-88.
- Enell, M., Löf, J. 1983a. Environmental impact of aquaculture - sedimentation and nutrient loadings from fish cage culture farming. (Miljöeffekter av Vattenbruk - Sedimentation och närsaltbelastning från Fiskassöodlingar). Vatten 39: 364-375. [In Swedish].
- Enell, M., Löf, J. 1983b. Changes in sediment phosphorus-, iron- and manganese- dynamics caused by cage fish farming impact. Nordic Symp. Sediments 11: 80-89.
- Ervik, A., Johannessen, P., Aure, J. 1985. Environmental Effects of marine Norwegian fish farms. Int. Council Explor. of the Sea, C.M. 1985, F: 37, Sess. W.
- Eskelinen, P. 1986. The phosphorus balance of rainbow trout. Can. Transl. Fish. Aquat. Sci. No. 5274: 1-13. (Original Title: Fosforiainestase kirjohella. Vestihallituksen monistesarja 241: 33-42, 1984).
- Evelyn, T.P.T. 1972. A multispecies fish kill in Nanoose Harbour, B.C., September 29, 1971. Tech Rep. Fish Res. Board Ca. 319: 1-14.
- Evlson, L.M. 1985. Bacterial pollution of coastal waters in the UK and Mediterranean. J. Appl. Bacteriol. Symp. Suppl 1985: 81S-93S.
- Fenchel, T.M., Reldi, R.J. 1970. The sulfide system: a new biotic community underneath the oxidized layer of marine sand bottoms. Mar. Biol. 7: 255-268.
- Fletcher, M. 1979. The aquatic environment. In J.M. Lynch and N.J. Poole (eds.), Microbial Ecology: a Conceptual Approach, pp. 92-114. London: Blackwells.
- Forfar, J.O., Keay, A.J., Maccabe, A.F., Gould, J.C., Bain, A.D. 1966. Liberal use of antibiotics and its effect in neonatal staphylococcal infection, with particular reference to erythromycin. Lancet 11: 295-300.
- Fribourg, J.H., Meyer, F.P., Robinson, J.A. 1969. Oxytetracycline leaching from medicated fish feeds. U.S. Bureau Sport Fish. Wildlife, Tech. Paper 40: 1-7.
- From, J., Rasmussen, G. 1984. Growth, gastric evacuation and body composition in rainbow trout, *Salmo gairdneri* Richardson. Dana 3:61-139.
- Fryer, J.L., Rohovec, J.S. 1984. Principal bacterial diseases of cultured marine fish. Helgoländer Meeresunters. 37: 533-545.

- Gaines, G., Taylor, F.J.R.** 1986. A mariculturist's guide to potentially harmful marine phytoplankton of the Pacific Coast of North America. B.C. Ministry of Environment, Information Report No. 10: 1-55.
- Galkina, V.N., Kulakovskij, Eh.E., Kunin, B.L.** 1982. Effect of mussel culture on environmental conditions in the White Sea. *Okeanologiya* 22 (2): 321-324.
- Gates, K.W., Perkins, B.E., EuDaly, J.G., Harrison, A.S., Bough, W.A.** 1985. The impact of discharges from seafood processing on southeastern estuaries. *Estuaries* 8 (2B): 244-251.
- Gavalchin, J.** 1983. The persistence and significance of antibiotic residues in soils. Ph.D. thesis, Rutgers State University, New Brunswick, New Jersey.
- Gergel, J., Blaha, J., Lavicky, K., Kalenda, M.** 1983. The effect of duck farming on the quality of pond water. *Bul. Vyzk. Ustav. Ryb. Hydrobiol., Vodnany* 19 (3): 24-31.
- Gibbs, P.E., Bryan, G.W.** 1986. Reproductive failure in populations of the dog-whelk, *Nucella lapillus*, caused by imposex induced by tributyltin from antifouling paints. *J. mar. biol. Assoc. U.K.* 66: 767-777.
- Gomez-Parra, A., Establier, R., Blasco, J.** 1984. Heavy metal distribution in superficial sediments in the sea channels around the fish ponds of the Bay of Cadiz, southwestern Spain. *Inf. Tech. Inst. Invest. Pesq.* 0(120): 3-20.
- Gordon, M.S., Chapman, D.J., Kawasaki, L.Y., Tarifeño-Silva, E., Yu, D.P.** 1982. Aquacultural approaches to recycling of dissolved nutrients in secondarily treated domestic wastewaters. IV. -conclusions, design and operational considerations for artificial food chains. *Water Res.* 16: 67-71.
- Gowen, R. J.** 1984. The ecology of phytoplankton in Scottish coastal waters with particular reference to toxic species and their importance to mariculture. A final report for the Highlands and Islands Development Board with a Preface by P. Tett. 91 pp. (Mimeographed, unpublished).
- Gowen, R.J., Bradbury, N.B.** 1987. The ecological impact of salmonid farming in coastal waters: a review. *Oceanogr. Mar. Biol. Ann. Rev.* 25: 563-575.
- Gowen, R.J., Lewis, J., Bullock, A.M.** 1982. A flagellate bloom and associated mortalities of farmed salmon and trout in Upper Loch Fyne. *Scottish Marine Biol. Assoc., Internal Rep.* 71: 1-15.
- Gowen, R.J., Bradbury, N.B., Brown, J.R.** 1985. The ecological impact on salmon farming in Scottish coastal waters: A preliminary appraisal. *ICES, C.M. 1985/F:* 35, 1-11. (Mariculture Committee).
- Grabacka, E.** 1977. The influence of beet sugar factory wastes on the bottom microfauna in fish ponds. (Wpływ 'ścieko'w cukrowniczych na mikrofaunę dna w stawach rybnych), *Acta Hydrobiol.* 19(4): 373-387.
- Granéli, E., Granéli, W.** 1982. Eutrophication and dinoflagellate blooms in Swedish coastal waters - possible causes and counter measures. *Int. Symp. on Utilization of Coastal Ecosystems, Brazil, Nov. 22-27*, 32 pp.
- Granéli, E., Persson, H., Edler, L.** 1986. Connection between trace metals chelators and red tide blooms in the Laholm Bay, SE Kattegatt. An experimental approach. *Mar. Environ. Res.* 18: 61-78.
- Grovhoug, J.G., Sellgman, P.F., Vata, G., Fransham, R.** 1986. Baseline measurements of butyltin in U.S. harbors and estuaries. pp. 1283-1288. In *Oceans 86 Conference Record, Washington, D.C., September 23-25, 1986.*
- Gulral, D.** 1986. Modification et transformations des écosystèmes sédimentaires par des élevages piscicoles en Lagune Ebrié (Cote D'Ivoire). *Aquaculture* 52: 287-302.

- Günther, U.** 1976. Hydrochemische Untersuchungen in zwei schleswig-holsteinischen Forellennastbetrieben verschiedener Intensität und die durch die Betriebe verursachten Vorfluterbelastungen, 1972-1973. [Hydrochemical studies in two Schleswig-Holstein trout fattening units of different intensity and their effects on the receiving waters]. *Afz Fischwirt* 26(9): 49-52.
- Haavisto, P.** 1974. The magnitude of loading from natural fish ponds and from fish farms. *Rep. Nat. Bd. Wat. Finland* 74: 1-67.
- Hagino, S.** 1977. 3. Physical Properties of the pollutants. Source: Senkai Yoshoku to Jika Osen [Shallow-Sea Aquaculture and Self-Pollution]. *Suisan-gaku Shirizu* 21 [Fisheries Series 21]. By Koseisha Koseikaku. 134 pp. (In Japanese) pp.31-41.
- Hagström, Å., Larsson, A.-M.** 1985. Pelagic bacteria and mussel culture. [Pelagiska bakterier och mussellodling]. *Medd. Havsfisk. Lab. Lysekil* No 286: 1-21.
- Håkanson, L.** 1986a. Environmental impact of fish cage farms. In: *Aquaculture- an Industry for the future. IVA-Symposium (Royal Swedish Academy of Engineers, Stockholm). Rapport* 308: 179-199.
- Håkanson, L.** 1986b. The swedish coastal zone project: sediment types and morphometry. In: "Sediments and water interactions." *Proc. 3rd Int. Symp. on Interactions between Sediment and Water, Geneva, Switzerland, August 27-31, 1984*, pp. 35-51.
- Håkanson, L., Kvarnäs, H., Karlsson, B.** 1986. Coastal morphometry as regulator of water exchange -- a Swedish example. *Estuarine & Coast. Shelf Sci.* 23:
- Häkklilä, S.** 1982. Pohjaeläimistön tila Kustavin kalankasvatustaitosten vaikutuspiirissä 1980-81. (The condition of the bottom fauna in the area under the influence of the fish farms in the Kustavi area 1980-81.) *Vesihallituksen monistesarja* 110, 1982, 29 pp. [In Finnish].
- Häkklilä, S.** 1984. Särkängsalmen pohjaeläintutkimus 1983. (The study of the bottom animals in the sound of Särkängsalmi 1983.) *Turun yliopisto, Saaristomeren tutkimuslaitos. Mimeo*, 7 pp. [In Finnish].
- Hales, L.Z.** 1981. Floating Breakwaters: State-of-the-Art Literature Review. *Tech. Rep. No.* 81-1.
- Hall, P., Holby, O.** 1986. Environmental impact of a marine fish cage culture. *ICES, C.M.* 1986/F/46, 1-19. (Mariculture Committee, Ref. MEQC).
- Hall, P., Holby, O.** 1985. Environmental impact of marine fish farming - sedimentation and benthic solute in-situ fluxes under a fish cage culture in Bohuslän. *Svenska havsforskningsföreningens Medd.* No. 20: 141-159.
- Hallikainen, S.** 1986. Kalankasvatuksen vaikutukset luonnonkaloihin ja kalastukseen (The impact of fish farming on natural fish and fisheries.) *INSKO* 282-86 III, 11 pp. [In Finnish].
- Halonen, L.** 1985: Kalankasvatuksen vaikutukset rannikkovesissä. (The impacts of fish farming in coastal waters.) *Vesihallituksen monistesarja* nro 346, 1985, 42 pp. [In Finnish].
- Hata, Y, Katayama, H.** 1977. Shallow-Sea aquaculture and shelf-pollution. 5. Chemical processes of the pollutants. *Suisan-gaku Shirizu* 21: 52-66. (Fisheries Series).
- Hatakeyama, K., Nishimura, T., Nagasaki, S.** 1985. Study of the motion and the wave force on the hanging seaweed in the ocean food and energy farm. 20: 105-117.
- Haven, D.S., Morales-Alamo, R.** 1966a. Aspects of biodeposition by oysters and other invertebrate filter feeders. *Limnol. Oceanogr.* 11: 487-498.
- Haven, D.S., Morales-Alamo, R.** 1966b. Use of fluorescent marine particles to trace oyster biodeposits in marine sediments. *J. Cons. perm. int. Explor. Mer* 30: 267-269.
- Hays, T.** 1980. Impact on net pen culture on water quality and fish populations on Bull Shoals Reservoir. *Arkansas Game and Fish Commission Little Rock, AK (USA)*, 1980, 35 pp. (AGFC-Proj-2-338-R).

- Hechanova, R.G.** 1982. Some notes on site selection for coastal fishfarms in Southeast Asia. In: Report of Consultation /seminar on coastal fishpond engineering, 4-12 Aug. 1982. Surabaya, Indonesia. FAO/UNDP, Manila (Philippines) pp.72-75.
- Henderson, E.B.** 1985. Guidelines for Physical separation of salmonid aquaculture farms. (unpublished Manuscript). St. Andrews, Biol. Station DFO, Canada) 24pp.
- Henderson, R.S.** 1986. Effects of organotin antifouling paint leachates on Pearl Harbour organisms: A site specific flowthrough bioassay. Oceans '86 Conference Record, Organotin Symposium, Vol.4: 1226-1233.
- Héral, M., Deslous-Paoli, J.M., Sornin, J.M.** 1984. Transferts énergétiques entre l'hiure *Crassostrea gigas* et la nourriture potentielle disponible dans un bassin ostréicole: premières approches. Océanis 9(13): 169-194.
- Herman, R.L.** 1969. Oxytetracycline in fish culture - a review. U.S. Bureau Sport Fish. Wildlife, Tech Pap. 31:1-9.
- Hilge, V., Huschenbeth, E., Harms, U.** 1984. Trace contaminants in catfish raised under intensive culture conditions. ICES/CM @984/E:30, 1-13 (Marine Environmental Quality Committee).
- Hinshaw, R.N.** 1973. Pollution as a result of fish cultural activities. U.S. Environmental Protection Agency, Washington, D.C. EPA-R3-73-009.
- Hinshaw, R.W., Hilden, D.A.** 1973. Pollution as a result of fish culture activities. EPA Ecol. Res. Ser. 7-73-009.
- Hls, E., Robert, R.** 1985. Développement des véligeres de *Crassostra gigas* dans le bassin d'Arachon études sur les mortalités larvaires. Rev. Trav. Inst. Peches marit. 4 (1-2): 63-88. [In French].
- Hisaoka, M., Nogami, K., Takeuchi, O., Suzuki, M., Sugimoto, H.** 1966. Studies on sea water exchange in fish farms. II. Exchange of sea water in floating net. Bull. Nakai Reg. Res. Lab., Contr. No. 115:21-43. [In Japanese].
- Hoffmann, E.** 1986. Marin akvakultur og miljøeffekter. Vand & Miljø 5(3): 195-199.[In Danish].
- Holmes, P.R., Lam, C.W.Y.** 1985. Red tides in Hong Kong waters - response to a growing problem. Asian Mar. Biol. 2: 1-10.
- Holt, R.A., Conrad, J.F., Fryer, J.F.** Furanace for control of *Cytophaga psychrophila*, the causative agent of cold-water disease in coho salmon. Prog. Fish. Cult. 37(3): 137-139.
- Horiguchi, Y., Maegawa, M.** 1978. Studies on marine environments and mariculture resources in the waters around Zaga-shima (Ago Bay)- III. Effects of dredging works upon growth and mortality of operated pearl oyster and quality of pearl.
- Hurlburt, E.F.** 1984. Aquaculture in Puget Sound -its potential and its possible environmental impact. State of Washington, Department of Fisheries, Olympia, Washington.
- ICES**, 1985. Report of the working group on exceptional algal blooms. ICES, C.M. 1985/F:58/Sess. T.
- Iigura, T.** 1974. Sinking speed of faeces and uneaten food in hamachi culture. Suisan zoshoku 22: 34-39. [In Japanese].
- Inoue, H.** 1972. On water exchange in a net cage stocked with the fish hamachi. Bull. Jap. Soc. Sci. Fish. 38(2): 167-176. [In Japanese].
- Inoue, H., Tanaka, Y., Fukuda, K.** 1970. On water exchange in shallow marine fish farms. II. Hamachi fish farm at Tanoura. Bull. Jap. Soc. Sci. Fish. 36(8): 776-782. [In Japanese].
- Inoue, H., Tanaka, Y., Salto, M.** 1966. On the water exchange in the shallow marine fish farm. I. Hitsuishi fish farm of hamachi. Bull. Jap. Soc. Sci. Fish. 32(5): 384-392. [In Japanese].

- Isajl, T. Spaulding, M.L., Stace, J. 1985. Tidal exchange between a coastal lagoon and offshore waters. *Estuaries* 8 (28): 203-216.
- Ito, S., Imal, T. 1955. Ecology of oyster bed. I. On the decline of productivity due to repeated cultures. *Tohoku J. Agric. Res.* 5(4): 251-268.
- Jana, B.B., Patel, G.N. 1984. Measurements of some groups of microbial populations with special reference to denitrification in water and sediments of fish farming ponds under mono- and polyculture systems. *Int. Revue ges. Hydrobiol.* 69(2): 231-240.
- Jarrams, P., Starkle, A., Easton, K., Templeton, R.G. 1980. Salmonid rearing in floating net cages in freshwater reservoirs owned by the Severn Trent Water Authority. *Fish. Mgmt.* 3: 101-117.
- Jensen, R. 1972. Taking care of wastes from the trout farm. *Am. Fish. US Trout News* 16 (5): 4-6.
- Johnson, H.E., Duffield, D.J. 1980. Utilization of wastewater for intensive fish culture. Michigan State Univ, Rep. PC A03/MF AQ1, 41pp.
- Johnstone, A.K. 1984. Pathogenesis and life cycle of the myxozoan *Parvicapsula* sp infecting marine cultured coho salmon. Ph. D. Thesis, University of Washington, Seattle.
- Jones, K.J., 1983. Water quality at marine fish farms at the Hunterston A power station in relation to phytoplankton composition and abundance. Final report on work commissioned by Golden Sea Products Ltd., Easgan Ltd. and Sea Fish Industry Authority. *Scottish Mar. Biol. Assoc., Internal Rep*, 91: 1-31.
- Jones, K.J., Ayres, P.A., Bullock, A.M., Roberts, R.J., Tett, P. 1982. A red tide of *Gymnodinium aureolum* in sea lochs of the Firth of Clyde and associated mortalities of pond reared salmon. *J. mar. biol. Assoc. U.K.* 62: 771-782.
- Jørgensen, O.H. 1982. Legislative measures concerning trout farming in Denmark. EIFAC Tech. Paper 41: 137-140.
- Jory, D.E., Carrlker, M.R., Iversen, E.S. 1984. Preventing predation in molluscan mariculture: an overview. *J. World Maricult. Soc.* 15: 421-432.
- Jumppanen, K. 1986: AB Skärgårdens lax - saariston lohi OY fish farms environmental impacts. (in Finnish, AB Skärgårdens lax - Saariston lohi OY:n uivan kalankasvatuslaitoksen ympäristövaikutukset). Lounais-Suomen vesienhuoltoyhdistys r.y. Julkaisu 64, 100 pp.
- Kadowaki, S., Hirata, H. 1984. Distribution of dissolved oxygen in a shallow-sea fish farm. 1. Effects of feeding on the horizontal oxygen distribution. *Suisan Yoshoku (The Aquiculture)* 32(3): 142-147. [In Japanese].
- Kadowaki, S., Kasedo, T., Nakazono, T. 1978. Continuous records of DO contents by cruising in the coastal culture farms -I. Relation between DO content and fish density in cages. *Mem. Fac. Fish., Kagoshima Univ.* 27 (1): 273-280. [In Japanese].
- Kadowaki, S., Kasedo, T., Nakazono, T., Hirata, H. 1978. Continuous records of DO contents by cruising in the coastal culture farms -II. Diffusion of suspended particles by feeding. *Mem. Fac. Fish., Kagoshima Univ.* 27 (1): 281-288. [In Japanese].
- Kadowaki, S., Kasedo, T., Nakazono, T., Yamashita, Y., Hirata, H. 1980. The relation between sediment flux and fish feeding in coastal culture farms. *Mem. Fac. Fish., Kagoshima Univ.* 29: 217-224. [In Japanese].
- Kalresalo, T. 1979. Effects of rainbow trout breeding on winter decomposition activity and oxygen content in Lake Langelmavesi, south Finland. *Ann. Bot. Fenn.* 16(3): 176-180.
- Kalashnik, V.I., Kirilenko, N.S., Korobleva, A.I. 1984. Effectiveness of the use of piscicultural ponds in the practice of wastewater purification. *Samoochishchenie Vody Migr. Zagryaz. Troficheskoi Tsepi*, [Sb. Tr. Vses. Soveshch. Sanit. Gidrobiol.], 4th 1981 (Publ. 1984): 122-125. [In Russian].

- Kalfus, M., Korzenlewski, K. 1982. A mathematical model for optimization of intensive trout culture in running waters. Pol. Arch. Hydrobiol. 29(3-4): 693-702.
- Kapetsky, J.M. 1982. The problem of predicting seasonal and longterm variations in parameters affecting site suitability in coastal aquaculture. In: Report of consultation/seminar on coastal fishpond engineering, 4-12 Aug. 1982, Surabaya, Indonesia. FAO/UNDP Manila (Philippines) pp. 46-48.
- Kaspar, H.F., Gillespie, P.A., Boyer, I.C., MacKenzie, A.L. 1985. Effects of mussel aquaculture on the nitrogen cycle and benthic communities in Kenepuru Sound, Marlborough Sounds, New Zealand. Marine Biol. 85: 127-136.
- Katsuda, S., Taba, N., Chimi, K. 1984. Comparative study on the effects of gonyautoxins and tetrodotoxin on the cardiovascular function of rabbits. Shokuhin Eiseigaku Zasshi 25 (4): 329-333. [In Japanese].
- Katz, S.E. unpubl. Manuscript 1984. Environmental impact assessment for the use of oxytetracycline to control *Gaflkemia* infections in lobsters. November 26, 1984. Prepared for the U.S. Food and Drug Administration.
- Kautsky, N., Wallentinus, P.A. 1980. Nutrient release from a Baltic *Mytilus* -bed algal community and its role in benthic and pelagic productivity. Ophelia, Suppl. 1: 17-30.
- Kawasaki, L.Y., Tarifeño-Silva, E., Yu, D.P., Gordon, M.S., Chapman, D.J. 1982. Aquacultural approaches to recycling of dissolved nutrients in secondarily treated domestic wastewaters - I. Nutrient uptake and release by artificial food chains. Water Research 16: 37-49.
- Kay, B.H. 1981. Shellfish growing water control program, Annual Review 1980-1981. Okeover Inlet and surrounding area, Desolation Sound, Quadra Island, Cortes Island, Redonda Island, Texada Island, Lund Harbour and Scuttle Bay to Powell River. Environment Canada, Environmental Protection Service, Pacific Region, Regional Program Report 81-2, 105 pp.
- Kay, B.H., Walker, D. 1985. Shellfish growing water bacteriological and sanitary survey of Ladysmith Harbour, Davis Lagoon, Boulder Point, and Sharpe Point to Yellow Point, British Columbia, 1983-1984. Department of Environment, Environmental Protection Service, Pacific Region, Regional Program Report 84-18, 134 pp.
- Kehitysaluearahasto, OY. 1987. Kalanviljelyn verkkoalaskasvatuksen vesistökuormituksen vähentäminen. (Reduction of loading by net cage fish farming activities.) Kehitysaluearahasto OY, Tutkimus A 7:1987. 73 pp. [In Finnish].
- Kilambi, R.V., Adams, J.C., Wickizer, W.A. 1978. Effects of cage culture on growth, abundance, and survival of resident Largemouth Bass (*Micropterus salmoides*). J. Fish. Res. Board Can. 35: 157-160.
- Kilambi, R.V., Hoffman, C.E., Brown, A.V., Adams, J.C., Wickizer, W.A. 1976. Effects of cage culture fish production upon the biotic and abiotic environment of Crystal Lake, Arkansas. Arkansas Game and Fish Commission, Little Rock, Ak. U.S. Department of Commerce, National Technical Information Service. 1-145.
- Kille, U. 1979. Oxygen-regime and artificial aeration of net-cages in mariculture. Meereforsch. 27(4): 236-243.
- Kim, Y., Yoon, D.S., Kang, P.A. 1980. Studies on the hanging culture of oyster, *Crassostrea gigas*, in the Korean coastal waters. 7. Environmental condition of oyster culture beds and oyster growth. Bull. Fish. Res. Dev. Agency, Busan 25: 35-43.
- Kitamori, R. 1977. Changes in the species composition (principally benthic organisms). In Japanese Soc. Scientific Fisheries (ed.), Senkai Yoshoku to Jika Osen (Shallow-Sea Aquaculture and Self-Pollution), Suisan-gaku Shirizu 21 (Fisheries Series 21). pp. 67-76. Koseisha Koseikaku, Tokyo. [In Japanese].
- Klontz, G.W. 1982. An applied coupled simulation-optimization model of water use efficiency in intensive fish culture systems. Tech. Completion Report PC A05/MF A01 IDA, USA.

- Klontz, G.W., Chacko, A.J., Bealeu, M.H. 1980. Environmental gill disease (EGD): What it is and what to do about it. Proc. No. Pac. Aquacult. Symp., Anchorage, Alaska.
- Knösche, R. 1970. Zur Verbesserung der Wasserqualität bei der Forellenmast in Netzkäfigen. [On improvement of water quality in trout fattening in net cages.]. Dt. Fischrei-Ztg. 17(11): 328-332. [In German].
- Koditschek, L.K., Guyre, P. 1974. Resistance transfer faecal coliforms isolated from the Whippany river. Water Res. 8(10): 747-752.
- Koganezawa, A. 1978. Pests and prevention. In: Imai, T. (Ed.). Aquaculture in Shallow Seas: Progress in shallow sea culture. Koseisha Koseiku, Tokyo. pp. 249-251. Translated by the National Marine Fisheries Service and published in 1977 by Amerind Publishing Co. Pvt. Ltd., New Delhi.
- Kolvisto, V. 1987. Summary working report. Living archipelago - living water. Consequences of fish farming on coastal and archipelago area in SW Finland. (In Swedish, Sammanfattande arbetsrapport, Levande skärgård - levande vatten, fiskodlingens konsekvenser i kust- och skärgårdsområdet i SW Finland. Manuscript, 25 pp.
- Kool, B., Walker, D., Kay, B. 1985. Shellfish growing water bacteriological and sanitary survey of Nanoose Harbour, Lasqueti Island, Deep Bay and Pipers Lagoon, British Columbia 1983-1984. Department of Environment, Environmental Protection Service, Pacific Region, Regional Program Report 84-22, 65 pp.
- Korzeniewski, K., Korzeniewska, J. 1982. Changes in the composition and physiological properties of the bacterial flora of water and bottom sediments in lake Letowo, caused by intensive trout culture. Pol. Arch. Hydrobiol. 29(3-4): 671-682.
- Korzeniewski, K., Trojanowski, J., Mrozek, B. 1982. Effect of intensive trout culture on contents of nutrients in water. Pol. Arch. Hydrobiol. 29 (3-4): 625-632.
- Korzeniewski, K., Salata, W. 1982. Effect of intensive trout culture on chemistry of Lake Letowo waters. Pol. Arch. Hydrobiol. 29 (3-4): 633-657.
- Korzeniewski, K., Banat, Z., Moczulska, A. 1982. Changes in water of the Uniesc and Skotawa Rivers, caused by intensive trout culture. Pol. Arch. Hydrobiol. 29 (3-4): 683-691.
- Krishnamoorthi, K.P., Abdulappa, M.K., Sarkar, R., Siddiqi, R.H. 1975. Productivity of sewage fertilized fish ponds. Water Research 9: 269-274.
- Krogh, P. et al. 1985. Outbreak of Diarrhetic Shellfish Poisoning on the west coast of Sweden. In: Anderson et al. (Eds.) Toxic Dinoflagellates. Elsevier, North Holland/ New York, pp. 501-503.
- Kuenzler, E.J. 1961. Phosphorus budget of a mussel population. Limnol. Oceanogr. 6: 400-415.
- Kusuki, Y. 1977a. Fundamental studies on the deterioration of oyster growing grounds. I. Production of faecal materials by the Japanese oyster. Bull. Jap. Soc. Sci. Fish. 43(2): 163-166. [In Japanese].
- Kusuki, Y. 1977b. Fundamental studies on the deterioration of oyster growing grounds. II. Organic content of faecal materials. Bull. Jap. Soc. Sci. Fish. 43(2): 167-171. [In Japanese].
- Lam, C.W.Y. 1984. Anoxic effect of red tides on fishkills in Hong Kong. Paper delivered at the Red Tide Workshop 18-20 June, 1984, held at the CSIRO Marine Laboratories, Cronulla, NSW, Australia.
- Landau, M., Pierce, R.H., Ryther, J.H., Zeigler, J.A. 1982. The quality of shrimp reared in a sewage/seawater aquaculture system. Amer. Zoologist 22(2): 912. (Abstract, No. 378).
- Lapointe, B.E., Niell, F.X., Fuentes, J.M. 1981. Community structure, succession and production of seaweeds associated with mussel-rafts in the Ria de Arosa, NW Spain. Mar. Ecol.-Progr. Ser. 5: 243-253.
- Larsson, A.-M. 1985. Blue mussel sea farming - effects on water quality. Vatten 41: 218-224.

- Laughlin, R., Frech, W., Guard, H.E.** 1983. Acute and sublethal toxicity of tributyltin oxide (TBTO) and its putative environmental products, tributyltin sulfide (TBTS) to zoeal mud crabs, *Rhithropanopeus harrisi*. *Water Air Soil Pollut.* 20(1): 69-79.
- Laughlin, R.B., Jr.** 1986. Bioaccumulation of tributyltin: The link between environment and organism. *Oceans'86 Conference Record, Organotin Symposium, Vol.4:* 1206-1209.
- Lazler, J.R.N.** 1963. Some aspects of the oceanographic structure in the Jervis Inlet System.
- Ledo, A., González, E., Barja, J.L., Toranzo, A.E.** 1983. Effect of depuration systems on the reduction of bacteriological indicators in cultured mussels (*Mytilus edulis* Linnaeus). *J. Shellfish Res.* 3(1): 59-64.
- Lee, V., Olsen, S.** 1985. Eutrophication and management initiatives for the control of nutrient inputs to Rhode Island coastal lagoons. *Estuaries* 8 (2B): 191-202.
- Leminen, E., Mäkinen, T. & Junna, J.** 1986. Kalanviljelyn vesistökuormituksen vähentäminen varkkokassilaitoksella - kenttätutkimus merilolosuhteissa. (Reduction of loading by a net gage fish farm - a field research at sea circumstances). Vesihallituksen monistesarja nro 6, 1986. 32 pp. [In Finnish].
- Levin, M.A.** 1977. Water quality health effects: development of criteria for shellfish-growing waters. In: Wilt, D.S. (ed.). *Proc. Natl. Shellfish Sanit. Workshop (U.S.)* (10), FDA, Washington, DC (USA), 212 pp.
- Liao, P.B.** 1970. Pollution potential of salmonid fish hatcheries. *Water and Sewage Works* 117 (8): 291.
- Liao, P.B., Mayo, R.D.** 1972. Salmonid hatchery water reuse systems. *Aquaculture* 1: 317-335.
- Lieffrig, D.V.M.** 1985. The effects of hydrogen sulfide (H₂S) on aquaculture production. MSc. Thesis, University of Stirling, Scotland.
- Lindbergh, J.M.** (unpubl. Manuscript). Potential interaction between net pen farmed Atlantic salmon and native salmonid species in the Pacific Northwest. Prepared for Sea Farm of Norway, Inc., Greenwich, Connecticut, USA.
- Lindqvist, O.V.** 1986. Fish farming in Finland. *Can. Transl. Fish. Aquat. Sci.* No. 5273: 1-7. (Original Title: Kalanviljely Suomessa. Vesihallituksen monistesarja 241: 5-8, 1984).
- Loo, L.O., Rosenberg, R.** 1983. *Mytilus edulis* culture: growth and production in western Sweden. *Aquaculture* 35(2): 137-150.
- Loosanoff, V.L., McKenzie, C.L., Jr., Shearer, L.W.** 1960. Use of chemicals to control shellfish predators. *Science* 131: 1522-1523.
- Loosanoff, V.L.** 1961. Recent advances in the control of shellfish predators and competitors. *Proc. 13th Ann. Meeting Gulf and Caribbean Fish. Inst.* 13: 118-128.
- López-Jamar, E.** 1985. Distribución espacial del poliqueto *Spiochaetopterus costarum* en las Rías Bajas de Galicia y su posible utilización como indicador de contaminación orgánica en el sedimento. *Bol. Inst. Esp. Oceanogr.* 2 (1): 68-76. [In Spanish].
- López-Jamar, E., Iglesias, J., Otero, J.J.** 1984. Contribution of infauna and mussel-raft epifauna to demersal fish diets. *Mar. Ecol. Progr. Ser.* 15: 13-18.
- Lovegrove, T.** 1979. Anti-fouling paint in farm ponds. *Fish Farming Int.* 6(2): 13-15,39.
- Loyacano, H.A., Jr., Smith, D.C.** 1976. Attraction of native fish to catfish culture cages in reservoirs. *Proc. Ann. Conf. Southeast. Assoc. Game Fish. Commn* 29: 63-73.
- Loyacano, H.A. Jr., Smith, D.C.** 1975. Attraction of native fish to catfish culture cages in reservoirs. *Southeastern Ass. Fish Comm.* vol. 29.

- Maguire, R.J.** 1986. Review of the occurrence, persistence and degradation of tributyltin in fresh water ecosystems in Canada. pp. 1252-1255. In Oceans 86 Conference Record, Washington, D.C., September 23-25, 1986.
- Maguire, R.J., Chau, Y.K., Bengert, A., Hale, E.J.** 1982. Occurrence of organotin compounds in Ontario lakes and rivers. Environ. Sci. Technol. 16: 698-702.
- Mäkinen, T.** 1986a. Reduction of the pollution effects of a net cage station. Can. Transl. Fish. Aquat. Sci. No. 5272: 1-4. (Original Title: Vesistokuurmituksen vahentaminen verkkokassilaitoksella. Suomen Kalankasvattaja 4: 14. 1984)
- Mäkinen, T.** 1986b. Loading effect of fish farming with net cages on coastal waters - can it be reduced? Can. Transl. Fish. Aquat. Sci. No. 5268: 1-11. (Original Title: Belastningen på kustvatten från fiskodlingar i nät-kassar -- kan den minskas? Fiskeritidskrift för Finland 28(2): 36-39, 1984).
- Mäkinen, T. & Naukkarinen, M.** 1982. Mallitutkimus pyörreselkeyttimen soveltuvuudesta kalanviljelyn poistovesien käsittelyyn. (Model study on the suitability of the whirl concentrator in fish farm effluent treatment.) Vesihoitituksen monistesarja nro 128, 1982, 41 pp. [In Finnish]
- Mäkinen, T.** 1984. Avskiljning av partikulärt material från fiskodlingsanstaltens avloppsvatten. (Removal of suspended solids from fish farm effluents.). NJF-utredning/Rapport nr. 14: pp. 53-89. [In Swedish].
- Mäkinen, T.** 1985. Tekniska erfarenheter av reningsåtgärder vid fiskodlingsanläggningar. (Technical experiences of fish farm effluent treatment). Nordforsk miljövårdserien, publikation 1985:2: pp. 185-196. [In Swedish].
- Mäkinen, T. & Pursiainen, M.** 1987: Environmental impacts of mariculture in Finland. ICES ad hoc study group on environmental impacts of mariculture, Hamburg 7.-9. 4.1987. Manuscript 15 pp.
- Mann, H.** 1978. Einfluß der Käfighaltung auf den Chemismus in stehenden Gewässern. (Influence of cage culture on the chemistry of stagnant waters.). unpubl. Manuscript, 5 pp.
- Mann, R., Ryther, J.H.** 1979. Trace contaminant accumulation by organisms grown in a waste recycling aquaculture system. Proc. World Maricult Soc. 10: 809-822.
- Mantle, G.J.** 1982. Biological and chemical changes associated with the discharge of fish-farm effluent. EIFAC Tech. Paper 41: 103-112.
- Margalef, R.** 1978. Life-forms of phytoplankton as survival alternatives in an unstable environment. Oceanologica Acta 1(4): 493-509.
- Markmann, P.N.** 1982. Biological effects of effluence from Danish fish farms. EIFAC Tech. Paper 41: 99-102.
- Matsuno, T., Katsuyama, M., Maoka, T., Hirano, T., Komori, T.** 1985. Trductive metabolic pathways of carotenoids in fish: (3S,3'S)-astaxanthin to tunaxanthin A,B, and C. Comp. Biochem. Physiol. B. Comp. Biochem. 80(4): 779-790.
- Mattsson, J., Lindén, O.** 1983. Benthic macrofauna succession under mussels, *Mytilus edulis* L. (Bivalvia), cultured on hanging long-lines. Sarsia 68: 97-102.
- Mazaev, V.T., Shlepina, T.G.** 1973. Experimental data on hygienic standardization of dibutyltin sulfide in water bodies. Gig. Sanit. 38(8): 10-13. [In Russian].
- McLaughlin, T.W.** 1981. Hatchery Effluent treatment - U.S. Fish and Wildlife Service. Proc. Bio-Eng. Symp. Fish Culture Publ. by Am. Fish Soc. Spec. Publ., pp. 167-173.
- Meador, J.P., U'Ren, S.C., Salazar, M.H.** 1984. A flow-through bioassay system for the evaluation of organotin antifouling compounds. Water Res. 18(5): 647-650.

- Menasveta, P.** 1982. Environmental considerations for the development of coastal fishfarms in the Indo-Pacific region. In: Report of consultation/seminar on coastal fishpond engineering, 4-12 Aug. 1982, Surabaya, Indonesia. FAO/UNDP Manila (Philippines) pp. 29-35..
- Merican, Z.O., Phillips, M.J.** 1985. Solid waste production from rainbow trout, *Salmo gairdneri* Richardson, cage culture. *Aqua. Fish. Mgmt.* 1: 55-69.
- Meyers, T.R.** 1984. Marine bivalve mollusks as reservoirs of viral pathogens: significance to marine and anadromous finfish aquaculture. *Mar. Fish. Rev.* 46(3): 14-17.
- Mildtylling, P.V.** 1985. Bruk av medikamenter og desinfeksjonsmidler i norske fiskoppdrettsanlegg. (Use of drugs and disinfectants in Norwegian fish farms). *Vann* 3: 177-180. [In Swedish].
- Milbrink, G., Holmgren, S.** 1981. Addition of artificial fertilizers as a means of reducing negative effects of "oligotrophication" in Lakes after impoundment. *Rep. Inst. Freshwater Res., Drottningholm* 59: 97-120.
- Miner-Rensel Assoc.** 1986. Aquatic conditions at the Seafarm of Norway net-pen site in Port Angeles harbour in April, 1986. Prepared for Sea Farm of Norway, Port Angeles, Washington.
- Minchew, C.D., Yarbrough, J.D.** 1977. The occurrence of fin rot in mullet (*Mugil cephalus*) associated with crude oil contamination of an estuarine pond ecosystem. *J. Fish. Biol.* 10: 319-323.
- Minchin, D., Duggan, C.B.** 1986. Organotin contamination in Irish waters. *ICES, CM/1986/F:48*, 1-9.
- Misdorp, R., Kohslek, L.H.M., Steyaert, F.H.I.M., Dijkema, R.** Environmental consequences of a large scale coastal engineering project on aspects of mussel cultivation in the Eastern Scheldt. *Nat. Sci. Tech.* 16: 95-105.
- Miyake, Y., Fujisawa, K.** 1985. Survey of sediment in the oyster *Crassostrea gigas* farms. III. Hinase waters. *Bull. Fish. Exp. Stat. Okayama Pref.* 59: 142-144. [In Japanese].
- Miyake, Y., Fujisawa, K.** 1985. Survey on water quality of oyster farms in eastern parts of Okayama Prefecture. *Bull. Fish. Exp. Stat. Okayama Prefecture* 59: 131-140. [In Japanese].
- Morantz, D.L., Barbour, S.E., Sweeney, R.K.** 1986. Source of error in water velocity measurement for aquatic studies. *Can. J. Fish. Aquat. Sci.* 43: 893-896.
- Mudrak, V.A.** 1981. Guidelines for economical commercial fish hatchery wastewater treatment systems. *Proc. Bio-Engng Symp. Fish Culture*. Allen, L.J., Kinney, E.C. (Eds.) Publ. by Am. Fish. Soc, Spec. Publ. pp. 174-182.
- Muir, J.F.** 1982. Economic aspects of waste treatment in fish culture. *EIFAC Tech. Paper* 41: 123-135.
- Müller-Haeckel, A.** 1986. Control of water quality around a cage fish farm in the Norrby archipelago (Northern Bothnian Sea). *Vatten* 42: 205-209.
- Munro, K.A., Sams, S.C., Nassichuk, M.D.** 1985. The effects of hatchery effluents on water chemistry, periphyton and benthic invertebrates of selected British Columbia Streams. *Can. Manuscript Rep. Fish. aquatic Sci.* 1830: 1-203.
- Murakami, A.** 1973. Eutrophication of the hydrosphere and the increase of fisheries culture. In : *Water Pollution and Eutrophication*. pp. 37-48. *Jap. Fish. Soc., Koseisha kosei Kakuhon* (cited in Arakawa, 1973). [In Japanese].
- Murakami, A.** 1977. An experiment in removal of organically polluted bottom mud from the Seto Inland Sea. *U.S. Environ. Prot. Agency, Off. Res. Dev., EPA-600/3-77-083*.
- Murakami, M., Oonishi, Y., Kunishi, H.** 1985. A numerical stimulation of the distribution of water temperature and salinity in the Seto Inland Sea (Japan). *J. Oceanogr. Soc. Jpn* 41 (4): 213-224.

- Murphy, J.P., Lippør, R.I. 1970. BOD production of channel catfish. The Progressive Fish-Culturist, 195-198 pp.
- Murphy, A.M., Grohmann, G.S., Christopher, P.J., Lopez, W.A., Davey, G.R., Millsom, R.H. 1979. An Australian-wide outbreak of gastroenteritis from oysters caused by *Norwalk virus*. Med. J. Aust. 2(7): 329-333.
- Murray, I, Parsons, J.W., Robinson, K. 1975. Inter-relationships between nitrogen balance, pH and dissolved oxygen in an oxidation ditch treating farm animal waste. Water Res. 9 (1): 25-30.
- Niemelä, M., Taipalinen, I. 1982. Faecal indicator bacteria at fish farms. Hydrobiologia 86: 171-175.
- Nishimura, A. 1982. Effects of organic matters produced in fish farms on the growth of red tide algae *Gymnodinium* type'65 and *Chattonella antiqua*. Bull. Plankt. Soc. Japan 29(1): 1-7.
- Nishimura, A. 1983. The process of decomposition and sedimentation of deposits in the pearl oyster and the yellowtail culture grounds. Bull. Jap. Soc. Fish. Oceanogr. Suisan Kaiyo Kenkyukaiho 44: 44-50. [In Japanese].
- Nishimura, A., Seki, M. 1983. Effects of lime for the improvement of mariculture grounds. [Yoshoku gyojo kairyo ni okeru sekkai no sayo.] Bull. Jap. Soc. Sci. Fish. 49(3): 353-358. [In Japanese].
- Nixon, S.W., Ovitat, C.A., Rogers, C. Taylor, K. 1971. Mass and metabolism of a mussel bed. Oecologia 8: 21-30.
- Nowicki, B.L., Nixon, S.W. 1985. Benthic nutrient remineralization in a coastal lagoon ecosystem. pp. 182-189.
- Odum, W.E. 1974. Potential effects of aquaculture on inshore coastal waters. Environ. Cons. 1(3): 225-230.
- Ohwada, K, Tanaka, N., Sugiyama, M. 1983. Penetration of microorganisms into deposited fish scales in the sediment of a coastal finfish culture ground. [Gyorui yoshokujo teido ai taiseki-shita gyorin eno biseibutsu no senko]. Bull. Jap. Soc. Sci. Fish. 49(4): 657. [In Japanese].
- Ohwada, K., Uemoto, H. Environmental conditions and some problems in pearl oyster culture grounds, a review. pp. 1-7.
- Ojanen, T., Salonen, S. 1986. Licencing requirements of fish farms and control under the Water Rights Act. Can Transl. Fish. Aquat. Sci. No. 5276: 1-3. (Original Title: Kalankasvatuslaitosten luvantarve ja vesilain mukainen valvonta. Suomen kalankasvattaja 4:16, 1984). [Original in Finnish].
- Okarchl, T. 1983. Red tides and fisheries damages. Eisei Kagaku 29: 1-4. [In Japanese].
- Olah, J., Sinha, V.P.R., Ayyappan, S., Purushothaman, C.S., Radheysayam, S. 1986. Community metabolism in tropical undrainable rural fish ponds. Int. Revue ges. Hydrobiol. 71(6): 807-812.
- Ottmann, F., Sornln, J.-M. 1985. Observations on sediment accumulation as a result of mollusk culture systems in France. Proc. Int. Symp. "Utilization of Coastal Ecosystems: Planning, Pollution and Productivity, (1982: Nov. 21-27: Rio Grande, RS, Brazil) Vol. 1: 329-337.
- Ovlatt, C.A., Gall, A.L., Nixon, S.W. 1972. Environmental effects of Atlantic Menhaden on surrounding waters. Ches. Sci. 13(4): 321-322.
- Page, J.W., Andrews, J.W. 1974. Chemical composition of effluent from high density culture of channel catfish. Water, Air, Soil, Pollution 3: 365-369.
- Pärjälä, E. et al. 1984. Pollution loadings from Finnish inland fish-farms: a case study. Aqua Fennica 14(2): 205-214.

- Parjala, E.** 1986. Reduction of the loading effects of fish farming - a literature review. Can. Transl. Fish. Aquat. Sci. No. 5252: 1-17. (Original Title: Kuormituksen vähentäminen kalanviljelyssä, kirjallisuuskatsaus. Vesitalous 5: 11-16, 1984). [Original in Finnish].
- Parsons, T.R., Dovey, H.M., Cochlan, W.P., Perry, R.I., Creon, P.B.** 1984. Frontal zone analysis at the mouth of a fjord - Jervis Inlet, British Columbia. Sarsia 69:133-137.
- Parsons, T.R., Stronach, J., Borstad, G.A., Loutitt, G., Perry, R.I.** 1981. Biological fronts in the Strait of Georgia, British Columbia, and their relation to recent measurements of primary productivity. Mar. Ecol. Prog. Ser. 6:237-242.
- Partanen, P.** 1986. A study of the zoobenthos in the environment of fish farms in the sea off Sipoo. Can. Transl. Fish. Aquat. Sci. 5267: 1-24. (Original Title: Pohjaeläintutkimus Sipoon edustan merialueen kalankasvatustilastoisten ympäristössä. Vesihallituksen monistesarja 297: 1-30, 1984).
- Paul, J.D., Davies, I.M.** 1986a. Effects of copper and tin-based anti-fouling compounds on the growth of scallops (*Pecten maximus*) and oysters (*Crassostrea gigas*). ICES, CM 1986/F:13, 1-13. (Mariculture Committee).
- Paul, J.D., Davies, I.M.** 1986b. Effects of copper and tin-based anti-fouling compounds on the growth of scallops (*Pecten maximus*) and oysters (*Crassostrea gigas*). Aquaculture 54: 191-203.
- Pearson, T.R., Rosenberg, R.** 1978. Macrobenthic succession in relation to organic enrichment and pollution of the marine environment. Oceanogr. Mar. Biol. Ann. Rev. 16: 229-311.
- Pearson, T.H., Stanley, S.O.** 1979. Comparative measurement of the redox potential of marine sediments as a rapid means of assessing the effect of organic pollution. Mar. Biol. 53: 371-379.
- Pease, B.C.** 1977. The effect of organic enrichment from a salmon mariculture facility on the water quality and benthic community of Henderson Inlet, Washington. Ph.D. thesis, Univ. Washington, Seattle, Washington.
- Penczak, T., Gallcka, W., Molinski, M., Kusto, E., Zalewski, M.** 1982. The enrichment of a mesotrophic lake by carbon, phosphorus and nitrogen from the cage aquaculture of rainbow trout, *Salmo gairdneri*. Journal of Applied Ecology 19: 371-393.
- Pettersson, K.** 1986. Betydelsen av fiskfoders fosforsammansättning för fosforläckage till vatten från foderspill och fekaler. (Importance of phosphorus composition in fish feeds for the leaching of phosphorus from wasted feed and faeces.). Slutrapport, LIU B: 18. [In Swedish].
- Phillips, M.J., Beveridge, M.C.M., Mulr, J.F.** 1985. Waste output and environmental effects of rainbow trout cage culture. ICES, C.M. 1985/F:21, 16 pp. (Mariculture Committee).
- Phillips, M.J., Beveridge, M.C.M., Stewart, J.A.** 1985. The environmental impact of cage culture on Scottish freshwaters. Water Research Conference, 1985, Poster Paper, 6 pp.
- Piedrahita, R.H., Brune, D.E., Tchobanoglous, G., Orlob, G.T.** 1984. A general model of the aquaculture pond ecosystem. J. World Maricul. Soc. 15: 355-366.
- Pingree, P.D.** 1978. Mixing and stabilization of phytoplankton distributions on the northwest European continental shelf. pp. 181-220. In J.H. Steele (ed.), Spatial Pattern in Plankton Communities. Plenum Press, New York.
- Pingree, P.D., Holligen, P.M., Mardell, G.T.** 1978. The effects of vertical stability on phytoplankton distributions in the summer on the northwest European shelf. Deep Sea Res. 25: 1011-1028.
- Plumb, J.A.** 1984. Relationship of water quality and infectious diseases in cultured channel catfish. Symposia Biologica Hungarica 23: 189-199.

- Poemomo, A., Singh, V.P.** 1982. Problems, field identification and practical solutions of acid sulfate soils for brackishwater fishponds. In: Report of consultation /seminar on coastal fishpond engineering, 4-12 Aug. 1982, Surabaya, Indonesia. FAO/UNDP, Manila (Philippines) SCS/GEN /82/42: pp. 49-61.
- Porras, D.** 1981. On the utilization of organic fertilizers (wastes and dung) in Aquaculture. (Sobre la utilización en acuicultura de fertilizantes orgánicos (deshechos y excretas)). Rev. Latinoam. Acuicult. 9: 6-10. [In Spanish].
- Purdom, C.E.** 1982. Analyses of influent and effluent water in UK trout farms - An Interim Report. EIFAC Tech. Paper 41: 83-86.
- Quayle, D.B.** 1969. Paralytic shellfish poisoning in British Columbia. Bull. Fish. Res. Board Can. 168: 1-69.
- Quereilou, J. Fauré, A., Fauré, C.** 1982. Pollution loads from rainbow trout farms in Brittany, France. EIFAC Tech. Paper 41: 87-97.
- Ramsey, H.** 1978. Fish hatchery investigations. 2. Fish hatchery effluent monitoring. IDFG, Boise, ID (USA), IDFG-Proj.-F-72R-2(2): 1-9 (Job Performance Rep. Idaho Dep. Fish Game).
- Roberts, R.J.** 1976. Bacterial diseases of farmed fishes. In: Microbiology in Agriculture, Fisheries and Food. (Eds. F.A. Skinner and J.G. Carr) The Society for Applied Bacteriology Symposium Series No. 4. (Academic Press) 55-62.
- Roberts, R.J., Bullock, A.M., Turner, M.F., Jones, K.J. Tøtt, P.** 1983. Mortalities of *Salmo gairdneri* exposed to cultures of *Gymnodinium aureolum*. J. Mar. Biol. Assoc. UK 63: 741-743.
- Robohm, R.A., Brown, C., Murchelano, R.A.** 1979. Comparison of antibodies in marine fish from clean and polluted waters of the New York Bight: relative levels against 36 bacteria. Appl. environm. Microbiol. 38: 248-257.
- Romero, P., González-Gurrlarén, E., Penas, E.** 1982. Influence of mussel rafts on spatial and seasonal abundance of crabs in the Ria de Arousa, North-west Spain. Mar. Biol. 72: 201-210.
- Rosenberg, R., Loo, L.O.** 1983. Energy-flow in a *Mytilus edulis* culture in western Sweden. Aquaculture 35(2): 151-161.
- Rosenthal, H.** 1980. Implications of transplantations to aquaculture and ecosystems. Mar. Fish. Rev. 42 (5): 1-14.
- Rosenthal, H.** 1983. Summary of the EIFAC Workshop document on "Fish Farm-Effluents". ICES, C.M. 1983/E.21, 1-15. (Mariculture Committee).
- Rosenthal, H.** 1983. Coastal aquaculture and water quality in protected bays of Japan. ICES, C.M. 1983/E: 22, 1-22. (Mariculture Committee).
- Rosenthal, H.** 1985a. Recent observations on environmental effects of large-scale net cage culture in Japanese coastal waters. ICES, C.M. 1985/F:34, 1-21 (Mariculture Committee).
- Rosenthal, H.** 1985b. Constraints and perspectives in aquaculture development. GeoJournal 10(3): 305-324.
- Rosenthal, H., Hansen, P.-D., Bergmann, H.** 1981. Accumulation of trace contaminants in fishes raised in recirculation systems. ICES/CM 1981/F:16, 1-8. (Mariculture Committee).
- Russels, C.S.** (Ed.). 1975. Ecological modeling in a resources management framework. John Hopkins Univ. Press, Baltimore, Maryland, USA.
- Ryther, J., Losordo, T.M., Furr, A.K., Parkinson, T.F., Guteman, W.H., Pakkala, I.S., Lisk, D.J.** 1979. Concentration of elements in marine organisms cultured in seawater flowing through coal-fly ash. Bull. Environm. Contam. Toxicol. 23: 207-210.

- Rzepecka, H., Kunert, A., Reczek, B.** 1974. Contamination of seawater in port docks with Enterobacteriaceae strains. Roczn. Panstw. Zaky. Hig., 25 (2): 237-243.
- Salto, T., Maruyama, J., Kanoh, S., Jeon, J.K., Noguchi, T., Harada, T., Murata, O., Hashimoto, K.** 1984. Toxicity of the cultured pufferfish *Fugu rubripes rubripes* along with their resistibility against tetrodotoxin. Nippon Suisan Gakkaishi 50(9): 1573-1575. [In Japanese].
- Salto, T., Noguchi, T., Takeuchi, T. Kammura, S., Hashimoto, K.** 1984. Ichthyotoxicity of paralytic shellfish poison. Bull. Jpn. Soc. Sci. Fish. 51 (2): 257-260.
- Sakamoto, I.** 1977. Estimating the carrying capacity for culturing fish". Senkai Yoshoku to Jika Osen [Shallow Sea Aquaculture and Self-Pollution]. Suisan-gaku Shirizu 21 [Fisheries Series 21]. Edited by the Japanese Society of Scientific Fisheries. Published by Koseisha Koseikaku 134 pp [In Japanese].
- Salonen, S., Ojanen, T.** 1986. Some aspects of licencing conditions applying to licence granted to fish farms by the Water Rights Court. Can. Transl. Fish. Aquat. Sci. No. 5275: 1-4. (Original Title: Nakokotia kalankasvatus- laitoksille annettujen vesioikeuden paatosten lupachdoista. Suomen kalankasvattaja 4:24, 1984. (Original in Finnish).
- Salonen, S.** 1983: Kuormituksen määrittämiseen käytettyjen menetelmien arviointi. (Evaluating of the methods to assess the loading.). Mimeo, 9 pp. [In Finnish].
- Salonen, S.** 1986. Kalankasvatuksen vesilakiin perustuva sääätely. (Regulation of fish farming based on the Water Law.). INSKO 282-86 I. 9 pp. [In Finnish].
- Sayler, G.S., Nelson, J.D.Jr., Colwell, R.R.** 1975. Role of bacteria in bioaccumulation of mercury in the oyster *Crassostrea virginica*. Appl. Microbiol 30 (1): 91-96.
- Scherf, E., Wassermann, M.** 1971. Neue Ergebnisse über die Wasserverdunstungshemmung durch monomolekulare Fettkoholschichten. (New results on the reduction of evaporation losses through the use of mono- molecular surface films.). Vom Wasser 38: 207-214. [In German].
- Schnick, R.A., Meyer, F.P., Walsh, D.F.** 1986. Status of fishery chemicals in 1985. Progr. Fish-Cult. 48(1): 1-17.
- Schomburg-Barrett, S.L.** 1982. The stability of animal feed antibiotics in surface waters. M.S. thesis, Rutgers State University, New Brunswick, New Jersey.
- Schroeder, G.L.** 1975. Some effects of stocking fish in waste treatment ponds. Water Research 9: 591-593.
- Sellgam, P.F., Valkirs, A.O., Lee, R.F.** 1986. Degradation of tributyltin in marine and estuarine waters. Oceans'86 Conference Record, Organotin Symposium, Vol.4: 1189-1195.
- Sheng, Y.P.** 1983. Mathematical modeling of three-dimensional coastal currents and sediment dispersion: model development and application. U.S. Army Corps of Engineers, Washington, D.C., Tech. Rep. CERC-83-2.
- Shepherd, B.G.** 1984. The biological design process used in the development of Federal Government Facilities during phase 1 of the salmonid enhancement program. Can. Tech. Rep. Fish. Aquat. Sci. 1275: 1-53.
- Short, J.W., Thrower, F.P.** 1986a. Tri-n-butyltin caused mortality of chinook salmon, *Oncorhynchus tshawytscha*, on transfer to a TBT-treated marine net pen. Oceans'86 Conference Record, Organotin Symposium, Vol.4: 1202-1205.
- Short, J.W., Thrower, F.P.** 1986b. Accumulation of butyltins in muscle tissue of chinook salmon reared in sea pens treated with tri-n-butyltin. Oceans'86 Conference Record, Organotin Symposium, Vol.4: 1177-1181.
- Shumway, D.L., Palensky, J.R.** 1973. Impairment of the flavor of fish by water pollutants. EPA Ecol. Res. Ser., EPA-R3-73-010, 80 pp.

- Simensen, T., Munthe-Kaas, H.** 1973. Development of numerical fjordmodels for the functional planning of a regional waste water disposal system. In: Jenkins, S.H. (Ed.) "Advances in water pollution research. Proc. 6th Int. Conf, Jerusalem, June 18-23, 1972. Pergamon Press Ltd, Oxford, UK, ISBN - 0-08- 017005-6., pp. 853-863.
- Sindermann, C.J.** 1977. Disease problems created by introduced species. In C.J. Sindermann (ed.), Disease Diagnosis and Control in North American Marine Aquaculture. pp. 309-313. Developments in Aquaculture and Fisheries Science, 6. Elsevier Scientific Publ. Co., Amsterdam.
- Sindermann, C.J.** 1984. Diseases in marine aquaculture. Helgoländer Meeresunters. 37: 505-532.
- Singhal, R.N., Jeet, S., Davies, R.W.** 1986. Chemotherapy of six ectoparasitic diseases of cultured fish. Aquaculture 54: 165-171.
- Smith, E.J.** 1985. Marine resources- information leaflet: regulations governing marine aquaculture. Prepared by the California Department of Fish and Game, Sacramento, California.
- Sneed, K.E.** 1971. Pesticides residues in catfish. Resour. Publ. U.S. Bur. Sport Fish. Wildl. 102: 193-196.
- Solbé, L.G.** 1982. Fish-farm effluents; A United Kingdom survey. EIFAC Tech. Paper 41: 29-55.
- Solberg, S.O., Bregnballe, F.** 1982. Pollution from farmed trout feed with minced trash fish. EIFAC Tech. Paper 41: 65-71.
- Solomons, I.A.** 1978. Antibiotics in animal feeds - human and animal safety issues. J. Anim. Sci. 46(5): 1360-1368.
- Sommer, D.A., Stulber, D.A., Bradley, R.L., Peterson, R.E.** 1982. Raising marketable yellow perch on an polychlorinated biphenyl contaminated diet: A feasibility study for the perch aquaculture industry. Arch. Environ. Contam. Toxicol. 11(5): 589-593.
- Sornin, J.M.** 1979. Enquete sur la sédimentation et l'exhaussement des fonds dans les zones conchylicoles des cotes de France. Rapprt DEA, Université de Nantes, 41 pp.
- Sornin, J.M.** 1981. Sedimentary process and biodeposition linked to different shellfish culture methods. Lab. de Geologie Marine et Appliquee 1981, 188pp.
- Sornin, J.M., Feuillet, M., Héral, M., Deslous-Paoli, J.M.** 1983. Effet des biodépôts de l'huitre *Crassostrea gigas* sur l'accumulation de matière organiques dans les parcs du bassin de Marennes-Oléron. Journal of Molluscan Study, Suupt. 12A: 157-197.
- Sornin, J.M., Feuillet, M., Héral, M., Deslous-Paoli, J.M.** 1986. Influence des cultures d'huitres *Crassostrea gigas* sur le cycle du phosphore en zone intertidale: role de la biodéposition. Oceanol. Acta 9(3): (in press).
- Sparrow, R.A.H.** 1981. Hatchery effluent water quality in British Columbia. In: Proceedings of the bio-engineering symposium for fish culture (Eds.: L.J. Allen and E.C. Kinney) American Fisheries Society pp. 162-166.
- Speece, R.E.** 1973. Trout metabolism characteristics and the rational design of nitrification facilities for water reuse in hatcheries. Trans. Am. Fish. Soc. 102(2): 323-334.
- Stang, P.M., Sellgman, P.F.** 1986. Distribution and fate of butyltin compounds in the sediment of San Diego Bay, California. pp. 1256-1261. In Oceans 86 Conference Record, Washington, D.C., September 23-25, 1986.
- Stebbing, A.R.D.** 1985. Organotins and water quality - some lessons to be learned. Mar. Pollut. Bull. 16: 383-390.

- Stephenson, M.D., Smith, D.R., Goetzi, J., Ichikawa, G., Martin, M.** 1986. Growth abnormalities in mussels and oysters from areas with high levels of tributyltin in San Diego Bay. Oceans'86 Conference Record, Organotin Symposium, Vol.4: 1246-1251.
- Stewart, K.I.** 1984. A study on the environmental impact of fish cage culture on an enclosed sea Loch. MSc. Thesis, University of Stirling, Scotland.
- Stirling, H.P., Beveridge, M., Ross, L.G., Phillips, M.J.** 1985. Chemical and biological methods of water analysis for aquaculturists. Institut of Aquaculture, Univ. Stirling, Scotland. 119 pp.
- Storstrøms amtskommune,** 1985. Notat om havbrug. Miljøkontoret, Storstrøms amtskommune pp. 1-28.
- Storstrøms amtskommune,** 1987 (in preparation). Tilsyn med havbrug, 1986. Miljøkontoret, Storstrøms amtskommune publication.
- Stössel, P.** 1986. Aflatoxin Contamination in Soybeans: Role of proteinase inhibitors, zinc availability, and seed coat integrity. Appl. Environm. Microbiol. 52(1): 68-72.
- Sumari, O.** 1982. A report of fish farm effluents in Finland. EIFAC Tech. Paper 41: 21-27.
- Sumari, O.** 1984. Feed and waste water. the International Conference Acquacultura'84. pp1-16.
- Sutherland, I.R., Roland, W.G.** 1986. Monitoring the development of intense oyster culture in Trevensen Bay. B.C. Aquaculture Newsletter 6(4):17-24.
- Suzuki, S., Sato, N., Takatsuki, K., Kikuchi, H., Ushizawa, I.** 1985. Pollution of corbiculae with chlorinated dibenzo-p-dioxins from a herbicide, chlornitrofen: residues of paddy herbicides in fish and shellfish. J. Food Hyg. Soc. Jpn 26(2): 137-143. [In Japanese].
- Swärdh, L., Thulin, J.** 1985. The parasite fauna of natural and farmed *Mytilus edulis* from the west coast of Sweden, with special reference to *Renicola roscovita*. Med. Havsfisk. Lysekil 312: 1-16.
- Swedish Steering Committee on Aquaculture** 1983. The environmental impact of aquaculture. Report from the working group on environmental effects to the Steering Committee on Aquaculture, Swedish Council for Planning and Coordination of Research in cooperation with the National Marine Resources Commission, Forskningsrådsnämnden, 74 pp.
- Swingle, H.S.** 1968. Fish kills caused by phytoplankton blooms and their prevention. FAO Fish. Rep. No. 44, Vol.5: 407-411.
- Tabira, N.** 1980. the fish culture under the optimum density and the fishery right administered by fishermen's cooperative associations: A juristic study of yellowtail culture. Mem. Fac. Fish., Kagoshima Univ. 29: 157-167.
- Takagi, M., Kamada, H., Tazaki, A.** 1980. A consideration on the occurrence of Protogonyaulax sp. at the scallop ground in Funka Bay. Bull. Fac. Fish. Hokkaido Univ. 31(2): 215-221. [In Japanese].
- Takahashi, M., Fukazawa, N.** 1982. A mechanism of "red-tide" formation. II. Effect of selective nutrient stimulation on the growth of different phytoplankton species in natural water. Mar. Biol. 70: 267-273.
- Takamiya, K.** 1985. Organochlorine residues in human tissues and some foods, and the role of yellowtails as an index of the residues. Nippon Eiseigaku Zasshi 40(5): 822-824. [In Japanese].
- Tangen, G.** 1977. Blooms of *Gyrodinium aureolum* (Dinophyceae) in north European waters, accompanied by mortality in marine organisms. Sarsia 63(2): 123-133.
- Tarfeno-Silva, E., Kawasaki, L.Y., Yu, D.P., Gordon, M.S., Chapman, D.J.** 1982a. Aquacultural approaches to recycling of dissolved nutrients in secondarily treated domestic wastewaters - II. Biological productivity of artificial food chains. Water Research 16: 51-57.

- Tarifeño-Silva, E., Kawasaki, L.Y., Yu, D.P., Gordon, M.S., Chapman, D.J. 1982b. Aquacultural approaches to recycling of dissolved nutrients in secondarily treated domestic wastewater - III. Uptake of dissolved heavy metals by artificial food chains. *Water Research* 16: 59-65.
- Tarnbuppa, P. 1980. Snail eradication in shrimp farm. *Thai Fish. Gaz.* 33(3): 305-307.
- Tenore, K.R., Gonzalez, N. 1975. Food chain patterns in the Ria de Arosa, Spain: An area of intensive mussel aquaculture. 10th European Symposium on Marine Biology, Ostend, Belgium, Vol. 2 Universa Press, Wetteren, pp. 601-619.
- Tenore, K.R., Boyer, L.F., Cal, R.M., Corral, J., Garcia-Fernandez, C., González, N., González-Gurriaran, E., Hanson, R.B., Iglesias, J., Krom, M., López-Jamur, E., McClain, J., Pamatmat, M.M., Perez, A., Rhoads, D.C., deSantiago, G., Tietjen, J., Westrich, J., Windom, H.L. 1982. Coastal upwelling in the Rias Bajas, NW Spain: contrasting the benthic regimes of the Rias de Arosa and de Muros. *J. Mar. Res.* 43: 701-772.
- Tenore, K.R., Corral, J., Gonzalez, N. 1985. Effects of intensive mussel culture on food chain patterns and production in coastal Galicia, NW Spain. Proc. SIUEC (Symposium international on Utilization of Coastal Ecosystems, Rio Grande, RS, Brazil), Vol 1: 321-328.
- Terashima, S., Takagi, M., Fujisawa, K. 1972. Survey on the condition of sea water and bottom mud in the culture grounds of marine bivalves, arc-shell *Anadara (scapharca) broughtnis* (Schrenck) and *Anadara (scapharca) subcrenata* (Lischke). *Bull. Fish. Expt. Stat. Okayama Prefect.* 46: 120-153. [In Japanese].
- Thain, J.E. 1986. Toxicity to TBT to bivalves: Effects on reproduction, growth and survival. Oceans'86 Conference Record, Organotin Symposium, Vol.4: 1306-1313.
- Thompson, J.A.J., Sheffer, M.G., Pierce, R.C., Chau, Y.K., Cooney, J.J., Cullen, W.R., Maguire, R.J. 1985. Organotin compounds in the aquatic environment: scientific criteria for assessing their effects on environmental quality. National Research Council of Canada, NRCC Associate Committee on Scientific Criteria for Environmental Quality, NRCC No 22494: 1-284.
- Thorslund, A. 1980. Water pollution from trout cage-culture. *Vatten* 4: 308-311.
- Ti, T.L., Hassan, R.B., Rajamanickam, L.D. 1982. Coastal mangrove swamps for fishponds: A focus on some environmental factors. In: Report of consultation/seminar on coastal fishpond engineering, 4-12 Aug. 1982, Surabaya, Indonesia. FAO/UNDP Manila (Philippines) SCS/GEN/82/42: pp. 62-66.
- Trojanowski, J., Trojanowska, C., Ratajczyk, H. 1982. Effect of intensive trout culture in lake letowo on its bottom sediments. *Pol. Arch. Hydrobiol.* 29(3-4): 659-670.
- Tucholski, S., Kok, J., Wojno, T. 1980. Studies on removal of wastes produced during cage rearing of rainbow trout (*Salmo gairdneri*, Richardson) in lakes. 1. Chemical composition of wastes. *Rocz. Nauk. Roln.* 82: 3-15.
- Tucker, C.S., Busch, R.L., Lloyd, W.W. 1983. Effects of simazine treatment on channel catfish production and water quality in ponds. *J. Aquat. Plant Manage.*, vol. 21: 7-11.
- Turun vesipiiririin vesitoimisto 1984. Ehdotus saaristomeren kalankasvatustoimintaa koskevaksi vesiensuojelusuunnitelmaksiksi. (A proposal for the plan of water conservation concerning fish farming activities in the Archipelago sea.) Mimeo 236. [In Finnish].
- Uyeno, F., Funahashi, S., Tsuda, A. 1970. Preliminary studies on the relation between faeces of pearl oyster (*Pinctata martensi* (Dunker)) and bottom conditions in an estuarine pearl oyster area. *J. Fac. Fish., Prefect. Univ. Mie* 8(2): 113-137. [In Japanese].
- Van Houwelling, C.D., Galner, J.H. 1978. Public health concerns relative to the use of subtherapeutic levels of antibiotics in animal feeds. *J. Anim. Sci.* 46(5): 1413-1424.
- Vaughn, J.M., Metcalf, T.G. 1975. Coliphages as indicators of enteric viruses in shellfish and shellfish raising estuarine waters. *Water Res.* 9: 613-616.

- Venkataramlah, A., Lakshmi, G.J., Best, C., Gunter, G., Hartwig, E., Wilde, P. 1982. Studies on toxicity of OTEC plant components on *Eucalanus* sp. from the Gulf of Mexico. Ocean Science and Engineering 7 (3): 353-401.
- Verwey, J. 1952. On the ecology of distribution of cockle and mussel in the Dutch Waddensea, their role in sedimentation and the source of food supply, with a short review of the feeding behaviour of bivalve mollusks. Arch. Neerl. Zool. 10: 172-239.
- VKI (Vandkvalitetsinst. & Jydsk Teknologisk Inst.). 1976. Forskellige driftsparameters indflytelse på foreningen fra dambrug. (The influence of different operational parameters on the pollution from fish farms). Rapport til Teknologiradet. Hørsholm, Denmark. [In Danish].
- Walker, D.B., Kay, B.H. 1983. Shellfish growing water bacteriological and sanitary survey of Saltspring Island, British Columbia, 1982. Environment Canada, Environmental Protection Service, Pacific Region, Regional Program Report 83-22, 66 pp.
- Walker, D.B., Kay, B.H. 1986. Shellfish growing water bacteriological and sanitary survey of Sooke Harbour and Basin, British Columbia, 1983. Environment Canada, Conservation and Protection, Environmental Protection Service, Pacific and Yukon Region, Regional Program Report 84-24, 69 pp.
- Warrer-Hansen, I. 1982. Evaluation of matter discharged from trout farming in Denmark. EIFAC Tech. Paper 41: 57-63.
- Warrer-Hansen, I. 1982. Methods of treatment of waste water from trout farming. EIFAC Tech. Paper 41: 113-121.
- WDF and WDOE. 1985. Use of the insecticide Sevin to control ghost and mud shrimp in oyster beds of Willapa Bay and Grays Harbor. Final Environmental Impact Statement. Prepared by the Washington Department of Fisheries and the Washington Department of Ecology.
- Weston, D.P. 1986. The environmental effects of floating mariculture in puget sound. School of Oceanography, College of Oceans and Fishery Sci., Univ. Washington, Seattle, USA, 148 pp.
- White, A.W. 1977. Dinoflagellate toxins as probable cause of an Atlantic herring (*Clupea harengus harengus*) kill, and pteropods as apparent vectors. J. Fish Res. Board Can. 34: 2421-2424.
- White, A.W. 1981a. Sensitivity of marine fishes to toxins from the red tide dinoflagellate *Gonyaulax excavata* and implications for fish kills. Mar. Biol. 65: 255-260.
- White, A.W. 1981b. Marine zooplankton can accumulate and retain dinoflagellate toxins and cause fish kills. Limnol. Oceanogr. 26(1): 103-109.
- White, A.W. 1982. Intensification of *Gonyaulax* blooms and shellfish toxicity in the Bay of Fundy. Can. Tech. Rep. Fish. Aquat. Sci. 1064: 1-12.
- White, A.W., Lewis, C.M. 1982. Resting cysts of the toxic red tide dinoflagellate *Gonyaulax excavata* in the Bay of Fundy sediments. Can J. Fish. Aquat. Sci. 39: 1185-1194.
- Wlugh-Mäsak, P. 1982. Season-dependent dynamics in mussel cultures in Mollösund and Lyresund, Bohuslän, Sweden. Med. Havslisk. Lysekil 284: 1-26.
- Wildish, D.J., Kristmanson, D.D. 1984. Importance to mussels of the benthic boundary layer. Can. J. Fish. Aquat. Sci. 41: 1618-1625.
- Wildish, D.J., Martin, J.D., Wilson, A.J., DeCoste, A.M. 1986. Hydrographic and sedimentary conditions in the L'Etang Inlet during 1985. Canadian Technical Report of Fisheries and Aquatic Sciences, pp 1-34.
- Willoughby, H., Larsen, H.N., Bowen, J.T. 1972. The pollutional effects of fish hatcheries. Amer. Fish. U.S. Trout News 17 (3): 6-7, 20-21.
- Wooten, R. 1979. Tapeworm threat to trout in floating freshwater cages. Fish Farmer 2(3): 5.

Wootton, R. 1985. Experience of sea lice infestations in scottish salmon farms. ICES, C.M. 1985 /F:7, 1-6. Henderson, E.B. Guidelines for physical separation of salmonid aquaculture farms.

Working Group Report 1986. A plan for a water protection programme for the sea off Kuivaniemi with reference to fish farming. Working Group Proposal. Can. Transl. Fish. Aquat. Sci. No. 5271: 1-86. (Original Title: Kuivaniemen edustan merialueen kalankasvatusta koskeva vesiensuojelusuunnitelma. Työryhmän ehdotus. Vesihallituksen monistessarja 298: 1-65, 1984).

Zanker, K. 1961. Verminderung der Verdunstungsverluste auf Seen und Talsperren. (Reducing evaporation losses on lakes and reservoirs.). Wasser Boden 13: 6-12. [In German].

Zleman, H., Grosswendt, U. 1985. Untersuchungen über die in industriemäßigen Anlagen der Forellenmast in Betonrinnen anfallenden Abwasserinhaltsstoffe. (Investigations on waste water substances derived from industrial high intensity trout fattening in raceways.). Z. Binnenfisch. DDR :14-19. [In German].

Zlaskowski, J., Murchelano, R. 1975. Fin erosion in winter flounder (*Pseudopleuronectes americanus*) from the New York Bight. Mar. Poll. Bull. 6: 26-28.

Zitko, V. 1986. Chemical contamination in aquaculture. Canadian Aquaculture 2(1): 9-10.

