


Handwritten scribbles

FISKERIDIREKTORATETS SKRIFTER

SERIE HAVUNDERSØKELSER

VOL. 18. No. 5



DIRECTORATE OF FISHERIES, BERGEN, NORWAY

1987

THE GREY SEAL, *HALICHOERUS GRYPUS* (FABRICIUS), IN FINNMARK, NORWAY

By

ØYSTEIN WIIG

Institute of Marine Research, Bergen, Norway

ABSTRACT

WIIG, Ø. 1987. The grey seal, *Halichoerus grypus* (Fabricius), in Finnmark, Norway. *FiskDir. Skr.Ser. HavUnders.*, 18: 241-246.

The distribution, abundance and breeding season of grey seals in Finnmark, Norway are reviewed. The seals are divided into eight hypothetical stocks: Sørøya, Magerøya, Laksfjord, Tanafjord, Kongsfjord, Syltefjord, Vardø and Varangerfjord. Breeding has been verified for five of these stocks. The total minimum observed stock size is 353 grey seals, but the real number is probably much higher. The breeding season seems to be in October-November west of Nordkapp and in December in the eastern part of the county.

INTRODUCTION

The grey seal, *Halichoerus grypus*, and the common seal, *Phoca vitulina*, are the only resident seals on the coast of the Norwegian mainland. The grey seals seem to prefer the outlying skerries for their breeding while the common seals select more sheltered areas.

ØYNES (1964) recorded 660 pups of grey seals in 1963 from Møre to Finnmark, but no pup production was recorded from southern Norway, although the grey seal has been observed at several localities also in that part of the country. Based on the figures of ØYNES (1964, 1966), SUMMERS *et al.* (1978) estimated the total population in Norway to 2000-3000 grey seals.

A study of coastal seals and their interactions with inshore fisheries along the Norwegian coast from Stad (about 62°N) to Lofoten (about 68°N) was initiated by the Institute of Marine Research in 1974. The study has later been extended to the entire Norwegian coast. One result of this study is new information on the distribution and abundance of grey seals along the Norwegian coast which has been reviewed by WIIG (1986). The surveys indicate a minimum total stock of 3100 grey seals on the coast of Norway.

ØYNES (1964) reported two known colonies of grey seals in Finnmark county and estimated the number of breeding females to be three. Recent surveys show that the stock of grey seals in Finnmark is much higher.

METHODS

The number of seals has been estimated from aerial and ground surveys.

The aerial surveys were made from a high-wing, twin-engined Islander aircraft. In addition to direct counting, the seals were photographed by 35 mm hand-held reflex-cameras, using colour reversal film as recommended by VAUGHAN (1971). By this method large areas could be surveyed during short periods of time.

The ground surveys were made from 12'–15' Zodiac inflatable boats which are well suited for fast landings on rocky shores and skerries. The seals were counted by sight. By this method only a restricted area could be surveyed each day.

RESULTS

Seal counts by aerial and ground surveys on the coast of Finnmark are shown in Table 1 and the localities are shown on Fig. 1.

Two ground surveys lasted for several days, which could allow double observations of the same seals. However, distances between the localities are so large that such effect is not believed to have biased the results.

In addition to the observations given in Table 1, single seals have been observed at different localities along the coast between Grense-Jacobselv and Sørøya.

Proof of breeding has been reported from five areas. At Sørøya seven pups were tagged at Store- and Lille-Kamøya in 1982 (ØRITSLAND, 1982). At Skarholmen in Tanafjord two white pups were observed in January 1979. In Kongsfjord one white pup and five pregnant females were observed on 16 December 1982, and two pups (2–3 weeks old) were observed on 5 January 1984. At Syltefjorden one pup (about 2 weeks old) was observed on the eastern shores of Makkaurfjord on 5 January 1985. In addition, local hunters have reported taking pups in this area in previous years. The fifth locality is at Vardø, where one pup (about 4 weeks old) was tagged and another pup was observed in the sea on 10 January 1985.

It is worth noting from Table 1 that in the western part of the county pups have been observed in late October to early November, while the pups observed in the eastern part seem to have been born in December. Such late pupping in this area has also been reported from local sources (BERGFLØDT *et al.* 1985). Two white pups were observed on Skarholmen in Tanafjord on 15 January 1979 and two pups (2–3 weeks old) were observed on Skarholmen in Kongsfjord on 5 January 1984.




Table 1. Observations of grey seals in Finnmark and minimum local stock sizes.

Localities	Recorded numbers								Minimum stock size	
	June-July 1981 (Björge <i>et al.</i> 1982)	November 1981 (Bergflødt and Fagerheim 1981)	October 1982 (Øritsland 1982)	November 1982 (Øritsland 1982)	December 1982 (Fagerheim and Bergflødt 1982)	June-July 1983 (Øritsland 1983)	April 1984 (Øien 1984)	January 1985 (Bergflødt <i>et al.</i> 1985)		March 1985 (Bergflødt and Wiig 1985)
Bondøy S.-and L.-Kamøy Revsholmen Skipsholmen	70		1 ¹	6 ²		2 8	125			125
Gjesvær Tunes	5 3						40			40
Lebesbyholmene							50			50
Skarholmen in Tana								× ⁴		50
Kongsøy Skarholmen				10 ³		20		11	29	29
Syltefjord Makkaurfjord	40	30						5 ⁵	3	40
Reinøy/Hornøy Kibergsholmene		2				4	7 1	15 16 ⁷	3 ⁶	19
Total										353

¹ White pup and many adults² Newborn pups and many adults³ One white up, five pregnant females and four males⁴ Tracks from 50–75 seals⁵ One pup about 2 weeks old⁶ Two pups about 3 weeks old⁷ One pup about 3 weeks old

DISCUSSION

Our knowledge of grey seals in Finnmark is relatively limited. The main reason for this is the extreme weather conditions in this area during early winter from October to January when the seals are expected to breed.

ØYNES (1964) reported two breeding sites of grey seals in Finnmark. The northernmost site was situated just west of Nordkapp, where three pups were born in October 1962. The other breeding site reported by Øynes was at Bondøya outside Sørøya. However, the number of pups born at this site was unknown because no one had visited the island at that time of the year. According to the observations reported here, five more breeding localities of grey seals in Finnmark can be added to the two reported by ØYNES (1964). As a hypothesis it thus seems convenient to separate the grey seals in Finnmark into eight local stocks. These are those of Sørøya, Magerøya, Laksfjord, Tanafjord, Kongsfjord, Syltefjord, Vardø and Varangerfjord. Breeding has been verified for five of these stocks.

In order to get a rough estimate of the number of grey seals in these local stocks, a minimum size is estimated from the largest number of seals observed in each area (table 1). These numbers add up to a minimum stock size of 353 grey seals in Finnmark. There are many inaccuracies attached to these figures and the actual number of grey seals in the county is probably much higher. BJØRGE (unpubl.) estimated the number of grey seals in Finnmark in 1983 to be about 400.

The breeding season in Finnmark seems to differ between areas west and east of Nordkapp. At Sørøya pups have been observed in October and November, whereas on the coast from Tana to Vardø pups have been observed from December to January. Further south, between 62°N and 68°N where most of the Norwegian stock of grey seals is found, the breeding season is in September and October (BENJAMINSEN *et al.* 1977). Thus, there seems to be a cline in the date of breeding of grey seals along the Norwegian coast from 62°N to Vardø. The breeding season in Rogaland county is, however, also late, occurring in November–December (WIIG 1986), so that the cline does not extend to the southern part of the coast. The breeding season in eastern Finnmark also appears to be later than on the Murman coast which is in November (KARPOVICH *et al.* 1968). The reason for such late breeding in eastern Finnmark is not clear.

ACKNOWLEDGEMENTS

I am grateful to T. Øritsland for commenting on the manuscript.

REFERENCES

- BENJAMINSEN, T., BERGFLODT, B., HUSE, I., BRODIE, P. and TOKLUM, K., 1977. Undersøkelser av havert på norskekysten fra Lofoten til Frøya, september–november 1976. *Rapp. Fondet for fiskeleting og Forsøk, 1977 (1)*: 24–33.
- BERGFLODT, B. and FAGERHEIM, K.A., 1981. Rapport om leting etter havertunger med fly på Finnmarkskysten i oktober–november 1981. *Rapp. Fiskeridir. Havforskn. inst., Bergen*. 3 pp.
- BERGFLODT, B., FAGERHEIM, K.A. and WIIG, Ø., 1985. Selundersøkelser i Østfinnmark 3.–14. januar 1985. *Rapp. Fiskeridir. Havforskn. inst., Bergen*. 4 pp.
- BERGFLODT, B. and WIIG, Ø., 1985. Flyleting etter havert mellom Kirkenes og Hammerfest 5. mars 1985. *Rapp. Fiskeridir. Havforskn. inst., Bergen*. 2 pp.
- BJØRGE, A., BERGFLODT, B., FAGERHEIM, K.A., ØRITSLAND, T. and SKADSHEIM, A., 1982. Undersøkelser av steinkobbe og havert i Rogaland og Finnmark i 1981. *Fisken Hav., 1982(2)*: 1–9.
- FAGERHEIM, K.A. and BERGFLODT, B., 1983. Selundersøkelser i Østfinnmark 10.–17. desember 1982. *Rapp. Fiskeridir. Havforskn. inst., Bergen*. 3 pp.
- KARPOVICH, V.N., KOKHANOV, V.D. and TATARINKOVA, I.P., 1967. The grey seal on the Murman coast. *Fish. Res. Board Can., Trans. Ser., 1030*. 10 pp.
- SUMMERS, C.F., BONNER, W.N. and HAAFTEN, J.L. VAN. 1978. Changes in the seal population of the North Sea. *Rapp. P.-v. Reun. Cons. int. Explor. Mer., 172*: 278–285.
- VAUGHAN, R.W., 1971. Aerial survey of seals in the Wash. *Seals Res. Unit, Nat. Env. Res. Council (NEREC) occasional publ. 2*. 11 pp.
- WIIG, Ø., 1986. The status of grey seal (*Halichoerus grypus*) in Norway. *Biol. Conserv., 38*: 339–349.
- ØIEN, N., 1984. Selundersøkelser i Østfinnmark 10.–13. april 1984. *Rapp. Fiskeridir. Havforskn. inst., Bergen*. 4 pp.
- ØRITSLAND, T., 1982. Norsk fangst og undersøkelse av sel i 1982. *Rapp. Fiskeridir. Havforskn. inst., Bergen*. 23 pp.
- ØRITSLAND, T., 1983. Norsk fangst og undersøkelse av sel i 1983. *Rapp. Fiskeridir. Havforskn. inst., Bergen*. 22 pp.
- ØYNES, P., 1964. Sel på norskekysten fra Finnmark til Møre. *Fiskets Gang, 50*: 694–707
- ØYNES, P., 1966. Sel i Sør-Norge. *Fiskets Gang, 52*: 834–839.

Received 15. December 1985

Printed 16. February 1987

SEASONAL VARIATIONS OF THE OCEANOGRAPHIC CONDITIONS OFF THE SOUTHWEST COAST OF INDIA DURING 1971-1975

By

O. M. JOHANNESSEN

Geophysical Institute, University of Bergen, Norway

and

G. SUBBARAJU

Central Marine Fisheries Research Institute, Cochin, India

and

J. BLINDHEIM

Institute of Marine Research, Bergen, Norway

ABSTRACT

JOHANNESSEN, O.M., SUBARAJU, G. and BLINDHEIM, J. 1981. Seasonal variations of the oceanographic conditions off the southwest coast of India during 1971-1975. *FiskDir. Skr. Ser. HavUnders.*, 18: 247-161.

As part of a survey of the coastal fish resources off the southwest coast of India, the UNDP/FAO (United Nations Development Programme/Food and Agriculture Organization) Pelagic Fishery Project, Cochin, collected environmental data from June 1971 through October 1975. Based on these data, the oceanographic conditions during the 4-year period are discussed. The seasonal variations are highly repetitive from year to year. An uplift of water onto the shelf begins in March or April. Associated with a south-flowing current, this type of upwelling lasts throughout the SW monsoon period until September-October. A northflowing coastal current from November-December through January-February is associated with an influx of low-salinity water from the south. The environmental variations on the shelf are reflected in seasonal biological events.

INTRODUCTION

Several authors have reported on the environmental conditions along the southwest coast of India. In 1957 RAMA SASTRY and MYRLAND (1960) observed

upwelling on the shelf, while BANSE (1959) described associated low values in dissolved oxygen and discussed its effects on the catch of demersal fish. Major contributions from DARBYSHIRE (1967) and BANSE (1968) established that upwelling takes place during the SW monsoon period, May–October. Due to a minimum in the oxygen distribution at intermediate depth, water with less than 0.5 ml O₂/l comes up on the shelf during the upwelling period. The south-flowing current transports Arabian Sea Water of high salinity southward along the coast, though during the rainy season runoff causes a significant lowering of the surface salinity in the coastal zone. In the NE monsoon period the current reverses, advecting less saline Equatorial Surface Water from the equatorial region and Bay of Bengal northward (DARBYSHIRE 1967, WYRTKI 1973), causing a sinking or retreat of the Arabian Sea Water.

The present paper describes some features of the hydrography off the southwest coast of India, based on a programme carried out by the UNDP/FAO Pelagic Fishery Project in Cochin. An element of this programme was the study of the relationship between environmental factors and the abundance, distribution and migration of fish resources along the coast. For this purpose the shelf region between 7 and 17°N was surveyed 7–8 times per year from the middle of 1971 to October 1975, and hydrographic sections were worked across the shelf from Ratnagiri in the north to Tuticorin on the south coast (Fig. 1).

The project presented its results in progress reports which had only a limited distribution. In the oceanographic community there is, however, a general interest in the oceanographic effects associated with the monsoon, which calls for a wider distribution of these findings. The results presented here are based mainly on Progress Report no. 16 (ANON. 1976) and partly on Progress Reports nos. 3, 7, 12, 13 and 17 (ANON. 1973, ANON. 1974, ANON. 1976 a, ANON. 1976 b, ANON. 1976 c), and presents a brief overview of the findings.

OBSERVATIONS

A total of 1500 Nansen casts with observations of temperature, salinity and dissolved oxygen were carried out from June 1971 to October 1975 in seven sections, as indicated in Fig. 1. During 1971 and 1972, salinity was determined by the classical titration method. However, from 1973 onward, an Auto-lab salinometer was used. A quality analysis of the temperature and salinity data in a T–S diagram revealed an excessive scatter in the salinity determination done by titration in 1971 and 1972. These data were therefore rejected in most of the present analysis. The Winkler method was used for oxygen determination.

Plankton was sampled by oblique Bongo net tows. BCF Bongo–20 nets (20 cm diameter) with 0.5 mm mesh size and calibrated flow meters were used.




Fig. 1. Standard sections.

On the shelf the vertical range of sampling was from near the bottom to the surface and at offshore stations between the top of the thermocline and the surface (ANON. 1974, GEORGE 1976).

Assessments of the fish resources were based on acoustic surveys of the area with course tracks as indicated in Fig. 10. In the application of this method, calibrated echo sounders were used to record the fish, and the density of the resources was obtained by echo integration as described, for example, by JOHANNESON and MITSON (1983). Trawl hauls at the bottom or in mid-water were made for biological sampling and identification of echo recordings.

WATER MASSES

RAMAMIRTHAM and PATIL (1965) and DARBYSHIRE (1967) have defined three water masses in the area. The T-S diagram in Fig. 2 for the 1974 Kasaragod section confirms the presence of these water masses. Indian Ocean Equatorial Water is found at temperatures below 17°C, and its boundary is associated with a minimum salinity of 34.9‰. The Arabian Sea Water is defined by temperatures between 17 and 26°C, the high temperature being associated with a maximum in salinity up to about 35.5‰ (DARBYSHIRE 1967). According to Fig. 2. Arabian Sea Water attains temperatures up to 28°C with associated salinities close to 36‰. Equatorial Surface Water is characterized by a small temperature range between 26 and 30°C, and a wide salinity range between 30 and 34‰. In addition, a brackish water mass associated with runoff, of salinity less than 30‰ is present in Fig. 2. The T-S diagrams for the Kasaragod section for the other years show the same pattern.

A good correlation has been established between the depth of the thermocline and oxycline (Fig. 4). This has also been observed by other investigators, for example, BANSE (1968). The T-O₂ diagram for the Kasaragod section during the four-year period is plotted in Fig. 3. It shows that below the thermocline, which during the SW monsoon terminates at approximately 21–22°C, the majority of the observations of dissolved oxygen concentration are below 1 ml O₂/l. Similar diagrams have been plotted for the other sections, establishing the same relationship for the entire shelf area.




Fig. 2. Temperature-salinity diagram in the Kasaragod section during 1974. ESW: Equatorial Surface Water, ASW: Arabian Sea Water, IOE: Indian Ocean Equatorial Water.




Fig. 3. Temperature-oxygen diagram in the Kasaragod section for the period 1971–1975.

SEASONAL VARIATIONS

A study of all the sections shows that the seasonal variations reported by previous authors are very dominant. Typical conditions on the shelf are, for example, described by the Kasaragod section during 1974. (Fig. 4). The density field in January indicates weak currents, but the salinity in the upper 50 m is below 33‰. This is the lowest salinity of all sections in the year and indicates presence of Equatorial Surface Water. As described for instance by WYRTKI (1973), a branch of the NE monsoon circulation flows north along the west coast from November to January, carrying this low-salinity water which originates in the Bay of Bengal. The upper-layer salinity maximum is at the 150 m level, which is typical for the tropical oceans, and the water on the shelf is well aerated.

The situation in March demonstrates a shoreward shoaling of the isopycnals over the shelf, indicating a southward flow, but in the offshore part of the section there are clearer indications in the density field of a north-flowing current than in January. The salinity in the upper 50 m layer has, however, increased and suggests a retreat of the Equatorial Surface Water or an increased admixture of more saline Arabian Sea Water. The waters on the shelf are still aerated.




Fig. 4. Typical seasonal hydrographic fluctuations off the southwest coast as observed in the Kasaragod section during 1974.

From May through November, during the SW monsoon period, the mean flow continues southward in the upper layer. Upwelling of the Arabian Sea Water, of salinity higher than 35 ‰, is the dominant process during this period. This is illustrated by the lifting of the isotherms, with a shoaling and sharpening of the thermocline, and the penetration of the low oxygen water of less than 0.5 ml O₂/l over the entire shelf. Another feature is the lowering of the surface salinity in the nearshore region associated with the runoff during the rainy season in the SW monsoon period. This is indicated by the July and August sections.

From November to December a dramatic change in the dynamics occurs. The current reverses in the course of a month. In December it is established as a deep-reaching north-going coastal flow, thereby completing the annual cycle and again transporting low-salinity, well aerated Equatorial Surface Water northward. A rapid sinking of the Arabian Sea Water takes place during this period, suggested by the descent of the thermocline and oxycline.

SURFACE LAYER VARIATIONS

The seasonal variations in the temperature and salinity of the surface layer are illustrated in the isopleth diagrams in Fig. 5. The Cochin section was chosen for a comparison with a similar salinity diagram by DARBYSHIRE (1967, Fig. 28) for the 1958–59 period. Fig 5 shows a more repetitive pattern of the seasonal variation from year to year than could be demonstrated by the short series studied by DARBYSHIRE, who concluded that «the two years showed very different conditions and no conclusions could be drawn».

The surface salinity is low during the rainy SW monsoon period with a minimum in July–August. For example, the 33‰ isohaline extends to a maximum distance of about 60 miles off the coast in 1973. At the 10 m level the effect of runoff is drastically reduced. Generally, the surface salinity is less than 30‰ in the near-shore region from Cochin and northward, while the salinity is above 34‰ south of Quilon as illustrated in Fig. 6. Although this higher salinity in the southern region at least to some extent may be explained by the small amount of runoff in this area, the nearshore salinity front between Cochin and Quilon is suggestive of no nearshore, south-flowing current in this area. The salinity distribution in Fig. 6 suggests an offshore diversion north of the front.

The 1974 situation in Fig. 5 shows that the appearance of low-salinity Equatorial Surface Water in November or December may be rather abrupt, and the same situation shows a slower retreat in March–April. This is associated with the reversal of the current and may indicate a shorter reversal time after the SW monsoon than in March–April. In spite of considerable differences in this respect between individual years, the figure suggests that




Fig. 5. Isoleth diagrams for the Cochin section; temperature at 10 m and salinity at 0 and 10 m depths. Distance from the coast in nautical miles.

this may be a general feature, but occurring to a varying extent from year to year.

The density distribution in the Kasaragod section for December 1974 (Fig. 4) shows a wedge-formed coastal current flowing northward. The considerably lower nearshore salinities off Cochin in the same period strengthen such a distribution, typical for a freshwater-induced coastal density current, theoretically described for example by HEAPS (1972). In this case, the influence of nearshore low-salinity water may therefore be a driving force. The wind field




Fig. 6. Surface salinity in August 1974 illustrating the effect of runoff.

over the area is, of course, also of importance, and the interannual variations indicated in Fig. 5 may be associated with variations in supply, or properties of Equatorial Surface Water or monsoonal fluctuations.

The upwelling process during the SW monsoon is not well indicated in the surface salinity diagram of Fig. 5, but is well documented in the temperature pattern. The nearshore temperature is reduced to less than 23°C , while the offshore temperature at the same time is several degrees higher. A similar pattern develops in the oxygen values (not shown in the figure), where the concentrations are less than $0.5 \text{ ml O}_2/\text{l}$ near the coast.

UPWELLING

Upwelling was demonstrated by the shoreward lifting of the isolines in the description of the Kasaragod section during 1974. The other sections along the coast, as indicated by the August situation for 1974 (Fig. 7), show that these features are most prominent in the Cochin and Kasaragod sections. Slightly less upwelling is observed in the Karwar and Ratnagiri sections, while a significant change occurs between Cochin and Quilon, where only the outer part of the shelf is covered with water of less than 1 ml O_2/l . This develops further off Cape Comorin and Tuticorin where the same isoline retreats beyond the edge of the shelf.

The time variability of the upwelling along the coast is illustrated in Fig. 8A, where the intersection between 1 ml O_2/l isoline and the bottom has been plotted for the Karwar, Kasaragod, Cochin and Quilon sections. Generally, the intersection is at its greatest depth from October or November to March or April. Then the uplifting starts and reaches its most shallow position within the period from July to October. A plot of temperature, up to January 1973, near the bottom at 50 m depth in the sections (ANON. 1973) indicated a northward delay of the occurrence of the upwelling. This is not evident in Fig. 8A, and it is concluded that upwelling on the entire shelf starts within a period of 1–2 months. The sinking or the retreat of the 1ml O_2/l isoline from the shelf indicates the decay of the south-flowing current and the onset of the north-flowing current. In 1973 and 1974 this took place first in the southern




Fig. 7. Upwelling along the coast in August 1974 illustrated by the temperature and oxygen (O_2 ml/l) sections.

area off Quilon and Cochin, 1–2 months earlier than the northern area, while in 1972 it occurred at approximately the same time over the entire shelf area. The «rate» of upwelling can be qualitatively indicated by the rise of the isolines for the different parameters, which for this coastal zone is on the order of 1.5 m/day.

The driving forces of the upwelling are not evident. Several investigators have discussed the matter, and different viewpoints have been formulated. DARBYSHIRE (1967) concluded that «wind upwelling can not take place as the prevailing winds are onshore from west or southwest in the SW monsoon period» and claimed that the upwelling or the upward slope of the density surfaces toward the coast was consistent with the observed southerly current. BANSE (1968), however, mentioned that upwelling started with the onset of the SW monsoon. The interpretation of the present data suggests that the wind becomes the important driving force when the influence of the low-salinity Equatorial Surface Water decays in February. The direction of prevailing winds in February and March is mainly between northwest and northeast, of average Beaufort 3 (ANON. 1979). This gives rise to the reversal of the current system well in advance of the SW monsoon period. During the SW monsoon period the prevailing wind direction is from west (ANON. 1979, RAMAGE 1969) and is consequently not in favour of intensifying a wind-driven upwelling. Due to higher wind force during this period, the south-flowing current is, however, speeded up and the associated tilt of the isolines is increased. It seems, therefore, likely that the upwelling is not only associated with the local wind, but also with the more large scale monsoonal conditions which drive the anticyclonic Arabian Sea monsoon gyre.

OCEANOGRAPHIC INFLUENCE ON THE BIOMASS

In a discussion of the relationship between oceanographic factors and distribution and abundance of biomass, upwelling stands out as a dominant feature. Upwelling boosts the phytoplankton production in the area. This is reflected in the zooplankton production (ANON. 1976 c). The general trend of zooplankton biomass shown in Fig. 8B follows the pattern of the upwelling in Fig. 8A. On higher levels in the food chain the present observations do not reveal such close correlation with environmental components, but nevertheless upwelling is indirectly of importance since the feeding and growth of the fish population depend on the primary and secondary production.

The behaviour and distribution of some fish species are also directly restricted by the upwelling onto the shelf of cool, oxygen-deficient water. It was well known among local fishermen that schools of mackerel (*Rastrelliger kanagurta*) and sardines (*Sardinella spp.*) were often visible on the surface just before and after the SW monsoon. The investigations of the Pelagic Fishery




Fig. 8. A: Depth of the intersection between the 1 ml O₂/l isoline and the bottom.
 B: Zooplankton biomass in ml plankton per m³ of water filtered.
 1: Karwar section, 2: Kasaragod section, 3: Cochin section, 4: Quilon section.

Project revealed that a stock of these species was present in the area also during the monsoon months, but even though the schools were not visible on the surface during the rougher weather conditions of the monsoon, they occurred in the aerated surface layer above the oxycline (ANON. 1976 b).

A stock of anchovy (*Stolephorus spp.*) showed migrations clearly phased with the monsoon (ANON. 1976 a). As depicted in Fig. 9, the anchovy stock was dispersed along the southwest coast during the NE monsoon, mainly north of Quilon. A southward migration started around April–May, and in August through October almost all the stock was concentrated in the Gulf of Manaar to the east of Cape Comorin as shown in Fig. 10. The figure also shows typical survey tracks on which the distributional mapping was based. After the SW monsoon the anchovy again dispersed along the west coast of Quilon.

As previously described by, for example, GEORGE (1958) and BANSE (1959), some demersal species disappear from the oxygen-deficient water on the shelf during the SW monsoon. The Pelagic Fishery Project observed the demersal species as part of a group consisting of a considerable variety of species. Due to the different migration patterns of various species, it was difficult to draw any conclusive picture of their overall migratory behaviour. Nevertheless, the general trend was that demersal fish occurred in abundance on the northern shelf during post-monsoon months while they were concentrated on the southern shelf area during the SW monsoon months.




Fig. 9 Distribution of anchovy (*Stolephorus spp.*) in March-April 1974. 1: very scattered, 2: scattered, 3: dense.




Fig. 10. Distribution of the anchovy in September-October 1974 at the end of the SW monsoon. 1: very scattered, 2: scattered, 3: dense.

SUMMARY AND CONCLUSIONS

The seasonal variations in the oceanographic conditions are repetitive from year to year. Generally, a shoreward lifting of isopleths starts in March and continues through the SW monsoon, reaching a maximum in August or September. This upwelling causes the penetration of low-oxygen water over the entire shelf north of Quilon, while south of Quilon it is less significant. The influence of runoff during the latter part of the SW monsoon can be traced up to 60 nautical miles offshore.

During the early part of the NE monsoon the current reverses. Due to the influence of low-salinity Equatorial Surface Water, this has the typical features of a coastal density current, opposing northerly winds of Beaufort 2-3. The characteristic time scale for reversing the circulation on the shelf is of the order of 1-2 months.

Some biological events clearly reflect the seasonal environmental variations. A peak in the plankton concentration is associated with upwelling. Schooling pelagic fish concentrate in the surface layer due to the oxygen deficiency below the oxycline, and some species disappear from the northern part of the shelf during the SW monsoon months.

ACKNOWLEDGEMENTS

The UNDP/FAO Pelagic Fishery Project (IND.69/593) was jointly sponsored by the United Nations Development Programme and the Government of India. The project was subcontracted to the Norwegian Agency for International Development (NORAD) and the Institute of Marine Research, Bergen. The authors acknowledge the extensive sampling work and data processing by the project staff.

REFERENCES

- ANON. 1973. Progress Report No. 3, Hydrographic investigations - June 1971-January 1973. *UNDP/FAO Pelagic Fishery Project (IND/69/593)*, Bergen/Cochin, 1973: 1-13, + 1 Table + 19 Figures.
- ANON. 1974. Progress Report no. 7, Plankton - fish eggs and larvae studies. *UNDP/FAO Pelagic Fishery Project (IND/69/593)*, Bergen/Cochin, 1974: 1-21, + 3 Tables + 23 Figures.
- ANON. 1976. Progress Report No. 16. Physical oceanography of the south-west coast of India based on the investigations of the UNDP/FAO Pelagic Fishery Project. *UNDP/FAO Pelagic Fishery Project (IND/69/593)*, Bergen/Cochin, 1976: 1-39.
- ANON. 1976 a. Progress Report No. 12, Survey results 1973/74. *UNDP/FAO Pelagic Fishery Project (IND/69/593)*, Bergen/Cochin, 1976: 1-116.
- ANON. 1976 b. Progress Report No. 13, Survey results 1974/75. *UNDP/FAO Pelagic Fishery Project (IND/69/593)*, Bergen/Cochin, 1976: 1-107.
- ANON. 1976 c. Progress Report No. 17, Plankton - fish eggs and larvae studies. *UNDP/FAO Pelagic Fishery Project (IND/69/593)*, Bergen/Cochin, 1976: 1-27.

- ANON. 1979. Atlas of pilot charts, Indian Ocean, First Edn, 1979. *Defence Mapping Agency, Hydrographic/Topographic Center. Washington DC. Pub., 109*: 12 maps.
- BANSE, K. 1959. On upwelling and bottom-trawling off the southwest coast of India. *J. Mar. biol. Ass. India*, 1 (1): 33-49.
- BANSE, K. 1968. Hydrography of the Arabian Sea shelf of India and Pakistan and effects on demersal fishes. *Deep-Sea Res.*, 15: 45-79.
- DARBYSHIRE, M. 1967. The surface waters off the coast of Kerala, south-west India. *Deep-Sea Res.*, 14: 296-320.
- GEORGE, P.C. 1958. Sole fisheries. P. 51-54 in Jones, S. ed. *Fisheries of the West Coast of India, Mandapam Camp, India*.
- GEORGE, K.C. 1976. Ichthyoplankton surveys along the south West coast of India by the UNDP/FAO Pelagic Fishery Project, Cochin. 1976 (C.5/11.) *Abstracts of papers. Joint Oceanographic Assembly, Edinburg, 1976. FAO*:132 pp.
- HEAPS, N.S. 1972. Estimation of density currents in the Liverpool Bay area of the Irish Sea. *Geophys. J.R. astr. Soc.*, 30: 415-432.
- JOHANNESSON, K.A. and MITSON, R.B. 1983. Fisheries acoustics, a practical manual for aquatic biomass estimation. *FAO Fish. Tech. Pap.* 240: 1-249.
- RAMAGE, C.S. 1969. Indian Ocean surface meteorology. *Oceanogr. Mar. Biol. ann. Rev.*, 7: 11-30.
- RAMAMIRTHAM, C.P. and PATIL, M.R. 1965. Hydrography of the west coast of India during the pre-monsoon period of the year 1962 - Part 2: In and offshore waters of the Konkan and Malabar coasts. *J. Mar. biol. Ass. India*, 7: 150-168.
- RAMA SASTRY, A.A. and MYRLAND, P. 1960. Distribution of temperature, salinity and density in the Arabian Sea along the south Malabar coast (South India) during the post-monsoon season. *Indian J. Fish.*, 6(2): 223-255.
- WYRTKI, K. 1973. Physical oceanography of the Indian Ocean. P. 18-36 in ZEITZHEL, B. ed. *Biology of the Indian Ocean*. Springer-Verlag, Berlin.

Received 14 March 1986

Printed 16. February 1987