

# A new maximum age observed within the family Labridae, ballan wrasse *Labrus bergylta*

Calum J. Pritchard<sup>1,2</sup>  | Marthe M. R. Stendal<sup>3,4</sup>  | Jim Treasurer<sup>5</sup>  |  
Kim T. Halvorsen<sup>3</sup>  | Anne B. Skiftesvik<sup>3</sup> 

<sup>1</sup>School of Biological Sciences, University of Aberdeen, King's College, Aberdeen, UK

<sup>2</sup>Scottish Government's Marine Directorate, Marine Laboratory, Aberdeen, UK

<sup>3</sup>Ecosystem Acoustics Group, Institute of Marine Research, Flødevigen Research Station, His, Norway

<sup>4</sup>Department of Biological Sciences, University of Bergen, Bergen, Norway

<sup>5</sup>FAI Farms, Oxford, UK

## Correspondence

Calum J. Pritchard, School of Biological Sciences, University of Aberdeen, King's College, Aberdeen AB24 3FX, Scotland.  
Email: [c.pritchard.21@abdn.ac.uk](mailto:c.pritchard.21@abdn.ac.uk)

## Funding information

Scottish Salmon Education and Research Foundation (SSERF); Scottish Government's Marine Directorate; Norwegian Research Council, Grant/Award Number: 335887; Institute of Marine Research, Grant/Award Numbers: 15638, 15626-02

## Abstract

Ballan wrasse *Labrus bergylta* is the largest species of wrasse inhabiting European waters and one of the longest-living species within the family Labridae. A large specimen was caught off the coast of Skjerjehamn, western Norway (total length = 410 mm; weight = 1274 g). The age of the specimen was determined to be 34 years old based on the analysis of its opercula bones. This specimen establishes a new maximum age for ballan wrasse, 5 years older than the previously observed maximum age.

## KEYWORDS

ballan wrasse, *Labrus bergylta*, life history, longevity, maximum age, opercula


Ballan wrasse *Labrus bergylta* Ascanius 1767 is the largest species of wrasse (family: Labridae) inhabiting European waters, with a maximum observed length of 660 mm (Villegas-Ríos et al., 2021). Ballan wrasse also have the greatest longevity of European Labrids, with a maximum observed age of 29 years, as determined for a male specimen caught on the Isle of Man (Dipper et al., 1977). This species is a protogynous sequential hermaphrodite, of which the oldest observed female was 25 years (Dipper et al., 1977). In northern Europe, ballan wrasse are a relatively newly commercially exploited species after the introduction of wrasse as cleaner fish on salmonid farms (Bjordal, 1988). More than 1 million wild ballan wrasse were landed annually in Norway between 2013 and 2016 (Blanco Gonzalez & de Boer, 2017). Prior to this, ballan wrasse have been exploited as lobster bait in some parts of Norway; however, the extent of this fishery is unknown due to it being poorly documented within the literature. Ballan wrasse are considered to be

vulnerable to overexploitation due to their life-history strategy as a protogynous sequential hermaphrodite, where populations have naturally female-skewed sex ratios. Fishery-independent research has estimated that males, who provide obligate parental care, make up as little as 8%–12% of the population (Dipper et al., 1977; Mucientes et al., 2019; Quignard, 1966). To our knowledge, a quantitative stock assessment has not been performed for any population of ballan wrasse in northern Europe to date. Therefore, fisheries managers rely on valid life-history information to implement appropriate management strategies. As a measure to avoid depletion of males and age- and size-truncation in exploited ballan wrasse populations, a slot size limit, with a minimum landing size of 22 cm and a maximum landing size of 28 cm, was recently implemented in Norway (pre-2022 season). Slot size limits have also been implemented for wrasse species in Sweden, Scotland, and some of the fishing districts in England (Halvorsen et al., 2021).

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2024 The Author(s). *Journal of Fish Biology* published by John Wiley & Sons Ltd on behalf of Fisheries Society of the British Isles.

**FIGURE 1** (a) Whole specimen; (b) backcalculated length-at-age; (c) magnified opercula, including marks on the annuli used to age the sample; and (d) unmagnified opercula.



A large specimen was caught in a scientific survey by gillnet in Skjerjehamn, Norway (60.9515° N, 4.966° E; depth between 3 and 30 m), on October 25, 2021 (Figure 1a; Otterå et al., 2024). The specimen was determined to be an intermediate morphotype, neither plain nor spotted, using Seljestad et al. (2020) as a guide. The specimen was frozen and processed at a later date. The specimen measured 410 mm (total length, accuracy = 1 mm) and weighed 1274 g (accuracy = 1 g). The sex of the specimen was identified as male through macroscopic inspection of the gonads. Age was determined by reading the number of annuli present within the opercula bones. Studies on the age and growth of marine fish have mainly used otoliths for age determination and, in some cases, scales. However, the opercula bone has been used previously in fishes, such as perch *Perca fluviatilis*, where scales and otoliths were less clear and reliable, reported for the first time by Nilsson (1921). Ballan wrasse otoliths are very small and often unclear, so opercula bones have been commonly used for age determination as described by Dipper et al. (1977). Both opercula bones were extracted, immersed in boiled water for 10 min, and wiped clean. Opercula were coated in a thin layer of oil and photographed using a microscope under reflected light. We assumed the birth date to be May 1 as described by Dipper et al. (1977). As the specimen was captured in October, the opercula edge was considered to be plus-growth instead of a further annulus. Length-at-age was backcalculated using the equation:

$$L_i - a = \left(\frac{R_i}{R}\right)(L_C - a)$$

where  $L_i$  is length at age,  $i$ ;  $R_i$  is the radius of annulus,  $i$ ;  $R$  is the radius of the opercula;  $L_C$  is the length at capture; and  $a$  is a correction factor

obtained as the intercept from a regression of fish length with opercula radius. This is a modification of the direct proportionality formula used by Lee (1912) to correct for allometric growth and to account for opercula size at age zero (32.5 mm; Dipper et al., 1977). The age of the specimen was estimated as 34 years, based on readings of both opercula bones by three independent analysts (C.P., J.T., and M.S.). Backcalculated length-at-age for the Skjerjehamn specimen is shown in Figure 1b with comparison to ballan wrasse from the Isle of Man (Dipper et al., 1977) and Scotland (Treasurer, 1994).

Backcalculated lengths-at-age suggest that the first annulus was correctly identified as  $L_1$  and is highly comparable with age-1 fish from Scotland and the Isle of Man. All ballan wrasse show a similar growth pattern up to 13 years, after which the Skjerjehamn specimen achieves a smaller length-at-age. The Skjerjehamn specimen is 20 mm shorter than that of the previous oldest observed ballan wrasse, despite being 5 years older (Dipper et al., 1977). Plain morphotype individuals have been documented to have slower growth rates and attain smaller sizes-at-age than spotted morphotype individuals (Villegas-Ríos et al., 2013), though Dipper et al. did not document the morphotype of their specimen. Whether growth rates and longevity differ among plain, intermediate, and spotty ballan wrasse in Scandinavian populations has not yet been investigated, but a recent study did not detect any genetic differences among morphotypes in Scandinavia, whereas they are clearly different in the Spanish population (Seljestad et al., 2020).

The time elapsed between the previously reported maximum age and the present paper, 47 years, initially suggests that the former was a valid representation of maximum age for this species. However, we argue that the new maximum age presented in this paper, 34 years,

may remain to be an underestimate of this species' longevity as wrasse are rarely sampled and aged outside of the fishery. The live wrasse fishery directly targets the younger demographic of ballan wrasse through the use of size-selective fishing gears and in compliance with maximum landing sizes (Halvorsen et al., 2019). Targeting the youngest demographic of ballan wrasse may also be indirect as the live wrasse fishery is generally restricted to shallow water to reduce the risk of barotrauma, but research suggests that larger ballan wrasse occupy deeper depths (Halvorsen et al., 2020). Therefore, fishery-dependent surveys are biased toward sampling young fish. Although the absence of interactions between large fish and the fishery is a positive indicator for the sustainability of this fishery, the absence of large specimens available for scientific research propagates ballan wrasse as a data-limited species. Therefore, future surveys should operate independent of the live wrasse fishery to sample the entire demographic of ballan wrasse and establish a valid representation of their population dynamics (e.g., growth rates, mortality rates, and recruitment). However, caution should be taken to ensure that sampling of old males is kept to a minimum to preserve natural sex ratios. Although anecdotal, this observation may indicate a healthy ballan wrasse population on the coast of Skjerjehamn due to the lack of age truncation. The maximum age presented in this paper contributes to a greater understanding on the longevity of ballan wrasse and updates the life-history information available for this species. This finding is particularly important within the context of fisheries management, as sampling the eldest demographic is crucial for producing valid and representative growth models and estimating natural mortality rates.

#### AUTHOR CONTRIBUTIONS

Kim T. Halvorsen and Anne B. Skiftesvik arranged the sampling. Calum J. Pritchard and Marthe M. R. Stendal performed the dissection. Marthe M. R. Stendal prepared the opercula for aging. Calum J. Pritchard, Jim Treasurer, and Marthe M. R. Stendal examined the opercula and determined age. Calum J. Pritchard wrote the manuscript. All the authors contributed to improving the initial draft.

#### ACKNOWLEDGMENTS

The authors would like to thank Håkon Otterå and Kjell Nedreaas for organizing the gillnet survey and Torkel Larsen for his assistance in the laboratory.

#### FUNDING INFORMATION

The study was funded by the Scottish Salmon Education and Research Foundation (SSERF), Scottish Government's Marine Directorate, and the Norwegian Research Council through the project "Understanding human-induced trophic cascades in coastal ecosystems" (project # 335887) and the Institute of Marine Research (project # 15638 and 15626-02).

#### ORCID

Calum J. Pritchard <https://orcid.org/0000-0001-7093-6785>

Marthe M. R. Stendal <https://orcid.org/0009-0002-7725-6453>

Jim Treasurer <https://orcid.org/0000-0002-3447-9449>

Kim T. Halvorsen <https://orcid.org/0000-0001-6857-2492>

Anne B. Skiftesvik <https://orcid.org/0000-0002-7754-5661>

#### REFERENCES

- Bjoridal, Å. (1988). *Cleaning symbiosis between wrasse (Labridae) and lice infested salmon (Salmo salar) in mariculture* (Vol. 188, pp. 1–8). International Council for the Exploration of the Sea Mariculture Committee.
- Blanco Gonzalez, E., & de Boer, F. (2017). The development of the Norwegian wrasse fishery and the use of wrasses as cleaner fish in the salmon aquaculture industry. *Fisheries Science*, 83, 661–670.
- Dipper, F. A., Bridges, C. R., & Menz, A. (1977). Age, growth and feeding in the ballan wrasse *Labrus bergylta* Ascanius 1767. *Journal of Fish Biology*, 11, 105–120.
- Halvorsen, K., Skiftesvik, A. B., & Jørgensen, T. (2019). Kunnskapsstøtte- anbefaling om redusert inngangsstørrelse i teiner i fisket etter leppefisk. *Havforskningsinstituttet Rapport*, 20-2019, 1–8.
- Halvorsen, K. T., Skiftesvik, A. B., Durif, C. M., Faust, E., Wennhage, H., André, C., Linnemann Rønfeldt, J., Moller, P. R., Carl, H., & Jørgensen, T. (2021). Towards a sustainable fishery and use of cleaner fish in salmonid aquaculture. In *TemaNord*, (2021: 545). Nordic Council of Ministers.
- Halvorsen, K. T., Sjørdalen, T. K., Larsen, T., Browman, H. I., Rafoss, T., Albretsen, J., & Skiftesvik, A. B. (2020). Mind the depth: The vertical dimension of a small-scale coastal fishery shapes selection on species, size, and sex in wrasses. *Marine and Coastal Fisheries*, 12, 404–422.
- Lee, R. M. (1912). An investigation into the methods of growth determination in fishes. *Conseil Permanent International l'Exploration de la Mer Publacion de Cironstance*, 63, 3–34.
- Mucientes, G., Irisarri, J., & Villegas-Ríos, D. (2019). Interannual fine-scale site fidelity of male ballan wrasse *Labrus bergylta* revealed by photo-identification and tagging. *Journal of Fish Biology*, 95, 1151–1155.
- Nilsson, D. (1921). *Nagra insjöfiskars alder och tillvaxt i Bottniska Viken och Malaren. Medd. Landtbr-Styr* (p. 231) [in Swedish]. Tiden.
- Otterå, H., Bjelland, O., Hinriksson, J., Næss, H., & Nedreaas, K. (2024). *Prøvefiske med trollgarn og dleruser på grunt vatn* (Vol. 24). Survey Report – Institute of Marine Research. [in Norwegian].
- Quignard, J. P. (1966). *Recherches sur les Labridae (poison téléostéen perciformes) des côtes européennes: systématique et biologie* (Vol. 5, pp. 7–248). Naturalia Monspel.
- Seljestad, G. W., Quintela, M., Faust, E., Halvorsen, K. T., Besnier, F., Jansson, E., Dahle, G., Knutsen, H., André, C., Folkvord, A., & Glover, K. A. (2020). 'A cleaner break': Genetic divergence between geographic groups and sympatric phenotypes revealed in Ballan wrasse (*Labrus bergylta*). *Ecology and Evolution*, 10, 6120–6135.
- Treasurer, J. W. (1994). The distribution, age and growth of wrasse (Labridae) in inshore waters of west Scotland. *Journal of Fish Biology*, 44, 905–918.
- Villegas-Ríos, D., Alonso-Fernández, A., Fabeiro, M., Bañón, R., & Saborido-Rey, F. (2013). Demographic variation between colour patterns in a temperate protogynous hermaphrodite, the ballan wrasse *Labrus bergylta*. *PLoS One*, 8, 71591.
- Villegas-Ríos, D., Bañón, R., & Fabeiro, M. (2021). A new maximum length of *Labrus bergylta* (Labriformes: Labridae) with notes on age determination for the species. *Cybius*, 45, 239–242.

**How to cite this article:** Pritchard, C. J., Stendal, M. M. R., Treasurer, J., Halvorsen, K. T., & Skiftesvik, A. B. (2024). A new maximum age observed within the family Labridae, ballan wrasse *Labrus bergylta*. *Journal of Fish Biology*, 105(4), 1354–1356. <https://doi.org/10.1111/jfb.15886>